

Table 4: HIV testing uptake by modes in different types of sexual relationship among reachable sexual contacts of newly diagnosed HIV positive MSM in June 2014 through June 2016, in Hangzhou and Ningbo Cities, China.

Type(s) of sexual relationship with sexual contacts			Modes of partner tracing and testing package	Tested	Not tested	χ^2	<i>p</i> -value
Stable, non-commercial same-sex relationship/ opposite-sex spouse			CHCT	120(88.2)	16(11.8)	2.827	0.419
			IDAPN	56(94.9)	3(5.1)		
			HIVST	10(90.9)	1(9.1)		
			Patient Referral	80(93.0)	6(7.0)		
Casual, non-commercial same-sex relationship			CHCT	39(68.4)	18(31.6)	-	<0.001

Type(s) of sexual relationship with sexual contacts	Modes of partner tracing and testing package	Tested	Not tested	χ^2	<i>p</i> -value
Commercial same-sex relationship / Unmarried opposite-sex relationship/ Others	IDAPN	89(93.7)	6(6.3)	-	0.392
	HIVST	5(100.0)	0(0.0)		
	Patient Referral	67(87.0)	10(13.0)		
	CHCT	3(75.0)	1(33.3)		
	IDAPN	1(100.0)	0(0.0)		
	HIVST	-	-		
	Patient Referral	6(75.0)	2(25.0)		