

-----原始邮件-----

发件人: gb-admin@ncbi.nlm.nih.gov
发送时间: 2014-11-13 21:34:10 (星期四)
收件人: xykong@scsio.ac.cn
抄送:
主题: GenBank KP134336-KP134337

Dear GenBank Submitter:

Thank you for your direct submission of sequence data to GenBank. We have provided GenBank accession numbers for your nucleotide sequences:

Psettina iijimae.sqn JP57 KP134336
Arnoglossus tenuis.sqn XYSP58 KP134337

The GenBank accession numbers should appear in any publication that reports or discusses these data, as it gives the community a unique label with which they may retrieve your data from our on-line servers. You may prepare and submit your manuscript before your accessions are released in GenBank.

Submissions are not automatically deposited into GenBank after being accessioned. Each sequence record is individually examined and processed by the GenBank annotation staff to ensure that it is free of errors or problems.

You have requested that your data are to be held confidential until:

Dec 31, 2019

They will not be released to the public database until this date, or until the data or accession numbers appear in print, whichever is first.

Since the flatfile record is a display format only and is not an editable format of the data, do not make changes directly to a flatfile. For complete information about different methods to update a sequence record, see: <http://www.ncbi.nlm.nih.gov/Genbank/update.html>

Any inquiries about your submission should be sent to gb-admin@ncbi.nlm.nih.gov

For more information about the submission process or the available submission tools, please contact GenBank User Support at info@ncbi.nlm.nih.gov.

Please reply using the original subject line.
This will allow for faster processing of your correspondence.

Sincerely,

Beverly Underwood
Contractor

The GenBank Direct Submission Staff
Bethesda, Maryland USA

-----原始邮件-----

发件人: gb-admin@ncbi.nlm.nih.gov
发送时间: 2018-12-06 22:53:49 (星期四)
收件人: xykong@scsio.ac.cn
抄送:
主题: GenBank MK256952

Dear GenBank Submitter:

Thank you for your direct submission of sequence data to GenBank. We have provided a GenBank accession number for your nucleotide sequence:

A.intermedius.sqn A.intermedius MK256952

The GenBank accession number should appear in any publication that reports or discusses these data, as it gives the community a unique label with which they may retrieve your data from our on-line servers. You may prepare and submit your manuscript before your accession is released in GenBank.

Submissions are not automatically deposited into GenBank after being accessioned. Each sequence record is individually examined and processed by the GenBank annotation staff to ensure that it is free of errors or problems.

You have requested that your data are to be held confidential until:

Nov 30, 2024

They will not be released to the public database until this date, or until the data or accession numbers appear in print, whichever is first.

Since the flatfile record is a display format only and is not an editable format of the data, do not make changes directly to a flatfile. For complete information about different methods to update a sequence record, see: <https://www.ncbi.nlm.nih.gov/Genbank/update.html>

Any inquiries about your submission should be sent to gb-admin@ncbi.nlm.nih.gov

For more information about the submission process or the available submission tools, please contact GenBank User Support at info@ncbi.nlm.nih.gov.

Please reply using the current Subject line.

Sincerely,

Anjanette Johnston, PhD

The GenBank Direct Submission Staff
Bethesda, Maryland USA

gb-admin@ncbi.nlm.nih.gov (for updates/replies to GenBank entries)
info@ncbi.nlm.nih.gov (for general questions regarding GenBank)

-----原始邮件-----

发件人: gb-admin@ncbi.nlm.nih.gov
发送时间: 2019-04-10 01:13:36 (星期三)
收件人: 976526480@qq.com, xykong@scsio.ac.cn
抄送:
主题: GenBank MK770643

Dear GenBank Submitter:

Thank you for your direct submission of sequence data to GenBank. We have provided a GenBank accession number for your nucleotide sequence:

BankIt2211545 Seq1 MK770643

The GenBank accession number should appear in any publication that reports or discusses these data, as it gives the community a unique label with which they may retrieve your data from our on-line servers. You may prepare and submit your manuscript before your accession is released in GenBank.

Submissions are not automatically deposited into GenBank after being accessioned. Each sequence record is individually examined and processed by the GenBank annotation staff to ensure that it is free of errors or problems.

You have requested that your data are to be held confidential until:

Apr 9, 2024

They will not be released to the public database until this date, or until the data or accession numbers appear in print, whichever is first.

Since the flatfile record is a display format only and is not an editable format of the data, do not make changes directly to a flatfile. For complete information about different methods to update a sequence record, see: <https://www.ncbi.nlm.nih.gov/Genbank/update.html>

Any inquiries about your submission should be sent to gb-admin@ncbi.nlm.nih.gov

For more information about the submission process or the available submission tools, please contact GenBank User Support at info@ncbi.nlm.nih.gov.

Please reply using the current Subject line.

Sincerely,

Anjanette Johnston, PhD

The GenBank Direct Submission Staff
Bethesda, Maryland USA

gb-admin@ncbi.nlm.nih.gov (for updates/replies to GenBank entries)
info@ncbi.nlm.nih.gov (for general questions regarding GenBank)

Case CAS-405542-L0K8C8 - National Library of Medicine Customer Service confirmation TRACKING:000435000239472

📧 📅 📄 📧 发起会议

精简信息 ^

发件人: NLM Supp... <nlm-supp...@nlm.nih.gov>

时 间: 2019年08月02日 16:06:17 (星期五)

收件人: 罗海容 <luohairong14@mails.ucas.edu.cn>

Thank you for contacting the U.S. National Library of Medicine (NLM). The details of your message are listed below. You can expect to hear from us within 2 to 5 business days. For additional information about NLM products and services, please visit <https://support.nlm.nih.gov>.

Case Information:

Case #: CAS-405542-L0K8C8

Summary: Questions for searching the submitted mitogenomic sequences

Details: Dear GenBank Submission Staff:

GenBank accession number of two complete mitochondrial DNA sequences, KP134336 (Psettina iijimae) and KP134337 (Arnoglossus tenuis), have appeared in the published paper, Concerted Evolution of Duplicate Control Regions in the Mitochondria of Species of the Flatfish Family Bothidae (Teleostei: Pleuronectiformes).

But when KP134337 and KP134336 are entered into the database, it is made clear that "This record has not yet been released. Please contact info@ncbi.nlm.nih.gov with the publication details if the accession has been published."

As the submitter of these two complete mitogenomes, we are wondering could you assist us release the sequence of KP134336 and KP134337.

In addition, other two complete mitochondrial DNA sequences, MK770643 (Grammatobothus polyophthalmus) and MK256952 (Asterorhombus intermedius) were submitted by us before Apr. 2019. However, when entered these two accession numbers into the database, it is made clear that "The following term was not found in Nucleotide: No items found."

We are wondering what the reason of this situation, and we

also want to make sure will these two sequences be release once the corresponding accessions have been published?

Thanks for your consideration and assistance.

If you have any question, please feel free to contact me or Dr. Xiaoyu Kong.

Sincerely,
Hairong Luo

LMB, The South China Sea Institute of Oceanology
No.164, Xingangxi Rd,
Haizhu District
Guangzhou 510301

Case Created: 8/2/2019 8:06 AM

RESEARCH ARTICLE

Concerted Evolution of Duplicate Control Regions in the Mitochondria of Species of the Flatfish Family Bothidae (Teleostei: Pleuronectiformes)

Dong-He Li^{1,2,3}, Wei Shi^{1,2}*, Thomas A. Munroe⁴, Li Gong^{1,2,3}, Xiao-Yu Kong¹*

1 Key Laboratory of Tropical Marine Bio-resources and Ecology and Guangdong Provincial Key Laboratory of Applied Marine Biology, South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou, 510301, China, **2** South China Sea Bio-Resource Exploitation and Utilization Collaborative Innovation Center, Guangzhou, 510301, China, **3** University of Chinese Academy of Sciences, Beijing, 100049, China, **4** National Systematics Laboratory NMFS/NOAA, Post Office Box 37012, Smithsonian Institution NHB, WC 60, MRC-153, Washington, D.C., 20013–7012, United States of America

* These authors contributed equally to this work.

* shiwei@scsio.ac.cn (WS); xykong@scsio.ac.cn (XYK)



OPEN ACCESS

Citation: Li D-H, Shi W, Munroe TA, Gong L, Kong X-Y (2015) Concerted Evolution of Duplicate Control Regions in the Mitochondria of Species of the Flatfish Family Bothidae (Teleostei: Pleuronectiformes). PLoS ONE 10(8): e0134580. doi:10.1371/journal.pone.0134580

Editor: Bi-Song Yue, Sichuan University, CHINA

Received: December 31, 2014

Accepted: July 12, 2015

Published: August 3, 2015

Copyright: © 2015 Li et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: The GenBank accession numbers for new complete mitogenomes sequenced in this study are KJ433561, KJ433566, KJ433567, [KP134336](#) and [KP134337](#). Tree files are available on the following webset: <http://purl.org/phylo/treebase/phyloids/study/TB2:S17532>

Funding: This work was supported by the Natural Science Foundation of China (grant numbers 31471979, 41206134, 30870283) and Science and Technology Planning Project of Guangdong Province, China (2014B030301064).

Abstract

Mitogenomes of flatfishes (Pleuronectiformes) exhibit the greatest diversity of gene rearrangements in teleostean fishes. Duplicate control regions (CRs) have been found in the mito-genomes of two flatfishes, *Samariscus latus* (Samaridae) and *Laeops lanceolata* (Bothidae), which is rare in teleosts. It has been reported that duplicate CRs have evolved in a concerted fashion in fishes and other animals, however, whether concerted evolution exists in flatfishes remains unknown. In this study, based on five newly sequenced and six previously reported mitogenomes of lefteye flounders in the Bothidae, we explored whether duplicate CRs and concerted evolution exist in these species. Results based on the present study and previous reports show that four out of eleven bothid species examined have duplicate CRs of their mitogenomes. The core regions of the duplicate CRs of mitogenomes in the same species have identical, or nearly identical, sequences when compared to each other. This pattern fits the typical characteristics of concerted evolution. Additionally, phylogenetic and ancestral state reconstruction analysis also provided evidence to support the hypothesis that duplicate CRs evolved concertedly. The core region of concerted evolution is situated at the conserved domains of the CR of the mitogenome from the termination associated sequences (TASs) to the conserved sequence blocks (CSBs). Commonly, this region is considered to regulate mitochondrial replication and transcription. Thus, we hypothesize that the cause of concerted evolution of the duplicate CRs in the mtDNAs of these four bothids may be related to some function of the conserved sequences of the CRs during mitochondrial replication and transcription. We hope our results will provide fresh insight into the molecular mechanisms related to replication and evolution of mitogenomes.

Competing Interests: The authors have declared that no competing interests exist.

Introduction

The vertebrate mitochondrial genome (mitogenome) typically codes for 37 genes, including 13 protein-coding genes, 22 transfer RNAs (tRNAs), two ribosomal RNAs (rRNAs) and one control region (CR) [1]. However, two or more CRs have been found in some vertebrate mitogenomes, including those of birds [2–4], turtles [5–7], snakes [8–10] and fishes [11–13]. Moreover, there is a special phenomenon in mitogenomes with duplicate CRs, where sequences of the duplicate CRs are extremely similar. This has been proposed to have evolved in a concerted fashion, a situation which has been found in birds, snakes, fishes, sea cucumbers, sea firefly and ticks [3, 4, 8, 9, 12, 14–18]. Based on Liao's opinion (1999), the concept of concerted evolution of duplicate CRs is a peculiar evolutionary phenomenon of the two CRs in mitogenomes, which may lead to homogenization of duplicate CRs within one species [19]. Shao *et al.* [17] further defined concerted evolution of CRs as including two main points: (1) the sequences of CR1 and CR2 of one species are highly similar; and (2) they are more similar to each other in the same species than either is to its namesake in the other species.

Molecular phylogenetic studies have made significant contributions to further understanding the process of concerted evolution of duplicate CRs in many different species [3, 8, 9, 12, 14]. For example, Eberhard *et al.* [15] conducted a phylogenetic analysis of the duplicate CRs from 21 individuals representing four *Amazona* parrots. The result revealed that two CRs of an individual were more closely related to one another than to corresponding segments of others in the same species, except for three subspecies in *Amazona ochrocephala*. The paralogous CRs of an individual in the same subspecies did not group together, but the orthologous CRs in different individuals clustered first. This phylogenetic analysis helps to understand the different evolutionary pattern of the two CRs, one was concerted evolution of the two CRs in some species and the other was independent evolution in other subspecies.

As more mtDNAs with duplicate CRs have been sequenced, different hypotheses explaining how duplicate CRs are generated and how they have evolved in a concerted fashion have been put forward [9, 16, 18]. Most of them explain for either the generation of CRs or for the concerted evolution of duplicate CRs, just one example could explain both cases.

Two mechanisms, illegitimate recombination and dimeric mitogenome, were used to account for the generation of duplicate CRs. The illegitimate recombination involves the breakage and rejoining of participating DNA strands [20], and if the recombined DNA strands include the CR, it would generate a mitogenome with two CRs. The dimeric mitogenome is formed by two monomeric mitogenomes joining head to tail (dimerization). This mitogenome would have two CRs and two sets of mitochondrial genes [21].

Another two mechanisms have been proposed to explain how duplicate CRs maintain concerted evolution. Ogoh and Ohmiya [18] illustrated that the duplicated CR1 inserted at the location where the old CR2 was deleted during mitochondrial replication, which made the sequences of the two CRs identical or highly similar. The mechanism of gene conversion proposed by Kumazawa *et al.* [9] indicated that the sequences of two control regions can be homogenized. The crossing over of nicked strands between two control regions within an mtDNA molecule leads to formation of a Holliday structure, and the sequence of one control region may be replaced by that of the other via repair of heteroduplex DNA intermediates.

The tandem duplication mechanism, proposed by Kumazawa *et al.* [9], account for not only the generation of duplicate CRs but also for the concerted evolution of them. It involves a replication error triggered by imprecise termination or slipped-strand mispairing, which would lead to duplication of a section. If this section included the CR, then the replication error would generate a mitogenome with two copies of the CRs.

Of the 1500 complete mitogenomes available for teleosts from GenBank (as of November, 2014, <http://www.ncbi.nlm.nih.gov/genome>), duplicate CRs have been found only in three species, including *Kryptolebias marmoratus* (Cyprinodontidae) [12], a righteye flounder, *Samariscus latus* (Samaridae) [13], and a lefteye flounder, *Laeops lanceolata* (Bothidae) [22]. Only a single species from each of these families has been found with this phenomenon. The studies related to these species mainly focused on gene rearrangements, or on the evolutionary pattern conducted only in a single species; both situations represent only narrow taxonomic coverage. No concerted evolution of CRs has been reported in these species or in broad taxa above the level of species. Even within the entire group of teleosts, no phylogenetic study about concerted evolution of duplicate CRs has ever been reported before.

The family Bothidae (excluding species now assigned to the Paralichthyidae) has been recovered as a monophyletic lineage within the Pleuronectiformes in several studies [22–27]. Based on the previous finding of duplicate CRs in the mitogenome of *L. lanceolata*, here we sequenced five new mitogenomes from species in the family Bothidae to test whether or not duplicate CRs exist in these species. Also, together with six previously reported complete mitogenomes in this family, we'll explore how duplicate CRs were generated and what the evolutionary pattern and molecular mechanism of CRs exist in this monophyletic lineage representing 11 species from seven genera. We hope our results will provide fresh insight into molecular mechanisms related to replication and the evolution of mitogenomes.

Materials and Methods

Ethics statement

This study was carried out in strict accordance with the recommendations and guidelines of the National Institutes of Health. No specific permits were required for specimens used in the present study because they are common marine fishes captured in commercial fisheries. *Arnoglossus tenuis* was collected from Shipu commercial fisheries, Zhejiang, China; *Chascanopsetta lugubris* and *Crossorhombus valderostratus* from commercial fisheries in Durban, South Africa; *Lophonectes gallus* from the Sydney Fish Market, Australia; *Psettina iijimae* from Xingda commercial fisheries, Kaohsiung, Taiwan. These species are not included in the endangered species list of the IUCN (<http://www.iucnredlist.org/>). All species are small- to moderate-sized species of flatfishes characterized by both eyes on the left side, with pectoral and pelvic fin rays not branched [28, 29].

Sampling, DNA extraction, PCR amplification and sequencing

Mitogenomes for *A. tenuis*, *C. lugubris*, *C. valderostratus*, *L. gallus* and *P. iijimae* were sequenced for the present study and have been submitted to GenBank (Table 1). Total genomic DNA was extracted from the muscle of these specimens using an SQ Tissue DNA Kit (OMEGA, Guangzhou, China) according to the standard manufacturer's protocol. Dozens of primer pairs (S1 Table) were designed for the fragment amplifications of five flatfish mitogenomes based on the method of aligning sequences of the flatfish mitogenomes previously reported by Shi *et al.* [30].

PCR was performed in a 25 µl reaction volume containing 2.0 mM MgCl₂, 0.4 mM of each dNTP, 0.5 µM of each primer, 1.0 U of Taq polymerase (Takara, China), 2.5 µl of 10x Taq buffer, and approximately 50 ng of DNA template. The amplification profile included an initial denaturation at 95°C for 3 min, 35 cycles of a denaturation at 94°C for 45 s, an annealing temperature of 45–55°C for 45 s, and elongation at 68–72°C for 1.5–5 min. The PCR reaction was completed by a final extension at 72°C for 5 min. The PCR products were detected in 1.0% agarose gels and purified with the TaKaRa Agarose Gel DNA Purification Kit (TaKaRa, China) and used directly as templates for cycle sequencing reactions in both directions with the ABI

Table 1. Flatfishes used in this study.

Classification	Species	Abbreviation	LEF/REF*	Accession No.*	Duplicate CRs
Bothidae	<i>Arnoglossus polyspilus</i>	A.po	LEF	NC_024946	NO
	<i>Arnoglossus tenuis</i>	A.te	.LEF.	<u>KP134337</u>	YES
	<i>Bothus myriaster</i>	B.my	LEF	KJ433563	NO
	<i>Bothus pantherinus</i>	B.pa	LEF	NC_024947	NO
	<i>Crossorhombus azureus</i>	C.az	LEF	JQ639068	NO
	<i>Crossorhombus kobensis</i>	C.ko	LEF	NC_024949	NO
	<i>Crossorhombus valderostratus</i>	C.va	LEF	<u>KJ433566</u>	NO
	<i>Chascanopsetta lugubris</i>	C.lu	LEF	<u>KJ433561</u>	NO
	<i>Lophonectes gallus</i>	L.ga	LEF	<u>KJ433567</u>	YES
	<i>Laeops lanceolata</i>	L.la	LEF	AP014591	YES
	<i>Psettina iijimae</i>	P.ii	LEF	<u>KP134336</u>	YES
Paralichthyidae	<i>Paralichthys olivaceus</i>	P.oli	LEF	NC_002386	NO
Pleuronectidae	<i>Platichthys stellatus</i>	P.ste	REF	NC_010966	NO

*: LEF/REF means Lefteye or Righteye flounder. The underlined number indicates newly-sequenced mtDNAs for five species of Bothids.

doi:10.1371/journal.pone.0134580.t001

3730 DNA sequencer (Applied Biosystems, USA). Sequence specific primers were further designed and used as walking primers for both strands of each fragment.

Sequence analysis

All sequenced fragments were assembled to create complete mitochondrial genomes using CodonCode Aligner (version 3.7.1) and BioEdit (version 7.0.1) [31], and then manually checked against possible sequencing errors. Annotation and boundary determination of protein-coding genes, rRNA and tRNA genes were performed using NCBI-BLAST. Identification of the CR is based on identifying the symbolic structures of CRs: Conserved Sequence Blocks (CSBs) and Termination-Associated Sequences (TASs) [30]. Tandem repeated sequences (TRs) were identified using Tandem Repeats Finder 4.03 in the default parameters. The genetic distances of Kimura-2 Parameter between the core region sequences of CRs were calculated in the MEGA 5.0 program [32].

Phylogenetic analysis and ancestral state reconstruction

Eleven mitogenomes of species of Bothidae, five from the present study and six others retrieved from Genbank, were used for phylogenetic analyses. Based on previous studies [22, 24–26], two species, *Platichthys stellatus* (Pleuronectidae) and *Paralichthys olivaceus* (Paralichthyidae), which have a close relationship with bothids, were chosen as the outgroup (Table 1). Two datasets were generated: one included the first and second codon position sites for each of 12 coding genes (without the third codon position site and the ND6 gene following studies by Campbell *et al.* [22]), the rRNA genes (R), and tRNA genes (T) (noted as 12RT, S1 Fig); the other dataset included only the core region of CRs excluding the tandem repeat sequences (S2 Fig). Before phylogenetic tree estimations, multiple alignment within each dataset was performed using Clustal X [33], then the two datasets were used for maximum likelihood (ML) analyses implemented in PhyML [34] and Bayesian inference (BI) in MrBayes 3.1.2 [35]. The support values for ML analyses were evaluated with 1000 non-parametric bootstrap replicates and the best-fit evolutionary models were determined using Modeltest 3.7 [36]. The best-fit

models of nucleotide substitution for the BI analyses were selected with MrModeltest 2.1 [37] and the BI trees were constructed using MrBayes ver.3.2.1 on 10,000,000 generations and 100 sampled generations [35]. The phylogenetic signal was tested based on the existence of CRs in 11 bothids with two other flatfishes as outgroup using the commonly employed metrics: Pagel's λ [38]. Calculations of Pagel's λ were conducted using the R package Geiger [39–41]. The ancestral state was reconstructed by using maximum likelihood in the Ape package of the R statistical environment [40, 41].

Results and Discussion

Genome organization

The mitogenomes of *A. tenuis*, *C. lugubris*, *C. valderostratus*, *L. gallus* and *P. iijimae* were all circular molecules, with 17,556 bp, 17,251 bp, 16,790 bp, 18,642 bp and 18,080 bp in length, respectively. All of them contained 37 genes, including 13 protein-coding genes, two rRNA genes and 22 tRNA genes. Most of these genes were encoded on the heavy-strand, while *ND6* and eight tRNA genes were encoded on the light-strand (S2 Table). All tRNA genes could be folded into typical cloverleaf structures except for the *tRNA-C*. Two large non-coding (NC) regions were found in these five species, one was located between *tRNA-P* and *tRNA-F*, the other was situated between *tRNA-T* and *tRNA-Q*. Comparison of these NC sequences with the CR sequences of other flatfishes revealed that *C. lugubris* and *C. valderostratus* had one CR located between *tRNA-T* and *tRNA-Q*, with lengths of 855 bp and 704 bp, respectively. However, *A. tenuis*, *L. gallus* and *P. iijimae* had two similar control regions, which were termed CR1 (between *tRNA-P* and *tRNA-F*) and CR2 (between *tRNA-T* and *tRNA-Q*). The length of CR1 ranged from 966–1,513 bp and CR2 ranged from 830–836 bp (S2 Table). The length differences between the CRs mainly resulted from the presence of tandem repeats with extensive variation in sizes and copy numbers (Fig 1A and 1B).

Concerted evolution of duplicate CRs

Comparisons of six previously determined mitogenomes of bothid species [22, 30, 42], reveal that only one species, *L. lanceolata*, has duplicate CRs (Table 1). Together with three other bothids in the present study, a total of four species from different genera in the Bothidae have duplicate CRs (Fig 1A and 1B). The core region sequences of duplicate CRs of mitogenomes within the same species were virtually identical or nearly identical (represented by cross-hatched areas in Fig 1B, about 600 bp in total). Sequence similarity of the core regions of the paralogous CRs (CR1 and CR2) was 99.8–100% in the same species, versus 65.4–76.1% between counterparts of the orthologous CRs in different species (S3 Table, Fig 1B and S2 Fig). Genetic distances of the Kimura-2 Parameter between the core regions of the paralogous CR1 and CR2 are 0–0.2%, versus 23.8–43.6% between the counterparts of the orthologous CRs (S3 Table, S2 Fig). The high similarity and small genetic distances between duplicate CRs within each of four species fit the predicted pattern of concerted evolution [3, 15, 17, 18].

Based on two datasets (12RT and CRs) from 11 bothid species with two other flatfishes as outgroup, ML and BI trees were constructed. The results showed that both trees were largely congruent with each other, therefore, only one topology with both support values was shown, including bootstrap values for the ML tree and posterior probability for Bayesian analysis (Figs 1 and 2). Phylogenetic analyses supported the hypothesis that all 11 species were the descendants of the monophyletic group of Bothidae, but the four flounders with duplicate CRs did not group to one clade. However, the CR1 and CR2 from one species always clustered together (Figs 1 and 2). These results indicated that the four flounders with duplicate CRs are not monophyletic, and that the concerted fashion of duplicate CRs is not a single event that happened




Fig 1. (A) Phylogenetic tree for 11 bothids with two other flatfishes as outgroup using the 12RT dataset. The phylogenetic tree was reconstructed by maximum likelihood and Bayesian methods. The first number beside the internal branch indicates Bayesian posterior probability, the second one is the bootstrap value. **(B)** Gene orders of fragments between tRNA-T and tRNA-F from 11 bothid species. Cross-hatched area in bold indicates the core region of duplicate CRs. Light-gray box indicates the Conserved Sequence Blocks (CSBs) and Termination-Associated Sequences (TASs); Dark-gray box indicates the Tandem Repeat Sequences (TRs); White box indicates the Non-coding region (NC).

doi:10.1371/journal.pone.0134580.g001

only in one lineage of bothid, but rather, it occurred in multiple cases within the family Bothidae.




Fig 2. Phylogenetic tree for 11 bothids with two other flatfishes as outgroup using the core region of CRs. The first number beside the internal branch indicates Bayesian posterior probability (values below 0.9 not shown); the second number represents the bootstrap value (values below 50% not shown).

doi:10.1371/journal.pone.0134580.g002

How are duplicate CRs generated?

So far, the phenomenon of four species with duplicate CRs from different genera within one family (Bothidae) is the first report in fishes. How these duplicate CRs were generated raises a very interesting question. Comparison of the mechanisms for the generation of duplicate CRs from previous reports indicates that the similar dimerization model is suitable to explain the cases in these four species, which is similar to the conclusion of Shi *et al.* [30] who used this mechanism to describe the process for generation and degeneration of duplicate CRs in the mitogenome of *Crossorhombus azureus*.

The complete mitogenome of *C. azureus*, the first one for species of lefteye flounder, has undergone genomic-scale gene rearrangements. The *ND6* and seven tRNA genes (*Q*, *A*, *C*, *Y*, *SI*, *E*, *P*) encoded by the light-strand have been translocated to the position between *tRNA-T*

and *tRNA-F*. It also contains two large non-coding regions (NC1 and NC2), one (NC2) is identified as the CR, the other (NC1) is considered as a remnant from one of the duplicate CRs. Considering the mechanisms previously used to explain the generation of duplicate CRs, Shi *et al.* [30] point out that the tandem duplication mechanism and illegitimate recombination model were not as parsimonious as the dimeric mitogenome model [21]. Therefore, they inferred that the duplicate CRs were generated from a dimeric molecule of mitogenome formed by two monomers linked head-to-tail. Subsequently, one of the CRs was degraded to the NC1 and the other became the new CR [30].

For 11 lefteye flounders in this study, the mitogenomes of four species have the two CRs (CR1 and CR2), whereas seven others have only one CR and in the position where a second CR would occur (Fig 1B), the mitogenome has a large non-coding (NC) region. Further analysis revealed that either the position of the NC or the CR, as well as the cluster of light-strand coding genes, is consistent with the arrangements observed in *C. azureus* (Fig 1B; S2 Table). Thus, we hypothesized that the duplicate CRs of four bothids were generated through the same mechanism outlined in the dimerization model and proposed for the evolution of the mtDNA of *C. azureus* as was described by Shi *et al.* [30].

What is the cause of concerted evolution of duplicate CRs in bothid species?

The evidence of duplicate CRs (CR1 and CR2) in four species and a large non-coding region in seven species with one CR supports the hypothesis that the mitogenomes of seven species once had the other CR at the corresponding position of NC [30]. That is, the mitogenomes of 11 bothid species in the ancestral state have two CRs. Accordingly, we hypothesize that duplicate CRs of the mitogenomes in these flounders are inherited from a common ancestor of present-day bothids.

In order to provide more evidence for this hypothesis, we reconstructed the ancestral state of the CRs character based on the existence of the CRs from 11 bothids with two other flatfishes as outgroup. First, we calculated the phylogenetic signal by Pagel's λ . Pagel's λ is a scaling parameter that ranges from 0 to 1.0 Lambda values. Lambda value of 0 indicates no phylogenetic signal, whereas a value of 1.0 indicates perfect phylogenetic signal. The result of Pagel's λ in this analysis was a 1.0 Lambda value that indicated perfect phylogenetic signal. Therefore, the ancestral state reconstruction (ASR) of the CRs character was deduced by using the maximum likelihood approach (Fig 3). From the highest likelihood values of the ASR result, the evolutionary mode of duplicate CRs was plotted out manually (Fig 4).

The ASR result showed that the copy of CRs had already occurred as early as the origin of the bothids. With expansion of bothid lineages, the duplicate CRs have been retained in some such as *Arnoglossus* (*Arnoglossus tenuis*), *Lophonectes*, *Laeops* and *Psettina*. Meanwhile, in the evolution of other lineages, i.e., *Crossorhombus*, *Arnoglossus* (*Arnoglossus polyspilus*), *Chascanopsetta*, CR1 was degraded and only CR2 was retained. In the *Bothus* lineage, there are two different evolutionary results: one branch lost CR1 and the other kept CR1, but lost CR2. This result supports the hypothesis that the two CRs have the same evolutionary tempo and direction, and it also provides other evidence for our hypothesis that the ancestral state in these bothid species has two CRs (Figs 3 and 4). Furthermore, it also shows that duplicate CRs do not result from recent evolutionary events in these fishes, but rather, they appeared as early as the origin of the bothid lineage (Figs 3 and 4). Subsequently, during evolutionary expansion of bothid lineages from two CRs ancestral state, one of the duplicate CRs was lost in different bothid species at different evolutionary times, whereas, in contrast, other species (lineages) still retained two CRs.




Fig 3. The analysis of ancestral state reconstruction for CR type based on maximum likelihood approach. The pie chart represents the relative likelihood value of alternative CR type. Numbers in color represent the relative likelihood value of the corresponding slice (values below 40% not shown). The blue slice indicates the likelihood that both the CR1 and CR2 were inherited from a common ancestor; the red slice indicates the likelihood of CR1; the green slice indicates the likelihood of CR2. NC represents the non-coding region that was degenerated from the CR.

doi:10.1371/journal.pone.0134580.g003

As for the time of origin of a common ancestor of present-day bothids, studies have discovered that bothids may have appeared about 30 million years ago (mya) based on calculations of the molecular clock [26], or to the late Eocene (ca. 40–56 mya) based on fossil evidence [43].

Over at least 30 mya of evolution, how did those mitogenomes keep two CRs without degeneration and how did duplicate CRs maintain concerted evolution in their core regions? Comparisons with the above three mechanisms about concerted evolution of CRs (the tandem duplication model [9], the deleted and duplicated hypothesis [18] and the gene conversion [9]), revealed that all of them are not suitable here. Primarily, because the first two models hypothesize that concerted evolution of CRs is maintained by generating new duplicate CRs through multiple independent origins. With respect to the gene conversion mechanism [9], it is mainly caused by homologous recombination. It is known that recombination is rarely found in the mitogenomes of teleostean fishes. Based on the phylogenetic trees, the four species of bothids possessing concerted evolution of their CRs belong to different branches rather than one lineage. If the events of concerted evolution of two CRs resulted from one of the three mechanisms mentioned above, the same replication errors should have happened several times in different species; the possibilities of these cases are too low to occur because the mitogenome structure is quite conserved and relatively stable. In addition, Shao *et al.* [17] also considered that it is unlikely that the same replication errors (tandem duplication model) arise over and over again, and independently in each species.

Although the mechanism of concerted evolution of CRs in the Bothidae is not quite sure, further analysis of the structure of CRs of four bothids having concerted evolution can still




Fig 4. The hypothesized evolutionary mode of duplicate CRs of 11 bothids with two flatfishes as outgroup. The mode was hypothesized based on the highest likelihood values of the ASR result. Number and CRs status beside the nodes are based on the highest likelihood value of the ASR result. The two close likelihood values of 40.6% and 42.5% for the node including *B. pantherinus* and *B. myriaster* are shown.

doi:10.1371/journal.pone.0134580.g004

reveal useful information. The symbolic structures of the CR in vertebrate mitogenomes contain some conserved sequences known as CSBs and TASSs. The fragment from the TASSs to CSBs is generally considered to be the core region for regulation of mitochondrial replication and transcription of the heavy-strand [44–48]. The CSBs play a crucial role in both transcription and replication since they are required for the stability of heavy-strand synthesis initiation [49], and the TASSs have a function that can arrest heavy-strand synthesis [47, 48].

In the mitogenomes of the four lefteye flounders with duplicate CRs, the concerted evolution region (identical or nearly identical sequences) in two CRs of each species is restricted in these conserved sections. Although some sequences are very close to these regions, such as the tandem repeat sequences in the mitogenome of *L. gallus*, which has only an 11-base-pair interval to the concerted evolution region of CR1 (Fig 1B), concerted evolution was not found in them because those tandem repeat sequences are not present in CR2. When Eberhard *et al.* [15] studied the concerted evolution of the CRs in parrots, they also pointed out that duplicate CRs may persist only if the duplication event gives rise to complete, functional copies. Thus, we infer that concerted evolution of the CRs in these bothid species may be related to some function of the conserved sequences of the CRs during mitochondrial replication and transcription.

Supporting Information

S1 Fig. Alignment of the sequences of 12RT dataset from mitogenomes of 13 species of flatfishes. Abbreviations of species names are shown in [Table 1](#).
(DOCX)

S2 Fig. Alignment of the core region sequences of the CRs from mitogenomes of 13 species of flatfishes. Abbreviations of species names are shown in [Table 1](#). Numbers following species names represent the types of the CRs: CR1 or CR2. “1” represents the CR1, and “2” represents the CR2, respectively.
(DOCX)

S1 Table. Primers used for fragment amplifications in five flatfish mitogenomes.
(DOCX)

S2 Table. Features and gene maps of the mitogenomes of five species of Bothidae. Intergenic region: non-coding bases between the feature on the same line and the line below, with a negative number indicating an overlap.
(DOCX)

S3 Table. Percent similarity (above triangle) and genetic distance (below triangle) calculated by Kimura-2 Parameter based on core regions of CRs and counterpart sequences from 11 bothids with two other flatfishes as outgroup. Abbreviations of species names are shown in [Table 1](#). Numbers following the species names represent the type of CRs: CR1 or CR2. Numbers in gray are sequence similarity and the genetic distances of Kimura-2 Parameter between core regions of the duplicate CRs within each of four species of flounders.
(DOCX)

Acknowledgments

We are very grateful to C. B. Dillman and X.-X Shen for their helpful comments on the manuscript. We thank A. D. Connell, B. Yau and H.-D Lin for sample collections.

Author Contributions


Conceived and designed the experiments: XYK WS. Performed the experiments: DHL LG. Analyzed the data: WS DHL. Contributed reagents/materials/analysis tools: WS. Wrote the paper: WS. Revised the manuscript critically for important intellectual content: TAM.

References

1. Shadel GS, Clayton DA. Mitochondrial DNA maintenance in vertebrates. *Annu Rev Biochem.* 1997; 66:409–35. doi: [10.1146/annurev.biochem.66.1.409](#) PMID: [WOS:A1997XH20100015](#).
2. Abbott CL, Double MC, Trueman JW, Robinson A, Cockburn A. An unusual source of apparent mitochondrial heteroplasmy: duplicate mitochondrial control regions in *Thalassarche albatrosses*. *Mol Ecol.* 2005; 14(11):3605–13. doi: [10.1111/j.1365-294X.2005.02672.x](#) PMID: [16156827](#)
3. Morris-Pocock JA, Taylor SA, Birt TP, Friesen VL. Concerted evolution of duplicated mitochondrial control regions in three related seabird species. *BMC Evol Biol.* 2010; 10(1):14. doi: [10.1186/1471-2148-10-14](#)
4. Schirtzinger EE, Tavares ES, Gonzales LA, Eberhard JR, Miyaki CY, Sanchez JJ, et al. Multiple independent origins of mitochondrial control region duplications in the order Psittaciformes. *Mol Phylogenet Evol.* 2012; 64(2):342–56. doi: [10.1016/j.ympev.2012.04.009](#) PMID: [22543055](#)
5. Parham JF, Feldman CR, Boore JL. The complete mitochondrial genome of the enigmatic bigheaded turtle (*Platysternon*): description of unusual genomic features and the reconciliation of phylogenetic hypotheses based on mitochondrial and nuclear DNA. *BMC Evol Biol.* 2006; 6:11. doi: [10.1186/1471-2148-6-11](#) PMID: [WOS:000236032500001](#).

6. Parham JF, Macey JR, Papenfuss TJ, Feldman CR, Türkozan O, Polymeni R, et al. The phylogeny of Mediterranean tortoises and their close relatives based on complete mitochondrial genome sequences from museum specimens. *Mol Phylogenet Evol.* 2006; 38(1):50–64. doi: [10.1016/j.ympev.2005.07.015](https://doi.org/10.1016/j.ympev.2005.07.015) PMID: [16150614](https://pubmed.ncbi.nlm.nih.gov/16150614/)
7. Peng Q-L, Nie L-W, Pu Y-G. Complete mitochondrial genome of Chinese big-headed turtle, *Platysternon megacephalum*, with a novel gene organization in vertebrate mtDNA. *Gene.* 2006; 380(1):14–20. doi: [10.1016/j.gene.2006.04.001](https://doi.org/10.1016/j.gene.2006.04.001) PMID: [16842936](https://pubmed.ncbi.nlm.nih.gov/16842936/)
8. Kumazawa Y, Ota H, Nishida M, Ozawa T. Gene rearrangements in snake mitochondrial genomes: highly concerted evolution of control-region-like sequences duplicated and inserted into a tRNA gene cluster. *Mol Biol Evol.* 1996; 13(9):1242–54. PMID: [8896377](https://pubmed.ncbi.nlm.nih.gov/8896377/)
9. Kumazawa Y, Ota H, Nishida M, Ozawa T. The complete nucleotide sequence of a snake (*Dinodon semicarinatus*) mitochondrial genome with two identical control regions. *Genetics.* 1998; 150(1):313–29. PMID: [9725849](https://pubmed.ncbi.nlm.nih.gov/9725849/)
10. Dubey B, Meganathan PR, Haque I. Complete mitochondrial genome sequence from an endangered Indian snake, *Python molurus molurus* (Serpentes, Pythonidae). *Mol Biol Rep.* 2012; 39(7):7403–12. doi: [10.1007/s11033-012-1572-5](https://doi.org/10.1007/s11033-012-1572-5) PMID: [WOS:000304404600020](https://pubmed.ncbi.nlm.nih.gov/WOS:000304404600020/)
11. Lee J-S, Miya M, Lee Y-S, Kim CG, Park E-H, Aoki Y, et al. The complete DNA sequence of the mitochondrial genome of the self-fertilizing fish *Rivulus marmoratus* (Cyprinodontiformes, Rivulidae) and the first description of duplication of a control region in fish. *Gene.* 2001; 280(1):1–7. doi: [10.1016/S0378-1119\(01\)00765-X](https://doi.org/10.1016/S0378-1119(01)00765-X)
12. Tatarenkov A, Avise JC. Rapid concerted evolution in animal mitochondrial DNA. *P Roy Soc B-Biol Sci.* 2007; 274(1619):1795–8. doi: [10.1098/rspb.2007.0169](https://doi.org/10.1098/rspb.2007.0169)
13. Shi W, Miao X-G, Kong X-Y. A novel model of double replications and random loss accounts for rearrangements in the Mitogenome of *Samariscus latus* (Teleostei: Pleuronectiformes). *BMC Genomics.* 2014; 15(1):352. doi: [10.1186/1471-2164-15-352](https://doi.org/10.1186/1471-2164-15-352) PMID: [WOS:000336066900002](https://pubmed.ncbi.nlm.nih.gov/WOS:000336066900002/)
14. Arndt A, Smith MJ. Mitochondrial gene rearrangement in the sea cucumber genus *Cucumaria*. *Mol Biol Evol.* 1998; 15(8):1009–16. Epub 1998/08/27. PMID: [9718728](https://pubmed.ncbi.nlm.nih.gov/9718728/)
15. Eberhard JR, Wright TF, Bermingham E. Duplication and concerted evolution of the mitochondrial control region in the parrot genus *Amazona*. *Mol Biol Evol.* 2001; 18(7):1330–42. PMID: [11420371](https://pubmed.ncbi.nlm.nih.gov/11420371/)
16. Ogoh K, Ohmiya Y. Complete mitochondrial DNA sequence of the sea-firefly, *Vargula hilgendorffii* (Crustacea, Ostracoda) with duplicate control regions. *Gene.* 2004; 327(1):131–9. doi: [10.1016/j.gene.2003.11.011](https://doi.org/10.1016/j.gene.2003.11.011) PMID: [14960368](https://pubmed.ncbi.nlm.nih.gov/14960368/)
17. Shao R, Barker SC, Mitani H, Aoki Y, Fukunaga M. Evolution of duplicate control regions in the mitochondrial genomes of metazoa: a case study with Australasian *Ixodes* ticks. *Mol Biol Evol.* 2005; 22(3):620–9. doi: [10.1093/molbev/msi047](https://doi.org/10.1093/molbev/msi047) PMID: [15537802](https://pubmed.ncbi.nlm.nih.gov/15537802/)
18. Ogoh K, Ohmiya Y. Concerted evolution of duplicated control regions within an ostracod mitochondrial genome. *Mol Biol Evol.* 2007; 24(1):74–8. doi: [10.1093/molbev/msl132](https://doi.org/10.1093/molbev/msl132) PMID: [17000719](https://pubmed.ncbi.nlm.nih.gov/17000719/)
19. Liao D. Concerted evolution: molecular mechanism and biological implications. *Am J Hum Genet.* 1999; 64(1):24–30. doi: [10.1086/302221](https://doi.org/10.1086/302221) PMID: [9915939](https://pubmed.ncbi.nlm.nih.gov/9915939/)
20. Lunt DH, Hyman BC. Animal mitochondrial DNA recombination. *Nature.* 1997; 387(6630):247. Epub 1997/05/15. doi: [10.1038/387247a0](https://doi.org/10.1038/387247a0) PMID: [9153388](https://pubmed.ncbi.nlm.nih.gov/9153388/)
21. Boore JL. The duplication/random loss model for gene rearrangement exemplified by mitochondrial genomes of deuterostome animals. In: Sankoff D, Nadeau J, editors. *Computational Biology. Computational Biology.* 1. Ann Arbor, MI: Springer Netherlands; 2000. p. 133–47.
22. Campbell MA, Lopez JA, Satoh TP, Chen WJ, Miya M. Mitochondrial genomic investigation of flatfish monophyly. *Gene.* 2014; 551(2):176–82. doi: [10.1016/j.gene.2014.08.053](https://doi.org/10.1016/j.gene.2014.08.053) PMID: [25172210](https://pubmed.ncbi.nlm.nih.gov/25172210/)
23. Chapleau F. Pleuronectiform relationships: A cladistic reassessment. *Bull Mar Sci.* 1993; 52(1):516–40. PMID: [BIOSIS:PREV199345039888](https://pubmed.ncbi.nlm.nih.gov/BIOSIS:PREV199345039888/)
24. Betancur-R. R, Li C, Munroe TA, Ballesteros JA, Orti G. Addressing Gene Tree Discordance and Non-Stationarity to Resolve a Multi-Locus Phylogeny of the Flatfishes (Teleostei: Pleuronectiformes). *Syst Biol.* 2013; 62(5):763–85. doi: [10.1093/sysbio/syt039](https://doi.org/10.1093/sysbio/syt039) PMID: [23749787](https://pubmed.ncbi.nlm.nih.gov/23749787/)
25. Betancur-R. R, Orti G. Molecular evidence for the monophyly of flatfishes (Carangimorpharia: Pleuronectiformes). *Mol Phylogenet Evol.* 2014; 73:18–22. doi: [10.1016/j.ympev.2014.01.006](https://doi.org/10.1016/j.ympev.2014.01.006) PMID: [24447882](https://pubmed.ncbi.nlm.nih.gov/24447882/)
26. Campbell MA, Chen WJ, López JA. Are flatfishes (Pleuronectiformes) monophyletic? *Mol Phylogenet Evol.* 2013; 69(3):664–73. doi: [10.1016/j.ympev.2013.07.011](https://doi.org/10.1016/j.ympev.2013.07.011) PMID: [23876291](https://pubmed.ncbi.nlm.nih.gov/23876291/)
27. Campbell MA, Chen WJ, López JA. Molecular data do not provide unambiguous support for the monophyly of flatfishes (Pleuronectiformes): A reply to Betancur-R. and Orti. *Mol Phylogenet Evol.* 2014; 75(0):149–53. doi: [10.1016/j.ympev.2014.02.011](https://doi.org/10.1016/j.ympev.2014.02.011)

28. Li SZ, Wang HM. Fauna Sinica (Osteichthyes): Pleuronectiformes. Beijing: Science Press; 1995.
29. Nelson JS. Fishes of the world. 4th ed. New York: John Wiley, Inc.; 2006. 601 p.
30. Shi W, Dong XL, Wang Z-M, Miao X-G, Wang S-Y, Kong X-Y. Complete mitogenome sequences of four flatfishes (Pleuronectiformes) reveal a novel gene arrangement of L-strand coding genes. BMC Evol Biol. 2013; 13:173. doi: [10.1186/1471-2148-7-247](https://doi.org/10.1186/1471-2148-7-247) PMID: [ISI:000253469400001](https://pubmed.ncbi.nlm.nih.gov/25346940/).
31. Hall TA. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. Nucleic acids symposium series. 1999; 41:95–8.
32. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: Molecular Evolutionary Genetics Analysis Using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods. Mol Biol Evol. 2011; 28(10):2731–9. doi: [10.1093/molbev/msr121](https://doi.org/10.1093/molbev/msr121) PMID: [WOS:000295184200003](https://pubmed.ncbi.nlm.nih.gov/22000003/).
33. Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG. The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. Nucleic Acids Res. 1997; 25(24):4876–82. Epub 1998/02/28. gka797 [pii]. PMID: [9396791](https://pubmed.ncbi.nlm.nih.gov/9396791/).
34. Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. Syst Biol. 2010; 59(3):307–21. doi: [10.1093/sysbio/syq010](https://doi.org/10.1093/sysbio/syq010) PMID: [20525638](https://pubmed.ncbi.nlm.nih.gov/20525638/).
35. Ronquist F, Huelsenbeck JP. MrBayes 3: Bayesian phylogenetic inference under mixed models. Bioinformatics. 2003; 19(12):1572–4. Epub 2003/08/13. PMID: [12912839](https://pubmed.ncbi.nlm.nih.gov/12912839/).
36. Posada D, Crandall KA. MODELTEST: testing the model of DNA substitution. Bioinformatics. 1998; 14(9):817–8. doi: [10.1093/bioinformatics/14.9.817](https://doi.org/10.1093/bioinformatics/14.9.817) PMID: [9918953](https://pubmed.ncbi.nlm.nih.gov/9918953/).
37. Nylander JA, Ronquist F, Huelsenbeck JP, Nieves-Aldrey JL. Bayesian phylogenetic analysis of combined data. Syst Biol. 2004; 53(1):47–67. Epub 2004/02/18. C5P30PXY4HAWE7BM [pii]. PMID: [14965900](https://pubmed.ncbi.nlm.nih.gov/14965900/).
38. Pagel M. Inferring the historical patterns of biological evolution. Nature. 1999; 401(6756):877–84. doi: [10.1038/44766](https://doi.org/10.1038/44766) PMID: [WOS:000083464700048](https://pubmed.ncbi.nlm.nih.gov/200083464700048/).
39. Harmon LJ, Weir JT, Brock CD, Glor RE, Challenger W. GEIGER: investigating evolutionary radiations. Bioinformatics. 2008; 24(1):129–31. doi: [10.1093/bioinformatics/btm538](https://doi.org/10.1093/bioinformatics/btm538) PMID: [WOS:000251865000018](https://pubmed.ncbi.nlm.nih.gov/2000251865000018/).
40. Paradis E, Claude J, Strimmer K. APE: Analyses of Phylogenetics and Evolution in R language. Bioinformatics. 2004; 20(2):289–90. PMID: [14734327](https://pubmed.ncbi.nlm.nih.gov/14734327/).
41. Ihaka R, Gentleman R. R: a language for data analysis and graphics. J Comput Graph Stat. 1996; 5(3):299–314.
42. Gong L, Shi W, Yang M, Li D, Kong X. Novel gene arrangement in the mitochondrial genome of *Bothus myriaster* (Pleuronectiformes: Bothidae): evidence for the Dimer-Mitogenome and Non-random Loss model. Mitochondrial DNA. 2015; 1–4. doi: [10.3109/19401736.2014.1003922](https://doi.org/10.3109/19401736.2014.1003922) PMID: [25629499](https://pubmed.ncbi.nlm.nih.gov/25629499/).
43. Schwarzhans W. A comparative morphological treatise of recent and fossil otoliths of the order Pleuronectiformes. Germany: Verlag F. Pfeil; 1999.
44. Clayton DA. Replication of animal mitochondrial DNA. Cell. 1982; 28(4):693–705. Epub 1982/04/01. doi: [10.1016/0092-8674\(82\)90049-6](https://doi.org/10.1016/0092-8674(82)90049-6) PMID: [6178513](https://pubmed.ncbi.nlm.nih.gov/6178513/).
45. Holt IJ, Lorimer HE, Jacobs HT. Coupled leading- and lagging-strand synthesis of mammalian mitochondrial DNA. Cell. 2000; 100(5):515–24. Epub 2000/03/18. doi: [10.1016/S0092-8674\(00\)80688-1](https://doi.org/10.1016/S0092-8674(00)80688-1) PMID: [10721989](https://pubmed.ncbi.nlm.nih.gov/10721989/).
46. Brown TA, Clayton DA. Genesis and wanderings—Origins and migrations in asymmetrically replicating mitochondrial DNA. Cell Cycle. 2006; 5(9):917–21. doi: [10.4161/cc.5.9.2710](https://doi.org/10.4161/cc.5.9.2710) PMID: [ISI:000238575200003](https://pubmed.ncbi.nlm.nih.gov/2000238575200003/).
47. Fernandez-Silva P, Enriquez JA, Montoya J. Replication and transcription of mammalian mitochondrial DNA. Exp Physiol. 2003; 88(1):41–56. doi: [10.1113/Eph8802514](https://doi.org/10.1113/Eph8802514) PMID: [WOS:000182357200008](https://pubmed.ncbi.nlm.nih.gov/2000182357200008/).
48. Clayton DA. Mitochondrial DNA replication: what we know. IUBMB Life. 2003; 55(4–5):213–7. Epub 2003/07/26. doi: [10.1080/1521654031000134824](https://doi.org/10.1080/1521654031000134824) PMID: [12880201](https://pubmed.ncbi.nlm.nih.gov/12880201/).
49. Fonseca MM, Harris DJ, Posada D. The Inversion of the Control Region in Three Mitogenomes Provides Further Evidence for an Asymmetric Model of Vertebrate mtDNA Replication. PLoS One. 2014; 9(9):e106654. doi: [10.1371/journal.pone.0106654](https://doi.org/10.1371/journal.pone.0106654) PMID: [25268704](https://pubmed.ncbi.nlm.nih.gov/25268704/).



Mitochondrial DNA Part B Resources

ISSN: (Print) 2380-2359 (Online) Journal homepage: <https://www.tandfonline.com/loi/tmdn20>

The complete mitochondrial genome sequence of *Asterorhombus intermedius* (Pleuronectiformes: Bothidae)

Shixi Chen, Hairong Luo, Wei Shi & Xiaoyu Kong

To cite this article: Shixi Chen, Hairong Luo, Wei Shi & Xiaoyu Kong (2019) The complete mitochondrial genome sequence of *Asterorhombus intermedius* (Pleuronectiformes: Bothidae), Mitochondrial DNA Part B, 4:1, 756-757, DOI: [10.1080/23802359.2019.1565969](https://doi.org/10.1080/23802359.2019.1565969)

To link to this article: <https://doi.org/10.1080/23802359.2019.1565969>



© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 08 Feb 2019.



Submit your article to this journal [↗](#)



Article views: 15



View Crossmark data [↗](#)

The complete mitochondrial genome sequence of *Asterorhombus intermedius* (Pleuronectiformes: Bothidae)

Shixi Chen^{a,b}, Hairong Luo^{a,b}, Wei Shi^a and Xiaoyu Kong^a

^aCAS Key Laboratory of Tropical Marine Bio-resources and Ecology, South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou, China; ^bUniversity of Chinese Academy of Sciences, Beijing, China

ABSTRACT

The complete mitogenome of *Asterorhombus intermedius* is 16,886 bp in length, containing 37 genes, among them, *ND6* and eight tRNA genes are encoded by L-strand, and other genes by H-strand, which are the same as those of typical mitogenome in fishes. A striking finding is a novel genomic-scale gene rearrangement related nine genes, including *ND6* and six tRNA encoded by the L-strand grouped to a cluster of *Gln-Ala-Cys-Tyr-Ser₁-ND6-Glu*. Also, the genes of this cluster with identical transcriptional polarities maintained the original gene order. In addition, the order of *12S-Val-16S-Leu₁* fragment rearranged to *12S-16S-Val-Leu₁*, and that of *Ser₁-Asp-COII* fragment rearranged to *Ser₁-Asp-ND6*.

ARTICLE HISTORY

Received 13 December 2018
Accepted 23 December 2018

KEYWORDS

Mitogenome; bothid species; *Asterorhombus intermedius*; gene rearrangement

Asterorhombus intermedius (Bleeker 1865) belongs to the family Bothidae of Pleuronectiformes (Amaoka 1969). The mitochondrial genome of *A. intermedius* were obtained (GenBank accession number: MK256952) based on one sample collected from Sanya City, Hainan, China (N18°16'33.87", E109°43'33.24"). This specimen was stored in the Marine Biodiversity Collection of South China Sea, Chinese Academy of Sciences, under the voucher number SCF2015121552.

The mitochondrial genome of *A. intermedius* is 16,886 bp in length, including 13 protein-coding genes, two ribosomal RNA (rRNA), 22 transfer RNA (tRNA), and a control region (CR). Among these genes, 28 genes are encoded on the H-strand, others including *ND6* and eight tRNA genes on the L-strand. The overall base composition is 27.6, 30.0, 16.5, and 25.9% for A, C, G, T, respectively, with a slight bias on AT content (53.5%).

The size of 13 protein-coding genes ranges from 168 bp (*ATP8*) to 1818 bp (*ND5*), that of *12S* and *16S* rDNA are 944 bp and 1653 bp, respectively and that of CR is 1042 bp. The size of 22 tRNA genes ranges from 65 bp (*tRNA-Cys*) to 73 bp (*tRNA-Leu₂*), which were identified using *tRNAscan-SE* 2.0 (Lowe and Chan 2016).

A striking finding of this study is a novel genomic-scale gene rearrangement related nine genes, which was not similar to both of the typical fishes and other bothid species (Wang et al. 2015; Gong et al. 2016; Li et al. 2015). In the mitogenome of this species, the *ND6* and six tRNA encoded by the L-strand grouped to a seven-gene cluster of

Gln-Ala-Cys-Tyr-Ser₁-ND6-Glu at the position between *tRNA-T* and *tRNA-P*. Besides, the genes of this cluster with identical transcriptional polarities maintained the original gene order. Also, the *tRNA-Asp* translocated from the position between *tRNA-Ser₁* and *COII* to the position between *tRNA-Ser₂* and *ND6*, which formed the eight-gene order of *Gln-Ala-Cys-Tyr-Ser₁-Asp-ND6-Glu*. In addition, the order of *12S-Val-16S-Leu₁* fragment rearranged to *12S-16S-Val-Leu₁*.

This mitogenome existed in 20 intergenic spacers with total 313 bp in length. Comparing with that of non-rearranged species, 10 abnormal intergenic regions with total 282 bp in length were found, among them, the longest is 98 bp and the shortest is 4 bp.

The phylogenetic tree was constructed based on the mitogenome of *A. intermedius* and other 11 bothid species with *Paralichthys olivaceus* and *Platichthys stellatus* as outgroup by using Bayesian inference in MrBayes 3.2 (Ronquist and Huelsenbeck 2003). The mitogenome sequences of 11 species and 3 outgroups were retrieved from GenBank. In the phylogenetic tree (Figure 1), three species of *Crossorhombus* and two species of *Bothus* formed two clades, respectively, both with 100% posterior probability. However, two species of *Arnoglossus* were not clustered together. *A. intermedius* and *Chascanopsetta lugubris* formed a clade with 100% posterior probability, which means *A. intermedius* has a relatively closer phylogenetic relationship to *C. lugubris*, compared with other 10 bothid species.




Figure 1. Phylogenetic tree of Bayesian Inference was constructed based on the mitogenome sequences from 12 species in Bothidae with two outgroup. The number beside the branch indicates Bayesian posterior probability.

Disclosure statement

The authors declare that there is no conflict of interest regarding the publication of this article. The authors alone are responsible for the content and writing of the paper.

Funding

This work was supported by the Natural Science Foundation of China [31471979 and 31872570].

Reference

Amaoka, K. 1969. Studies on the sinistral flounders found in the waters around Japan: taxonomy, anatomy and phylogeny. *J Natl Fish Univ.* 18:65–340.

Gong L, Shi W, Yang M, Li DH, Kong XY. 2016. Novel gene arrangement in the mitochondrial genome of *Bothus myriaster* (Pleuronectiformes: Bothidae): evidence for the dimer-mitogenome and non-random loss model. *Mitochondrial DNA A DNA Mapp Seq Anal.* 27:3089–3092.

Li DH, Shi W, Munroe TA, Gong L, Kong XY. 2015. Concerted evolution of duplicate control regions in the mitochondria of species of the flatfish family Bothidae (Teleostei: Pleuronectiformes). *PLoS One.* 10: e0134580.

Lowe TM, Chan PP. 2016. tRNAscan-SE On-line: integrating search and context for analysis of transfer RNA genes. *Nucleic Acids Res.* 44: W54–57.

Ronquist F, Huelsenbeck JP. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics.* 19:1572–1574.

Wang SY, Shi W, Wang ZM, Gong L, Kong XY. 2015. The complete mitochondrial genome sequence of *Aesopia cornuta* (Pleuronectiformes: Soleidae). *Mitochondrial DNA.* 26:114–115.