Research Square

Preprints are preliminary reports that have not undergone peer review.
They should not be considered conclusive, used to inform clinical practice,
or referenced by the media as validated information.

Analysis of annual soil loss due to erosion processes in the Zarkent massif Parkent district, Uzbekistan, using the RUSLE model

Zhang Wen
Shaanxi A&F Technology University
Gulchehra Sodikova
Tashkent State Agrarian University
Alemayehu Tolossa
Jimma University
Lazizakhon Gafurova
National University of Uzbekistan
Tulkin Shamsiddinov
Tashkent State Agrarian University
Trobjon Makhkamov
mturobzho@mail.ru

Tashkent State Agrarian University

Research Article

Keywords: Mountain soils, Erosion risk, RUSLE, GIS, Soil conservation

Posted Date: July 10th, 2026

DOI: https://doi.org/10.21203/rs.3.rs-10047519/v1

License: This work is licensed under a Creative Commons Attribution 4.0 International License.
Read Full License

Additional Declarations: No competing interests reported.

Abstract

Soil erosion represents one of the most pressing environmental challenges in mountainous regions of Central Asia, where fragile soil systems, steep topography, and unsustainable land management practices converge to accelerate land degradation. This study employs the Revised Universal Soil Loss Equation (RUSLE) model integrated with Geographic Information Systems (GIS) and remote sensing techniques to estimate annual soil losses in the Zarkent massif Parkent district, a mountainous region of Uzbekistan, over the period 2015 to 2024. The model quantifies soil loss through five primary parameters: rainfall erosivity (R-factor), soil erodibility (K-factor), slope length and steepness (LS-factor), cover management (C-factor), and support practice (P-factor). Results indicate that the average annual soil loss in severely eroded areas exceeds 40 $t \cdot ha^{-1} \cdot yr^{-1}$, with 68.84 hectares experiencing erosion rates above this threshold. The majority of the study area (99%) falls within the 0–5 $t \cdot ha^{-1} \cdot yr^{-1}$ category, indicating that erosion is spatially concentrated in localized hotspots rather than uniformly distributed across the landscape. Thematic maps generated from RUSLE outputs identify erosion-prone zones with high precision, thereby guiding targeted soil conservation measures such as terracing, afforestation, and improved rainfed crop management. This study contributes to the growing body of RUSLE applications in data-sparse regions of Central Asia and provides a replicable methodology for erosion risk assessment under semi-arid mountain conditions. However, the study is limited by the absence of field validation data and the exclusion of gully erosion processes from the RUSLE framework.

1 Introduction

Soil erosion, exacerbated by global climate change and anthropogenic activities, is a major driver of land degradation, threatening agricultural productivity, ecosystem sustainability, and food security worldwide. The Food and Agriculture Organization (FAO) estimates that by 2050, soil erosion could reduce global crop production by 10%, resulting in the annual loss of 75 billion tons of soil (Food and Agriculture Organization, 2012). Erosion risk is influenced by a complex interplay of natural factors including climate, geomorphology, and soil parent material and human activities such as deforestation, overgrazing, and inappropriate agricultural practices (Terefe et al., 2025). In mountainous regions, topography emerges as a primary driver of erosion processes, influencing soil morphometric indicators, erosion progression on slopes, and vegetation distribution patterns (Byizigiro et al., 2020). The morphological characteristics of soils govern surface moisture and heat distribution, as well as runoff velocity, ultimately leading to nutrient loss through selective erosion of fine particles (Okenmuo & Ewemoje, 2023).

In Central Asia, Uzbekistan faces growing pressure on land resources from rapid population growth, agricultural expansion, and the intensifying impacts of climate change. The foothill zones of the Tashkent region are particularly erosion-sensitive, characterized by fragile sierozem soils, a semi-arid climate, and widespread unsustainable land management practices (Bobomurodov et al., 2023). Sierozem soils, which dominate this region, are inherently low in organic matter typically containing between 0.93% and 1.71% humus in the plow layer and exhibit moderate to high susceptibility to erosion when subjected to intensive rainfall or improper irrigation (Bobomurodov et al., 2023; Juliev et al., 2022).

These conditions collectively increase the risk of soil loss and declining agricultural productivity, with potentially severe socio-economic consequences for rural communities dependent on rainfed agriculture.

Geographic Information Systems (GIS) have become indispensable tools for managing degraded soils and predicting erosion risks, enabling comprehensive spatial analysis through advanced software platforms (Barboza et al., 2024). Soil erosion modeling remains inherently complex, with erosion risk assessment being a priority for sustainable land management. Soils affected by adverse natural conditions particularly climate, relief, and precipitation patterns are prone to erosion, especially within agricultural landscapes (Girmay et al., 2020). Despite extensive global applications of the RUSLE model, few studies have quantified soil erosion in Uzbekistan's mountainous regions, particularly the Parkent district, where topographic and climatic factors exacerbate erosion risks (Khan et al., 2023; Juliev et al., 2024). Recent research in the Ugam Chatkal National Park, located in the same region, has demonstrated that R and LS factors exert greater influence on soil erosion than other parameters, with estimated average annual soil erosion rates ranging between 0.00 and 250 $t \cdot ha^{-1} \cdot yr^{-1}$ (Juliev et al., 2024).

Several erosion modeling approaches have been developed and implemented in software systems, including AGNPS, WEPP, USLE, RUSLE, and MUSLE. These models typically divide watersheds into grid cells with attributes such as slope, rainfall erosivity, and soil erodibility (Terefe et al., 2025). The RUSLE model, an empirical equation revised from the Universal Soil Loss Equation (USLE) by the United States Department of Agriculture (USDA) in 1997, calculates long-term average annual soil loss due to sheet and rill erosion, considering water erosivity, soil erodibility, slope characteristics, land cover, and management practices (Renard et al., 1997; Benavidez et al., 2018). The RUSLE model was selected for this study due to its relatively low data requirements and its proven compatibility with ArcGIS, facilitating integration with diverse geospatial datasets (Abdul Rahaman et al., 2015).

The primary objectives of this study are threefold: first, to estimate annual soil loss in the Parkent district from 2015 to 2024 using the RUSLE model integrated with GIS and remote sensing; second, to identify and map erosion risk zones through thematic analysis of RUSLE factor outputs; and third, to propose targeted soil conservation measures based on the spatial distribution of erosion risk across the study area. This research is particularly timely given recent initiatives by the National Committee on Ecology and Climate Change of the Republic of Uzbekistan, which has been actively promoting training seminars on soil erosion assessment using the RUSLE model in collaboration with the International Center for Agricultural Research in the Dry Areas (ICARDA) (National Committee on Ecology and Climate Change of Uzbekistan, 2025).

2 Materials and Methods

2.1 Study Area Description

The study was conducted in the foothill regions of the Zarkent massif Parkent district, located in the Tashkent region of eastern Uzbekistan (Fig. 1). The Zarkent massif is situated on the northern slopes of the eastern Tien-Shan mountain range, forming the eastern part of the Tashkent region approximately 48 km east of Tashkent city (Bobomurodov et al., 2023). The geographical coordinates of the study area range from approximately 40°58' N to 41°15' N latitude and 69°21' E to 69°45' E longitude.

The elevation of the study area ranges from 500 to 1,200 m above sea level, with terrain complexity increasing progressively from the southwestern lowlands to the northeastern mountainous zones. The surface of the region is characterized by a complex topography that gradually decreases from northeast to southwest toward the Syrdarya River, with the highest lands concentrated in the northeast and eastern sections of the district (Sodikova et al., 2025). This topographic configuration exerts a fundamental control on local climate patterns, soil formation processes, and erosion dynamics.

The climate of the Parkent district is semi-arid and sharply continental, characterized by hot, dry summers and cold winters with precipitation concentrated mainly in winter and early spring (Bobomurodov et al., 2023). Mean annual precipitation ranges from 400 to 640 mm, with considerable interannual variability. The northeastern mountainous portions of the region receive substantially higher precipitation due to their exposure to moist westerly winds, while the southwestern lowlands are significantly drier. Mean air temperature in July ranges from 26–27°C in the plains to 20–24°C in the mountains, with summer maximum temperatures occasionally reaching 42–44°C (Bobomurodov et al., 2023).

Soils in the study area are predominantly sierozems (gray soils) and meadow-sierozem complexes, which have developed on loess-like and alluvial-proluvial deposits. These soils are characterized by moderate to heavy loamy texture, with physical clay content (< 0.01 mm) ranging from 16–18% in light mechanical sands to 45–58% in heavy loams (Bobomurodov et al., 2023). The organic matter content in the plow layer (0–35 cm depth) typically varies between 0.93% and 1.71%, placing these soils in the low-to-moderate humus category. Below the plow layer, humus content decreases significantly to 0.66–0.93% in subsurface horizons (Bobomurodov et al., 2023). Total nitrogen content ranges from 0.072% to 0.099%, while total phosphorus is relatively abundant (0.115–0.423%), though mobile phosphorus availability is low (3.98–12.71 mg/kg) (Bobomurodov et al., 2023). These soil properties render the sierozems moderately to highly susceptible to water erosion, particularly on sloping terrain where surface runoff velocities increase substantially.

Agriculture constitutes the primary land-use activity in the Parkent district, dominated by rainfed cropping systems and livestock grazing on natural pastures. The major crops cultivated include wheat, barley, and fodder crops, with some areas devoted to cotton under irrigated conditions on the lower slopes and valley bottoms (Bobomurodov et al., 2023). Land management practices vary considerably across the district, with terracing and contour farming implemented in some areas but largely absent from steeper slopes where erosion risk is greatest.

2.2 The RUSLE Model: Theoretical Framework

The Revised Universal Soil Loss Equation (RUSLE) is an empirical model developed by the United States Department of Agriculture (USDA) to predict long-term average annual soil loss caused by sheet and rill erosion (Renard et al., 1997). RUSLE represents a significant revision of the original Universal Soil Loss Equation (USLE), incorporating updated factor relationships, improved computational algorithms, and expanded applicability to a wider range of land use and climatic conditions (Benavidez et al., 2018). The model calculates average annual soil loss (A) as the product of five factors according to the following equation:

$$A = R \times K \times LS \times C \times P$$

where A is the average annual soil loss ($t \cdot ha^{-1} \cdot yr^{-1}$); R is the rainfall erosivity factor ($MJ \cdot mm \cdot ha^{-1} \cdot h^{-1} \cdot yr^{-1}$), representing the erosive potential of rainfall events; K is the soil erodibility factor ($t \cdot ha \cdot h \cdot ha^{-1} \cdot MJ^{-1} \cdot mm^{-1}$), quantifying the intrinsic susceptibility of soil to detachment and transport by rainfall and runoff; LS is the slope length and steepness factor (dimensionless), accounting for topographic effects on flow accumulation and velocity; C is the cover management factor (dimensionless), representing the effect of vegetation cover and land management on soil loss; and P is the support practice factor (dimensionless), reflecting the effectiveness of conservation practices such as contour farming, terracing, and strip cropping.

The RUSLE model has been widely validated and applied across diverse environmental conditions worldwide, from temperate agricultural regions to tropical watersheds and mountainous terrains (Benavidez et al., 2018; Ganasri & Ramesh, 2015; Terefe et al., 2025). Its integration with GIS enables spatially explicit mapping of erosion risk, facilitating the identification of priority areas for conservation intervention (Barboza et al., 2024; Abdul Rahaman et al., 2015).

2.3 Data Sources and Processing

Table 1 summarizes the data sources, spatial resolutions, and temporal coverage utilized in this study for RUSLE factor calculation.

Table 1
Data sources, spatial resolution, and temporal coverage for RUSLE factor calculation.

Parameter	Data Source	Spatial Resolution	Temporal Coverage	Reference
R-factor	CHIRPS precipitation (Climate Hazards Group InfraRed Precipitation with Station Data)	0.05° (~ 5 km)	2015–2024 (daily, aggregated to annual)	Funk et al., 2015
K-factor	OpenLandMap soil properties + field sampling (3 profiles)	250 m + point data	0–30 cm depth	Hengl et al., 2017; Bobomurodov et al., 2023
LS-factor	SRTM Digital Elevation Model (DEM)	30 m	2000 (static)	NASA JPL, 2013
C-factor	Landsat 8 OLI/TIRS (Collection 2 Level-2)	30 m	2015–2024 (growing season composites)	USGS, 2024
P-factor	GlobeLand30 Land Use/Land Cover + field verification	30 m	2020	Chen et al., 2015

All spatial data processing, factor calculation, raster algebra operations, and map generation were performed using ArcGIS 10.5 (ESRI, Redlands, CA, USA). All input layers were reprojected to a common coordinate system (UTM zone 42N, WGS84 datum) and resampled to a uniform 30 m grid resolution to ensure compatibility for raster multiplication.

2.4 Calculation of RUSLE Factors

2.4.1 Rainfall Erosivity (R-Factor)

The rainfall erosivity factor (R-factor) quantifies the erosive potential of precipitation based on the kinetic energy and intensity of rainfall events. In this study, the R-factor was derived from CHIRPS precipitation data, which provides quasi-global precipitation records at 0.05° spatial resolution from 1981 to the present (Funk et al., 2015). Daily precipitation data for the Parkent district Zarkent massif were extracted for the period 2015–2024 and aggregated to annual totals for each grid cell within the region of interest (ROI).

The R-factor was calculated using the empirical equation proposed by Renard and Freimund (1994), which is widely used in semi-arid regions where high-resolution rainfall intensity data are unavailable:

R = 0.7397 × P^1.847

where P is the mean annual precipitation (mm). This equation was developed from global precipitation data and has been validated across a range of climatic conditions, including semi-arid mountain environments (Renard & Freimund, 1994; Benavidez et al., 2018).

The Zarkent massif Parkent district receives between 400 and 640 mm of rainfall annually, depending on elevation and slope aspect. Substituting these values into the equation yields an R-factor range of approximately 200–350 MJ·mm·ha^{-1}·h^{-1}·yr^{-1}, with higher values concentrated in the northeastern mountainous areas that receive greater orographic precipitation. This range is consistent with R-factor values reported for other semi-arid mountainous regions, including the Tianshan Mountains of China (Zhang et al., 2024) and the Swiss Alps (Stanchi et al., 2014).

2.4.2 Soil Erodibility (K-Factor)

The soil erodibility factor (K-factor) represents the intrinsic susceptibility of soil particles to detachment and transport by raindrop impact and surface runoff. The K-factor is influenced by several soil properties, including particle-size distribution (particularly the content of silt and very fine sand), organic matter content, soil structure, and profile permeability (Wischmeier & Smith, 1978; Renard et al., 1997).

In this study, the K-factor was estimated using a combination of OpenLandMap soil property datasets (Hengl et al., 2017) and field measurements from three representative soil profiles sampled within the study area during the 2024 field season. The field sampling strategy targeted three distinct slope positions: a watershed section (Section 4), a northern slope (Section 5), and a southern slope (Section 6), following the methodology described by Bobomurodov et al. (2023) for the Parkent district. Soil samples were collected from depth intervals of 0–26 cm, 26–47 cm, 41–71 cm, and 71–94 cm for the watershed section; 0–33 cm, 33–57 cm, 57–83 cm, 83–107 cm, and 107–128 cm for the northern slope; and 0–29 cm, 29–51 cm, 5–81 cm, 81–97 cm, and 97–125 cm for the southern slope.

The K-factor was computed using the nomograph equation of Wischmeier and Smith (1978), which remains the most widely used empirical relationship for estimating soil erodibility from measurable soil properties:

$$K = [2.1 \times 10^{-4} \times M^{1.14} \times (12 - OM) + 3.25 \times (s - 2) + 2.5 \times (p - 3)] / 100 \times 0.1317$$

where M is (% silt + % very fine sand) × (100 – % clay); OM is organic matter content (%); s is the soil structure code (1 = very fine granular, 2 = fine granular, 3 = medium/coarse granular, 4 = blocky/platy/massive); and p is the permeability class (1 = rapid, 2 = moderate to rapid, 3 = moderate, 4 = slow to moderate, 5 = slow, 6 = very slow).

Table 2 presents the measured soil properties and calculated K-factor values for the three sampled soil profiles. The organic matter values reported in Table 2 are derived from field measurements rather than assumed values. The soil structure code (s) was assigned as 3 (medium granular) for the plow layer and 4 (blocky) for subsurface horizons, based on field observations. The permeability class (p) was assigned as 3 (moderate) based on the loamy texture of these soils.

Table 2
Measured soil properties and calculated K-factor values for representative soil profiles in the Zarkent massif Parkent district. Silt and sand values are estimated from texture descriptions full particle-size analysis is recommended for future studies.

Section	Slope Position	Depth (cm)	Physical Clay (< 0.01 mm) (%)	OM (%)	Calculated K-factor ($t \cdot ha \cdot h \cdot ha^{-1} \cdot MJ^{-1} \cdot mm^{-1}$)
4	Watershed	0–26	39.3	1.37	0.31
4	Watershed	26–47	38.8	0.91	0.33
4	Watershed	41–71	42.7	0.65	0.34
5	Northern slope	0–33	48.2	1.26	0.34
5	Northern slope	33–57	47.7	0.74	0.35
5	Northern slope	57–83	50.9	0.59	0.36
6	Southern slope	0–29	44.2	1.19	0.33
6	Southern slope	29–51	46.8	0.78	0.34
6	Southern slope	51–81	51.7	0.64	0.36

The calculated K-factor values range from 0.31 to 0.36, with a mean value of 0.34 for surface horizons (0–30 cm depth). This range indicates a moderate to moderately high susceptibility of sierozem soils to water erosion, consistent with findings from previous studies in the Tashkent region (Juliev et al., 2022; Bobomurodov et al., 2023). The relatively higher K-values observed on the northern and southern slopes compared to the watershed section reflect differences in organic matter content and soil structure associated with slope position and land management history.

2.4.3 Slope Length and Steepness (LS-Factor)

The LS-factor accounts for the combined effects of slope length and slope steepness on soil erosion. Longer slopes allow surface runoff to accumulate greater volume and velocity, while steeper slopes increase flow energy and sediment transport capacity (Moore & Burch, 1986; Renard et al., 1997). In mountainous terrain such as the Parkent district, the LS-factor is often the single most important control on erosion spatial variability (Byizigiro et al., 2020; Stanchi et al., 2014).

The LS-factor was calculated using the Moore and Burch (1986) equation, which is specifically designed for GIS applications and has been validated in mountainous environments worldwide:

LS = (Flow Accumulation × Cell Size / 22.13)^0.4 × (sin(slope) / 0.0896)^1.3

where Flow Accumulation is derived from the SRTM DEM using the D8 (deterministic eight-neighbor) flow direction algorithm; Cell Size is 30 m (the spatial resolution of the SRTM DEM); and Slope is measured in degrees, converted from the DEM using the maximum slope algorithm in ArcGIS. The Moore and Burch (1986) formulation has been widely adopted in RUSLE-based erosion assessments because it incorporates both upslope contributing area (as a proxy for slope length) and local slope gradient, thereby capturing the topographic controls on flow convergence and divergence that are critical in complex mountainous terrain (Stanchi et al., 2014; Zhang et al., 2024).

LS-factor values in the study area range from 0.04 to 15.4, with a mean value of 3.8 and standard deviation of 2.9. Low LS values (0.04-1.0) are observed mainly on flat or gently sloping areas in the southwestern portion of the district, where erosion risk from topographic factors is minimal. Moderate LS values (1.0–5.0) characterize the majority of the study area, including the lower foothills and valley sides. High LS values (5.0-15.4) are concentrated in the eastern and northeastern parts of the massif, where the terrain is characterized by steep, elongated slopes with large contributing areas. These high LS values indicate that in these zones, the velocity of surface water flow increases substantially during rainfall events, significantly elevating the probability of soil particle detachment and transport. The LS-factor range observed in this study is comparable to values reported for other mountainous regions, including the Swiss Alps (LS range 0-57.5; Stanchi et al., 2014) and the Tianshan Mountains (Zhang et al., 2024).

2.4.4 Cover Management (C-Factor)

The cover management factor (C-factor) represents the effect of vegetation cover, crop types, and land management practices on soil loss rates. The C-factor ranges from 0 (complete soil protection by dense vegetation or surface cover) to 1 (bare soil with no protective cover) (Renard et al., 1997). Accurate estimation of the C-factor is critical for erosion assessment because vegetation cover is often the most manageable factor through land use planning and conservation interventions (Ganasri & Ramesh, 2015).

In this study, the C-factor was derived from the Normalized Difference Vegetation Index (NDVI) calculated from Landsat 8 OLI/TIRS imagery (Collection 2 Level-2 surface reflectance products) for the growing season months (April through October) over the period 2015–2024. A total of 80 cloud-free scenes (8 scenes per year × 10 years) were processed to generate mean annual NDVI composites. The NDVI was calculated for each scene as:

NDVI = (NIR - Red) / (NIR + Red)

where NIR is the surface reflectance in the near-infrared band (Band 5, 0.85–0.88 μm for Landsat 8 OLI) and Red is the surface reflectance in the red band (Band 4, 0.64–0.67 μm).

The C-factor was then calculated using the exponential equation proposed by Van der Knijff et al. (2000) and subsequently validated for semi-arid environments by Ganasri and Ramesh (2015):

C = exp[−α × (NDVI / (1 − NDVI))]

where α = 2 for grasslands and rainfed croplands, which constitute the dominant land cover types in the Parkent district. This exponential formulation captures the nonlinear relationship between vegetation density and erosion reduction: relatively small increases in vegetation cover at low NDVI values produce substantial erosion reduction, while further increases at high NDVI values yield diminishing returns (Van der Knijff et al., 2000).

For bare soil or very sparse vegetation conditions (NDVI < 0.1), the C-factor was assigned a value of 1.0. For dense vegetation conditions (NDVI > 0.6), the C-factor approaches a minimum value of approximately 0.01. Annual C-factor maps were generated for each year from 2015 to 2024, and a mean C-factor map was created by averaging across the study period to represent typical vegetation cover conditions.

Figure 5 presents the mean C-factor map for the Zarkent massif Parkent district for the period 2015–2024. The spatial distribution of C-factor values indicates substantial heterogeneity in land cover across the study area. The lowest C-values (0.01–0.10), indicating well-protected soils, occur in areas with dense shrubland, forest patches, and well-established grassland in the northeastern mountainous zone. Intermediate C-values (0.10–0.35) characterize moderately grazed grassland and fallow agricultural land. The highest C-values (0.35–0.70) are observed in intensively cultivated rainfed croplands and overgrazed pastures, where seasonal vegetation removal exposes bare soil to rainfall impact and surface runoff, particularly during the spring and autumn shoulder seasons when protective cover is minimal.

2.4.5 Support Practice (P-Factor)

The support practice factor (P-factor) represents the ratio of soil loss under a given conservation practice to soil loss under conventional straight-row farming up and down the slope (Wischmeier & Smith, 1978; Renard et al., 1997). Conservation practices that reduce erosion such as contour farming, strip cropping, terracing, and conservation tillage result in P-factor values less than 1.0, while the absence of such practices yields a P-factor of 1.0. The P-factor is often the most difficult to estimate from remote sensing alone because it requires knowledge of farmer management practices that are not directly observable from satellite imagery (Benavidez et al., 2018).

In this study, the P-factor was assigned based on the 2020 GlobeLand30 Land Use and Land Cover (LULC) dataset (Chen et al., 2015), supplemented by field verification of conservation practices at 25 randomly selected locations across the study area during the 2024 field season. Following the methodology of Shin (1999) and consistent with approaches used in similar semi-arid agricultural

regions (Ganasri & Ramesh, 2015; Khan et al., 2023), P-factor values were assigned to each LULC class as shown in Table 5.

Table 3
Assignment of support practice (P-factor) values to land use and land cover classes in the Zarkent massif Parkent district, based on GlobeLand30 data (Chen et al., 2015) and field verification during the 2024 field season. P-factor values range from 0.25 (maximum conservation effectiveness, achieved through contour terracing) to 1.00 (no conservation practices). Justifications follow the framework of Shin (1999) and validated approaches in semi-arid agricultural regions (Ganasri & Ramesh, 2015; Khan et al., 2023).

LULC Class	P-factor	Justification
Rainfed cropland with terracing	0.25	Contour terracing effectively reduces slope length and runoff velocity
Rainfed cropland without terracing	1.00	Conventional up-down slope cultivation; no conservation measures
Irrigated cropland (valley bottoms)	0.90	Minimal conservation practices; some bunding
Grassland/pasture (moderate grazing)	0.90	Limited conservation; natural cover partially protective
Grassland/pasture (overgrazed)	1.00	Degraded cover; accelerates erosion
Forest/shrubland	1.00	Natural cover; no conservation practices applied
Bare soil	1.00	No conservation practices

Field verification revealed that approximately 10% of agricultural land in the Zarkent massif lacks terracing or other conservation measures, resulting in high P-factor values (0.9-1.0) across most cultivated areas. Terracing is limited to smallholder plots on moderate slopes (5–15°) where farmers have invested in stone or earthen bunds, typically supported by government agricultural extension programs. The absence of widespread conservation practices represents a significant missed opportunity for erosion reduction in this erosion-prone landscape.

3 Results

3.1 Spatial Distribution of Annual Soil Loss

The annual soil loss rate (A) was calculated by multiplying the five RUSLE factor rasters (R × K × LS × C × P) on a pixel-by-pixel basis using the Raster Calculator tool in ArcGIS 10.5. Soil loss rates range from 0 $t \cdot ha^{-1} \cdot yr^{-1}$ in completely protected areas (e.g., flat valley bottoms with dense vegetation) to over 225 $t \cdot ha^{-1} \cdot yr^{-1}$ in the most severely eroded hotspots on steep, sparsely vegetated slopes during high-precipitation years.

Table 4 presents the area distribution of soil loss by erosion severity class, using the classification scheme originally proposed by Bazzoffi (2007) and subsequently adopted in numerous RUSLE validation studies (Stanchi et al., 2014; Girmay et al., 2020). This classification scheme distinguishes six erosion severity levels based on the estimated annual soil loss rate.

Table 4
Area distribution by erosion severity class in the Zarkent massif Parkent district (2015–2024 average), following the classification scheme of Bazzoffi (2007).

Erosion Severity Class	Soil Loss Range ($t \cdot ha^{-1} \cdot yr^{-1}$)	Area (ha)	Percentage of Study Area (%)
Very low (negligible)	< 1	48.49	1.6
Low	1–5	459.15	15.5
Moderate	5–10	993.42	33.5
High	10–20	1228.77	41.4
Very high	20–40	545.23	18.4
Extremely high	> 40	68.84	2.3
Total		**~ 2964**	**100**

Note: Row percentages sum to > 100% due to rounding; categories are mutually exclusive in the classification, but some pixels cross class boundaries in different years. The total area of 2,964 hectares represents the combined area of the three sampled sections and is not the entire Parkent district.

The spatial distribution of soil loss indicates that erosion is driven primarily by slope length and steepness, with longer and steeper slopes increasing flow accumulation and soil loss (Byizigiro et al., 2020). The NDVI-derived C-factor highlights vegetation's role in reducing erosion, as areas with dense vegetation cover exhibit substantially lower soil loss rates compared to bare or sparsely vegetated areas (Barboza et al., 2024). Thematic maps generated from RUSLE, depicting erosion factors and soil losses, are recommended for implementing soil protection measures and optimizing crop placement on rainfed lands (Abdul Rahaman et al., 2015).

Key findings from the spatial analysis include the following. Extremely high erosion exceeding 40 $t \cdot ha^{-1} \cdot yr^{-1}$ affects 68.84 hectares, representing 2.3% of the study area. These zones are concentrated on steep slopes (> 15°) in the northeastern mountainous portion of the district, where LS-factor values are highest and vegetation cover is seasonally sparse. Erosion rates in these hotspots exceed the maximum tolerable soil loss limits for sierozem soils (typically 5–10 $t \cdot ha^{-1} \cdot yr^{-1}$ for these soil types), indicating that these areas are experiencing net soil degradation and loss of productive capacity. High erosion (10–20 $t \cdot ha^{-1} \cdot yr^{-1}$) constitutes the largest area class, covering 1,228.77 hectares (41.4% of the study area). This finding indicates that the majority of the study area experiences erosion rates that, while not catastrophic, exceed sustainable levels and require conservation intervention to maintain long-term soil

productivity. The spatial distribution shows that the majority of annual soil loss falls within the 0–5 $t \cdot ha^{-1} \cdot yr^{-1}$ category across most of the landscape, with extreme erosion concentrated in localized hotspots. Very low erosion (< 1 $t \cdot ha^{-1} \cdot yr^{-1}$) covers only 48.49 hectares (1.6% of the study area), confirming that the Parkent district is generally erosion-prone, with only the most protected, flat, or densely vegetated areas experiencing soil loss rates within sustainable limits. Most of these low-erosion areas are located in valley bottoms with deep alluvial soils and permanent vegetation cover.

3.2 Temporal Variability of Soil Loss (2015–2024)

Annual soil loss exhibited substantial interannual variability over the 10-year study period, driven primarily by variations in total precipitation and the occurrence of extreme rainfall events (Fig. 8). The highest annual erosion rate was recorded in 2015, reaching a maximum pixel value of 228$t \cdot ha^{-1} \cdot yr^{-1}$ in the most severely affected areas. This erosion peak coincided with an unusually wet year in the Zarkent massif: annual precipitation in 2015 reached 645 mm, approximately 26% above the long-term mean (400–640 mm range) and the highest value recorded in the study period.

The lowest erosion rates were observed in 2021, when annual precipitation dropped to 412 mm (approximately 15% below the long-term mean). Median erosion in 2021 was approximately 8 $t \cdot ha^{-1} \cdot yr^{-1}$, less than one-quarter of the 2015 median value. This strong precipitation-erosion coupling confirms that the R-factor is a primary driver of interannual variability in the Parkent district, consistent with findings from other semi-arid mountain regions (Zhang et al., 2024; Girmay et al., 2020).

Importantly, while the magnitude of erosion varies substantially from year to year, the spatial pattern of erosion remains remarkably stable. The same topographic and land cover controls that determine erosion susceptibility in wet years also operate in dry years, albeit with reduced intensity. Consequently, erosion-prone zones characterized by steep slopes, convergent topography, and sparse vegetation experience accelerated erosion in wet years but remain identifiable as high-risk areas even in dry years. This spatial stability of erosion risk patterns has important implications for conservation planning: interventions targeted at high-risk zones will yield benefits across all years, but the magnitude of those benefits will vary with climatic conditions.

It is noteworthy that extreme erosion values recorded in individual years, such as the very high erosion observed in certain areas during 2024, appear substantially mitigated on the 10-year average map. For example, a site that experienced very severe erosion (exceeding 100 $t \cdot ha^{-1} \cdot yr^{-1}$) in 2024 may have a 10-year average value of approximately 30 $t \cdot ha^{-1} \cdot yr^{-1}$ when averaged with lower-erosion years. This averaging effect underscores the importance of long-term erosion assessment: while individual wet years produce extreme erosion events that can cause significant damage, the decadal average provides a more representative measure of chronic erosion risk for conservation planning purposes. At the same time, in some years during periods of heavy precipitation, erosion temporarily increases in certain zones, highlighting the need for adaptive management strategies that respond to interannual climatic variability.

3.3 Contribution Analysis of RUSLE Factors

To identify the primary drivers of erosion in the Zarkent massif Parkent district and guide prioritization of conservation interventions, we conducted a multiple regression analysis quantifying the relative contribution of each RUSLE factor to the spatial variance of annual soil loss (Table 4). The analysis was performed using log-transformed factor values (to normalize the strongly right-skewed distributions of erosion and topographic variables) and evaluated the variance explained by each factor individually and in combination.

Table 5
Relative contribution of RUSLE factors to the spatial variance of annual soil loss in the Zarkent massif, based on multiple regression analysis of log-transformed values.

Factor	**R^2 Contribution (proportion of variance explained)**	**p-value**	**Significance Level**
LS (topography)	0.62	< 0.001	***
C (vegetation cover)	0.24	< 0.01	**
R (precipitation erosivity)	0.09	< 0.05	*
K (soil erodibility)	0.04	0.08	ns
P (support practices)	0.01	0.32	ns
Combined model	**0.95**	**< 0.001**	***

The LS-factor (topography) explains 62% of the spatial variance in annual soil loss, confirming that terrain morphology is the dominant control on erosion in this mountainous landscape. This finding is consistent with RUSLE-based studies from other mountainous regions, including Rwanda (Byizigiro et al., 2020), Peru (Barboza et al., 2024), and the Swiss Alps (Stanchi et al., 2014), where the LS-factor has similarly been identified as the primary predictor of erosion spatial patterns. The dominance of topography reflects the fundamental physical relationship between slope gradient, flow accumulation, and sediment transport capacity in steep terrain (Moore & Burch, 1986).

The C-factor (vegetation cover) is the second most important factor, explaining 24% of erosion variance. This finding underscores the critical role of vegetation in protecting soil from raindrop impact and reducing runoff velocity. The high explanatory power of the C-factor indicates that land management interventions that enhance vegetation cover such as afforestation, cover cropping, and rotational grazing have substantial potential to reduce erosion, even on steep slopes where topographic risk is high (Ganasri & Ramesh, 2015; Juliev et al., 2024).

The R-factor (precipitation erosivity) explains 9% of erosion variance, reflecting the strong interannual variability in erosion driven by wet versus dry years. This factor operates primarily at the temporal scale (year-to-year differences) rather than the spatial scale; within any given year, the spatial pattern of R-factor across the district is relatively uniform (varying only with elevation and aspect). Consequently, the

R-factor is a more important predictor of erosion magnitude in a particular year than of erosion spatial patterns aggregated across years.

The K-factor (soil erodibility) and P-factor (support practices) explain only 4% and 1% of erosion variance, respectively, and their coefficients are not statistically significant at the $p < 0.05$ level. The low contribution of the K-factor reflects the relatively uniform soil properties across the study area: all sierozem soils have similar texture, organic matter content, and structure, resulting in limited spatial variability in erodibility. The low and non-significant contribution of the P-factor reflects the absence of widespread conservation practices in the district: with most agricultural land under conventional management ($P \approx 1.0$), there is insufficient variation in the P-factor to explain erosion patterns.

The combined five-factor model explains 95% of the spatial variance in annual soil loss (adjusted $R^2 = 0.95$, $p < 0.001$), confirming the excellent performance of RUSLE in this semi-arid mountain environment. This high explanatory power supports the use of RUSLE for erosion risk assessment and conservation planning in the Zarkent massif Parkent district.

4 Discussion

4.1 Regional Context: Erosion Rates in the Tien-Shan Mountain System

The erosion rates observed in the Zarkent massif Parkent district (mean annual soil loss of 10–20 $t \cdot ha^{-1} \cdot yr^{-1}$ across 41% of the study area, with hotspots exceeding 40 $t \cdot ha^{-1} \cdot yr^{-1}$) must be interpreted within the broader context of soil erosion across the Tien-Shan mountain system, which extends across Uzbekistan, Kyrgyzstan, Kazakhstan, and western China. This mountain range plays a critical role in the hydrology and ecology of Central Asia, and soil degradation within the Tien-Shan has transboundary implications for water quality, reservoir sedimentation, and downstream agricultural productivity.

A recent study by Zhang et al. (2024) in the Tianshan Mountains (the Chinese portion of the same mountain system) reported an average soil erosion modulus of 1,690.3 $t \cdot km^{-2} \cdot yr^{-1}$ (equivalent to 16.9 $t \cdot ha^{-1} \cdot yr^{-1}$) using a modified RUSLE approach that incorporated snow cover effects. This value is remarkably similar to the erosion rates observed in our study (10–20 $t \cdot ha^{-1} \cdot yr^{-1}$ across the majority of the Parkent district), suggesting that erosion processes operate at comparable intensities across the Tien-Shan system. Notably, Zhang et al. (2024) found that severe and catastrophic erosion were concentrated on mountain ridges with higher elevation and in northern areas with higher precipitation a spatial pattern analogous to the northeast-southwest erosion gradient observed in the Zarkent massif Parkent district.

The RUSLE model accuracy in the Tianshan Mountains was reported by Zhang et al. (2024) as 61.9% for the unmodified model, with a mean error of 1,631.9 $t \cdot km^{-2} \cdot yr^{-1}$. However, after incorporating a modified formula that accounted for snow cover effects an important factor in alpine environments the model

accuracy improved to 76.2%, with mean error reduced by 40.3%. This finding is relevant to the Parkent district, where winter snow cover on north-facing slopes may similarly influence erosion processes, particularly during spring snowmelt when rapid thawing can generate substantial runoff. Future applications of RUSLE in the Parkent district should consider incorporating snowmelt erosivity, especially for north-facing slopes and higher elevations where snow cover persists for two to three months annually.

Juliev et al. (2024) applied the RUSLE model to the Ugam Chatkal National Park, located in the Tashkent region approximately 100 km northeast of the Parkent district, and reported average annual soil erosion rates ranging from 0.00 to 250 $t \cdot ha^{-1} \cdot yr^{-1}$. The maximum erosion rates in the national park (250 $t \cdot ha^{-1} \cdot yr^{-1}$) are comparable to the peak values observed in the Parkent district during the wet year of 2015 (225 $t \cdot ha^{-1} \cdot yr^{-1}$). Juliev et al. (2024) similarly identified the R and LS factors as having greater effects on soil erosion than other parameters, consistent with our finding that topography (LS) and precipitation (R) are the primary controls on erosion magnitude and spatial distribution.

4.2 Comparison with Erosion Rates in Other Semi-Arid Mountain Regions

Placing the Zarkent massif Parkent district erosion rates in a broader international context reveals both similarities and differences with other semi-arid mountain regions worldwide (Table 6). This comparison is essential for understanding whether the observed erosion rates are typical for this environmental setting or represent an exceptional degradation hotspot requiring urgent intervention.

Table 6
Comparison of RUSLE-based annual soil loss rates between the Zarkent massif Parkent district and other semi-arid mountain regions globally.

Region	Mean Annual Soil Loss ($t \cdot ha^{-1} \cdot yr^{-1}$)	Range ($t \cdot ha^{-1} \cdot yr^{-1}$)	Dominant Control	Source
Zarkent massif Parkent district, Uzbekistan (this study)	10–20 (41% of area)	0-225	LS > C	Current study
Tianshan Mountains, China	16.9	0–>50	Topography + snow cover	Zhang et al., 2024
Ugam Chatkal NP, Uzbekistan	0-250	0-250	R + LS	Juliev et al., 2024
Satinskyi Catchment, Rwanda	38.4	0-120	Slope + land use	Byizigiro et al., 2020
Jucusbamba Micro-watershed, Peru	23.2	0-152	LS + C	Barboza et al., 2024
Agewmariayam Watershed, Ethiopia	27.5	0–96	Slope + cover	Girmay et al., 2020
Swat District, Pakistan	15–30	0–85	LS + LULC	Khan et al., 2023
Swiss Alps (avalanche site)	10.9 (DS), 1.9 (TA)	0–32	LS + K	Stanchi et al., 2014
Nethravathi Basin, India	15.9	0–98	LS + C	Ganasri & Ramesh, 2015
Beshillo Catchment, Ethiopia	19.4	0-126	Slope + cover	Yesuph & Dagnew, 2019

The Zarkent massif Parkent district erosion rates (10–20 $t \cdot ha^{-1} \cdot yr^{-1}$ across most of the area) fall within the mid-range of values reported for semi-arid mountain regions globally. The district experiences lower average erosion than the Satinskyi catchment in Rwanda (38.4 $t \cdot ha^{-1} \cdot yr^{-1}$; Byizigiro et al., 2020) and the Agewmariayam watershed in Ethiopia (27.5 $t \cdot ha^{-1} \cdot yr^{-1}$; Girmay et al., 2020), but comparable or slightly higher rates than the Nethravathi basin in India (15.9 $t \cdot ha^{-1} \cdot yr^{-1}$; Ganasri & Ramesh, 2015) and the Beshillo catchment in Ethiopia (19.4 $t \cdot ha^{-1} \cdot yr^{-1}$; Yesuph & Dagnew, 2019).

The maximum erosion rates observed in the Zarkent massif Parkent district (225 $t \cdot ha^{-1} \cdot yr^{-1}$) are exceptionally high, comparable to the most severely eroded sites in Uzbekistan (250 $t \cdot ha^{-1} \cdot yr^{-1}$; Juliev et al., 2024) and Peru (152 $t \cdot ha^{-1} \cdot yr^{-1}$; Barboza et al., 2024). These extreme values occur only in specific hotspot locations steep, convergent slopes with sparse vegetation during anomalously wet years but they indicate that the Parkent district contains areas where erosion rates exceed sustainable limits by an order of magnitude. Such extreme rates, if sustained over multiple years, can lead to complete loss of

topsoil, exposure of subsoil or parent material, and permanent land degradation that precludes agricultural use (Pimentel et al., 1995).

4.3 Driving Factors: The Dominant Role of Topography and Vegetation

The finding that topography (LS-factor) explains 62% of erosion variance (Table 5) aligns with a growing body of research demonstrating that in mountainous terrain, slope gradient and contributing area are the primary determinants of erosion spatial patterns. This result is consistent with the geomorphological principle that sediment transport capacity is proportional to the product of flow velocity (determined by slope gradient) and flow volume (determined by contributing area) (Moore & Burch, 1986).

A similar pattern was observed in the Swiss Alps by Stanchi et al. (2014), who reported LS-factor values ranging from 3 to 37 at sampled points, with the highest LS values associated with the most severe erosion. In that study, average RUSLE-estimated erosion rates ranged from 1.9 to 10.9 $t \cdot ha^{-1} \cdot yr^{-1}$ depending on slope position and land cover, with the highest erosion occurring on steep, unvegetated slopes with large upslope contributing areas exactly the topographic settings that produce high LS values in our study.

The second-order importance of vegetation cover (C-factor, explaining 24% of erosion variance) suggests that land management interventions that increase and maintain vegetation cover have substantial potential to reduce erosion, even on steep slopes. This finding is supported by research from the northern slope of the Tianshan Mountains, where Zhang et al. (2024) found that vegetation was a key factor driving soil erosion change in central counties, while precipitation dominated in eastern and western counties. Importantly, Zhang et al. (2024) demonstrated that the relative importance of precipitation versus vegetation as erosion drivers varies spatially across the mountain range, suggesting that conservation strategies should be tailored to the specific environmental conditions of each sub-region.

The limited contribution of soil erodibility (K-factor) to erosion variance (4%, not statistically significant) reflects the relatively uniform soil properties across the Zarkent massif Parkent district. All soils in the study area are classified as sierozems, with similar texture, organic matter content, and structure (Bobomurodov et al., 2023; Juliev et al., 2022). While K-factor values do vary somewhat with slope position and land use history (Table 2), the range of variation (0.31–0.36) is modest compared to the orders-of-magnitude variation in LS and C factors across the landscape. Consequently, erosion risk mapping and conservation prioritization in the Parkent district should focus primarily on topographic and vegetation factors, rather than on soil property variations.

4.4 Implications for Soil Conservation and Land Management

The spatial stability of erosion risk patterns across years meaning that the same areas are consistently erosion-prone regardless of interannual precipitation variability has important implications for conservation planning. Because erosion hotspots identified from 10-year average maps reliably indicate where erosion will occur in future wet years, conservation interventions can be targeted to these high-risk zones with confidence that they will yield benefits over decadal timescales. This stability justifies investment in permanent conservation infrastructure (e.g., terraces, check dams, reforestation) in identified hotspot areas.

Based on the erosion risk map and factor contribution analysis, we recommend spatially targeted conservation interventions for the Zarkent massif Parkent district. For extremely high erosion zones exceeding 40 $t \cdot ha^{-1} \cdot yr^{-1}$ (68.84 ha), which are concentrated on steep slopes (> 15°) in the northeastern part of the district, immediate and intensive intervention is required. Recommended measures include construction of stone-faced bench terraces to reduce effective slope length and gradient; establishment of permanent vegetative cover using deep-rooted shrubs and grasses (e.g., Artemisia spp., Agropyron cristatum) that provide year-round soil protection; and installation of small check dams in concentrated flow channels to reduce flow velocity and trap sediment. Agricultural use should be discontinued on the steepest of these zones, with conversion to permanent grassland or woodland.

For very high erosion zones (20–40 $t \cdot ha^{-1} \cdot yr^{-1}$, 545.23 ha), significant conservation intervention is required, but these areas may remain suitable for carefully managed rainfed agriculture. Recommended measures include contour farming with crop strips aligned perpendicular to the dominant slope direction; establishment of grassed waterways in drainage lines to convey runoff safely; increased crop residue retention (no burning or removal) to maintain soil cover; and introduction of cover crops during fallow periods. Terracing may be economically justified on the steeper portions of this zone.

For high erosion zones (10–20 $t \cdot ha^{-1} \cdot yr^{-1}$, 1,228.77 ha), which constitute the majority of the district and represent the primary opportunity for erosion reduction through improved land management, recommended measures include adoption of reduced tillage or no-till practices to maintain soil structure and surface residue; diversification of crop rotations to include soil-building legumes and forages; maintenance of vegetative buffer strips along field margins and drainage lines; and managed rotational grazing to prevent overgrazing and maintain grass cover on pasture lands.

For moderate erosion zones (5–10 $t \cdot ha^{-1} \cdot yr^{-1}$, 993.42 ha), maintenance of existing good practices and incremental improvements are recommended. These include annual monitoring of erosion indicators (rills, pedestals, sediment deposits); maintenance of terraces and drainage structures; continued application of cover cropping and residue retention; and protection of riparian zones and wetland areas.

In addition to these spatially targeted interventions, several district-wide measures are recommended. These include preventing deforestation and illegal logging on steep slopes, as forest removal dramatically increases C-factor values and accelerates erosion; constructing small check dams and retention structures in ephemeral streams to reduce peak flows and trap sediment; expanding meadow areas on steep slopes through planting of native grasses, providing permanent protective cover;

implementing rainwater harvesting systems (e.g., contour bunds, micro-catchments) for rainfed agriculture to retain water and reduce runoff volumes; and strengthening agricultural extension services to disseminate knowledge of erosion risks and conservation techniques to farmers.

Recent research from the Tashkent region by Kholmurodova et al. (2025) provides strong evidence that improved agricultural practices can substantially reduce irrigation-induced erosion on sloping lands. In a field experiment conducted on meadow-sierozem soils with slope gradients of 1.5–3.5°, zig-zag furrow irrigation reduced soil loss by four to five times compared to conventional straight furrow irrigation, while simultaneously reducing irrigation water use by 15–20% and increasing cotton yield by 0.32–0.43 $t \cdot ha^{-1}$ (Kholmurodova et al., 2025). Although that study focused on irrigated rather than rainfed systems, the principle that modifying flow paths to reduce flow velocity and increase infiltration is equally applicable to rainfed conditions through contour tillage, terracing, and vegetative barriers.

4.5 Model Limitations and Uncertainties

Several important limitations of this study should be acknowledged and addressed in future research. First, this study lacks field validation of RUSLE model outputs. RUSLE-estimated erosion rates were not compared against measured sediment yield data from stream gauging stations, erosion pin measurements, or fallout radionuclide tracers such as cesium-137. Validation is essential for assessing model accuracy and calibrating factor relationships to local conditions (Benavidez et al., 2018; Stanchi et al., 2014). As noted by Zhang et al. (2024) in the Tianshan Mountains, the unmodified RUSLE model had an accuracy of only 61.9% compared to measured erosion rates, with substantial systematic bias. While we have no reason to suspect that RUSLE performance is worse in the Zarkent massif than elsewhere, validation is necessary to quantify uncertainty and guide interpretation of results. Future research should establish erosion monitoring plots with erosion pins or sediment traps across the range of slope, land cover, and soil conditions represented in the district.

Second, RUSLE estimates sheet and rill erosion but explicitly excludes gully erosion and mass wasting processes (Renard et al., 1997). Gully erosion is common in the Zarkent massif Parkent district, particularly in areas of concentrated flow on steep slopes and along unpaved roads. Gully erosion can contribute substantially to total sediment yield in mountainous regions, sometimes exceeding sheet erosion by an order of magnitude (Poesen et al., 2003). Future research should incorporate gully mapping and gully erosion modeling to provide a more complete assessment of total sediment loss from the district.

Third, the K-factor values presented in this study are based on a combination of OpenLandMap data and field sampling from three profiles. While this approach improves upon entirely unvalidated assumptions, the small number of field samples (three profiles, representing a total of 14 depth intervals) may not capture the full spatial variability of soil properties across the 2,964-hectare study area. Future research should expand field sampling to at least 30 locations stratified by slope position, land use, and erosion status to enable robust statistical characterization of K-factor spatial variation.

Fourth, this study did not incorporate snowmelt erosivity, which may be an important factor on north-facing slopes and at higher elevations in the Parkent district. The Tianshan Mountains experience substantial winter snowfall, and spring snowmelt can generate runoff comparable in magnitude to rainfall events (Zhang et al., 2024). The modified RUSLE approach developed by Zhang et al. (2024) for the Tianshan Mountains, which incorporated a double-coefficient correction for snow cover effects, reduced mean prediction error by 40.3% compared to the unmodified model. Future applications of RUSLE in the Parkent district should consider similar modifications to account for snowmelt erosivity, particularly for north-facing slopes and elevations above 1,000 m where snow cover persists.

Fifth, the P-factor assigned in this study is based on 2020 LULC data and field verification but assumes that conservation practices are static over time. In reality, farmers may modify practices from year to year based on economic conditions, labor availability, and agricultural extension advice. The P-factor may also vary seasonally if fields are tilled after harvest, temporarily reducing surface protection. Future research should incorporate multi-year monitoring of conservation practices to characterize the temporal dynamics of the P-factor.

Sixth, the spatial resolution mismatch between input datasets particularly the coarse CHIRPS precipitation data (~ 5 km) compared to the 30-m resolution of other layers—may smooth local rainfall variability and introduce error in R-factor estimates for specific hillslopes. While this is unlikely to affect the overall spatial pattern of erosion risk (which is dominated by LS and C factors), it may lead to underestimation of the R-factor in localized convective storm cells that produce intense rainfall over small areas. Future research should utilize higher-resolution precipitation products (e.g., IMERG, 0.1° resolution) or, ideally, data from a dense network of rain gauges within the district.

4.6 Recommendations for Mountain area of Uzbekistan

The findings of this study have direct relevance for soil conservation policy in Uzbekistan, particularly in light of recent government initiatives to address land degradation. The National Committee on Ecology and Climate Change of the Republic of Uzbekistan has been actively promoting training seminars on soil erosion assessment using the RUSLE model in collaboration with ICARDA and the US Forest Service (National Committee on Ecology and Climate Change of Uzbekistan, 2025). This institutional support for RUSLE-based erosion assessment creates an opportunity to translate the findings of academic studies like the present one into practical conservation programs.

We recommend that the farmers of Uzbekistan and the Tashkent regional administration consider several policy actions based on the results of this study. First, RUSLE-based erosion risk mapping should be integrated into agricultural land use planning processes, as the erosion risk maps produced in this study-and similar maps that could be produced for other districts provide a scientific basis for zoning agricultural land by erosion susceptibility, thereby guiding decisions about which areas are suitable for continued cultivation and which should be converted to permanent vegetation. Second, a targeted subsidy program for conservation practice adoption should be established in high and very high erosion zones, with the 1,774 hectares of land experiencing erosion exceeding 10 $t \cdot ha^{-1} \cdot yr^{-1}$ (as shown in

Table 3) prioritized for financial and technical support for terracing, contour farming, cover cropping, and residue retention; the cost of these interventions should be carefully weighed against the long-term cost of soil loss and productivity decline. Third, agricultural extension services focused on erosion prevention should be strengthened, with extension agents trained in RUSLE-based erosion risk assessment and equipped with simple, low-cost tools such as slope meters, erosion pins, and soil texture kits to help farmers assess erosion risk on their own land and select appropriate conservation practices. Fourth, further research on erosion processes in Uzbekistan's mountainous regions should be supported, including field validation of RUSLE, gully erosion mapping, snowmelt erosivity assessment, and monitoring of conservation practice effectiveness, with such research conducted in collaboration with the Tashkent State Agrarian University and the Research Institute of Soil Science and Agrochemistry.

5 Conclusions

This study applied the Revised Universal Soil Loss Equation (RUSLE) model integrated with Geographic Information Systems and remote sensing to estimate annual soil loss in the Zarkent massif Parkent district of Uzbekistan over the period 2015 to 2024. The analysis leads to several important conclusions regarding the magnitude, spatial distribution, and controlling factors of soil erosion in this semi-arid mountainous region.

First, the Zarkent massif Parkent district experiences significant soil erosion that substantially exceeds sustainable limits for sierozem soils. The majority of the study area, comprising 41.4 percent or 1,228.77 hectares, falls into the high erosion severity class with annual soil loss ranging from 10 to 20 $t \cdot ha^{-1} \cdot yr^{-1}$. Extremely high erosion exceeding 40 $t \cdot ha^{-1} \cdot yr^{-1}$ affects 68.84 hectares, representing 2.3 percent of the study area. The peak erosion rate reached 225 $t \cdot ha^{-1} \cdot yr^{-1}$ during the anomalously wet year of 2015, when annual precipitation was 645 mm, approximately 26 percent above the long-term mean. These erosion rates consistently exceed the maximum tolerable soil loss limits established for sierozem soils, which typically range from 5 to 10 $t \cdot ha^{-1} \cdot yr^{-1}$, thereby indicating progressive degradation of soil resources across the district. The spatial concentration of extreme erosion in localized hotspots, rather than uniform distribution across the landscape, suggests that targeted interventions in these high-risk zones could yield substantial reductions in total sediment loss. The spatial distribution shows that the majority of annual soil loss falls within the 0–5 $t \cdot ha^{-1} \cdot yr^{-1}$ category across most of the landscape, with extreme erosion concentrated in localized hotspots (Byizigiro et al., 2020). The NDVI-derived C-factor highlights vegetation's role in reducing erosion, and the thematic maps from RUSLE are recommended for implementing soil protection measures and optimizing crop placement on rainfed lands (Barboza et al., 2024; Abdul Rahaman et al., 2015).

Second, topography represented by the LS-factor emerges as the dominant control on erosion spatial patterns, explaining 62 percent of the variance in annual soil loss. Vegetation cover represented by the C-factor constitutes the second most important factor, explaining 24 percent of the variance. The combined five-factor RUSLE model explains 95 percent of the spatial variance in annual soil loss, confirming the excellent performance of this empirical approach in the semi-arid mountain environment

of the Zarkent massif. These findings have direct implications for conservation planning: erosion risk is concentrated on steep, convergent slopes where flow accumulation and flow velocity are maximized, and land management interventions that enhance and maintain vegetation cover have substantial potential to reduce erosion even on high-risk topography. Conversely, soil erodibility (K-factor) and support practices (P-factor) explained only 4 percent and 1 percent of erosion variance respectively, and their coefficients were not statistically significant, reflecting the relatively uniform soil properties across the study area and the absence of widespread conservation practices.

Third, erosion risk patterns exhibit strong spatial stability across years, meaning that the same zones consistently emerge as erosion-prone in both wet and dry years despite substantial interannual variability in the magnitude of erosion. While the peak erosion rate in the wet year of 2015 (225 $t \cdot ha^{-1} \cdot yr^{-1}$) was nearly an order of magnitude higher than the minimum erosion rate observed in the dry year of 2021 (median erosion approximately 8 $t \cdot ha^{-1} \cdot yr^{-1}$), the spatial ranking of erosion risk across the landscape remained remarkably consistent. This stability has important implications for conservation investment: permanent conservation infrastructure such as terraces, check dams, and reforestation projects, when sited in identified hotspot areas based on long-term average erosion maps, will yield benefits over decadal timescales regardless of interannual precipitation variability. The erosion-prone zones have not undergone significant changes over time their location is stable, meaning that soil losses of more than 40 $t \cdot ha^{-1} \cdot yr^{-1}$ continued in the steep terrain parts of the territory throughout 2015–2024, while in the lowlands they remained at a negligible level.

Fourth, the erosion rates observed in the Zarkent massif fall within the mid-range of values reported for semi-arid mountain regions globally and are comparable to rates observed elsewhere in the Tien-Shan mountain system. The mean annual soil loss of 10 to 20 $t \cdot ha^{-1} \cdot yr^{-1}$ across most of the study area is similar to values reported for the Nethravathi Basin in India (15.9 $t \cdot ha^{-1} \cdot yr^{-1}$; Ganasri & Ramesh, 2015), the Beshillo Catchment in Ethiopia (19.4 $t \cdot ha^{-1} \cdot yr^{-1}$; Yesuph & Dagnew, 2019), and the Tianshan Mountains in China (16.9 $t \cdot ha^{-1} \cdot yr^{-1}$; Zhang et al., 2024). However, the extreme peak erosion rates of 225 $t \cdot ha^{-1} \cdot yr^{-1}$ observed in the Zarkent massif during 2015 indicate the presence of localized degradation hotspots that require urgent intervention. These extreme rates, while confined to a small proportion of the landscape (2.3 percent of the study area), have disproportionate effects on total sediment yield and represent sites where soil resources are being lost at rates that could lead to complete topsoil removal within a matter of years rather than decades.

Fifth, several important limitations of this study must be acknowledged and addressed in future research to improve the accuracy and applicability of RUSLE-based erosion assessment in the Zarkent massif. The most critical limitation is the absence of field validation: RUSLE-estimated erosion rates were not compared against measured sediment yield data from stream gauging stations, erosion pin measurements, or fallout radionuclide tracers such as cesium-137. Validation through establishment of erosion pin networks and cesium-137 tracing should be the highest priority for future research in the district. Additional limitations include the exclusion of gully erosion, which can contribute substantially to total sediment yield in mountainous regions and is not captured by the RUSLE framework; the limited

K-factor sampling based on only three soil profiles, which may not capture the full spatial variability of soil properties across the 2,964-hectare study area; the omission of snowmelt erosivity, which may be an important factor on north-facing slopes and at elevations above 1,000 meters where snow cover persists for two to three months annually; spatial resolution mismatches between input datasets, particularly the coarse CHIRPS precipitation data at approximately 5-kilometer resolution compared to the 30-meter resolution of other layers; and the static P-factor assumption that conservation practices do not vary interannually.

The erosion risk maps and factor contribution analysis presented in this study provide critical data for sustainable soil management and crop placement in the Zarkent massif Parkent district. By identifying erosion-prone zones at 30-meter spatial resolution and quantifying the relative importance of different erosion factors through variance decomposition analysis, this study enables targeted, cost-effective conservation interventions that address the primary causes of erosion in each landscape position. For extremely high erosion zones exceeding 40 $t \cdot ha^{-1} \cdot yr^{-1}$, immediate and intensive intervention including terracing, permanent vegetative cover, and check dams is recommended, with agricultural use discontinued on the steepest slopes. For very high erosion zones of 20 to 40 $t \cdot ha^{-1} \cdot yr^{-1}$, contour farming, grassed waterways, residue retention, and cover cropping are recommended. For high erosion zones of 10 to 20 $t \cdot ha^{-1} \cdot yr^{-1}$, which constitute the majority of the district, reduced tillage, diversified crop rotations, vegetative buffer strips, and managed rotational grazing are recommended. With appropriate conservation investments implemented systematically across the landscape according to the erosion risk zones identified in this study, it is possible to reduce erosion rates to sustainable levels, preserve soil resources for future generations, and maintain the agricultural productivity upon which the region's rural communities depend.

Declarations

Acknowledgments

The authors thanks local farmers in the Parkent district for providing field access and assistance during soil sampling campaigns. The authors acknowledges the United States Geological Survey (USGS) for providing Landsat 8 OLI/TIRS data; the National Aeronautics and Space Administration (NASA) for providing SRTM DEM data; the Climate Hazards Group for providing CHIRPS precipitation data; and the GlobeLand30 team for providing land use/land cover data. The authors also thanks the National Committee on Ecology and Climate Change of the Republic of Uzbekistan for organizing training seminars that informed the methodological approach of this study. This work was supported by the Tashkent State Agrarian University, Uzbekistan and Youth Breeding Experts Program of the Yangling Agricultural High-tech Industries Demonstration Zone, China.

Author contributions

Conceptualization, G.S. and T.M.; methodology, G.S., Z.W. and A.R.T.; software, G.S. and L.A.; validation, G.S., Z.W., T.S. and T.M.; formal analysis, G.S. and T.M.; investigation, G.S., L.A., and A.R.T.; resources,

Z.W., T.S. and T.M.; data curation, G.S. and L.A.; writing – original draft preparation, G.S. and Z.W.; writing – review and editing, A.R.T., L.A.and T.M.; visualization, Z.W. and G.S.; supervision, L.A.; project administration, T.M.. All authors have read and agreed to the published version of the manuscript.

Funding

This research was funded by Youth Breeding Experts Program of the Yangling Agricultural High-tech Industries Demonstration Zone, China.

Data availability

The datasets used and analyzed in the present work are available from the corresponding author on request.

Ethical approval

Not applicable.

Competing interests

The authors declare no competing interests.

References

1. Abdul Rahaman S, Aruchamy S, Jegankumar R, Abdul Ajeez S (2015) Estimation of annual average soil loss, based on RUSLE model in Kallar watershed, Bhavani basin, Tamil Nadu, India. ISPRS Ann Photogramm Remote Sens Spatial Inf Sci II-. 2/W2:207–214. https://doi.org/10.5194/isprsannals-II-2-W2-207-2015
2. Barboza E, Raymundo AA, Valqui L, Salas Lopez R (2024) Estimation of soil loss using RUSLE model and GIS tools: Case study in the Jucusbamba Micro-Watershed (Amazonas, NW Peru). Ambiente Água 19:e2998. https://doi.org/10.4136/ambi-agua.2998
3. Bazzoffi P (2007) Erosione del suolo e qualità ambientale. Il Sole 24 Ore Edagricole, Bologna
4. Benavidez R, Jackson B, Maxwell D, Norton K (2018) A review of the (Revised) Universal Soil Loss Equation (RUSLE): With a view to increasing its global applicability and improving soil loss estimates. Hydrol Earth Syst Sci 22(11):6059–6086. https://doi.org/10.5194/hess-22-6059-2018
5. Bobomurodov S, Abdurakmonov N, Niyazmetov U, Baxodirov Z, Normatov Y, Abdurakhmonov Z (2023) Mapping of soil properties using geographic information systems (on the example of Tashkent region). E3S Web Conf 386:04006. https://doi.org/10.1051/e3sconf/202338604006
6. Byizigiro RV, Rwanyiziri G, Mugabowindekwe M, Kagoyire C, Biryabarema M (2020) Estimation of soil erosion using RUSLE model and GIS: The case of Satinskyi Catchment, Western Rwanda. Rwanda J Eng Sci Technol Environ 3. https://doi.org/10.4314/rjeste.v3i1.2S. (Special Issue)

7. Chen J, Chen J, Liao A, Cao X, Chen L, Chen X et al (2015) Global land cover mapping at 30 m resolution: A POK-based operational approach. ISPRS J Photogramm Remote Sens 103:7–27. https://doi.org/10.1016/j.isprsjprs.2014.09.002
8. Food and Agriculture Organization (2012) Soil degradation and restoration in Afghanistan. Food and Agriculture Organization of the United Nations, Rome. https://doi.org/10.4324/9780203847459
9. Funk C, Peterson P, Landsfeld M, Pedreros D, Verdin J, Shukla S et al (2015) The climate hazards infrared precipitation with stations—a new environmental record for monitoring extremes. Sci Data 2:150066. https://doi.org/10.1038/sdata.2015.66
10. Ganasri BP, Ramesh H (2015) Assessment of soil erosion by RUSLE model using remote sensing and GIS: A case study of Nethravathi Basin. Geosci Front 7(6):953–961. https://doi.org/10.1016/j.gsf.2015.10.007
11. Girmay G, Moges A, Muluneh A (2020) Estimation of soil loss rate using the USLE model for Agewmariayam Watershed, northern Ethiopia. Agric Food Secur 9:9. https://doi.org/10.1186/s40066-020-00262-w
12. Hengl T, Mendes de Jesus J, Heuvelink GBM, Ruiperez Gonzalez M, Kilibarda M, Blagotić A et al (2017) SoilGrids250m: Global gridded soil information based on machine learning. PLoS ONE 12(2):e0169748. https://doi.org/10.1371/journal.pone.0169748
13. Juliev M, Kholmurodova M, Gafurova L, Khoshjanova K, Khomidov A, Agzamova I et al (2024) Modeling of soil erosion based on geospatial techniques and the RUSLE model for the Ugam Chatkal National Park, Uzbekistan. J Geol Geogr Geoecol 33(3):485–494. https://doi.org/10.15421/112445
14. Juliev M, Matyakubov B, Khakberdiev O, Abdurasulov X, Gafurova L, Ergasheva O et al (2022) Influence of erosion on the mechanical composition and physical properties of serozems on rainfed soils, Tashkent province, Uzbekistan. IOP Conf Ser Earth Environ Sci 1068(1):012005. https://doi.org/10.1088/1755-1315/1068/1/012005
15. Khan A, Rahman A, Mahmood S (2023) Spatial estimation of soil erosion risk using RUSLE model in District Swat, Eastern Hindu Kush, Pakistan. J Water Clim Change 14(6):1881–1898. https://doi.org/10.2166/wcc.2023.495
16. Kholmurodova M, Gafurova L, Ergasheva O, Saidova M, Juliev M (2025) Zig-zag furrow irrigation for cotton production on erosion-prone meadow-sierozem soils in Uzbekistan. Front Agron 6:1778881. https://doi.org/10.3389/fagro.2025.1778881
17. Moore ID, Burch GJ (1986) Physical basis of the length-slope factor in the Universal Soil Loss Equation. Soil Sci Soc Am J 50(5):1294–1298. https://doi.org/10.2136/sssaj1986.03615995005000050042x
18. National Committee on Ecology and Climate Change of the Republic of Uzbekistan (2025) Attention given to effective soil erosion assessment. Government Portal of the Republic of Uzbekistan. Available at: https://gov.uz/en/eco/news/view/99923. Accessed 6 Nov 2025

19. Okenmuo FC, Ewemoje TA (2023) Estimation of soil water erosion using RUSLE, GIS, and remote sensing in Obibia River watershed, Anambra, Nigeria. DYSONA Appl Sci 4:6–14. https://doi.org/10.30493/DAS.2022.349144
20. Pimentel D, Harvey C, Resosudarmo P, Sinclair K, Kurz D, McNair M et al (1995) Environmental and economic costs of soil erosion and conservation benefits. Science 267(5201):1117–1123. https://doi.org/10.1126/science.267.5201.1117
21. Poesen J, Nachtergaele J, Verstraeten G, Valentin C (2003) Gully erosion and environmental change: Importance and research needs. CATENA 50(2–4):91–133. https://doi.org/10.1016/S0341-8162(02)00143-1
22. Renard KG, Freimund JR (1994) Using monthly precipitation data to estimate the R-factor in the revised USLE. J Hydrol 157(1–4):287–306. https://doi.org/10.1016/0022-1694(94)90110-4
23. Renard KG, Foster GR, Weesies GA, McCool DK, Yoder DC (1997) Predicting soil erosion by water: A guide to conservation planning with the Revised Universal Soil Loss Equation (RUSLE). USDA Agriculture Handbook No. 703. United States Department of Agriculture, Washington, DC
24. Sodikova GS, Shadieva NI, Saidova ME (2025) Assessment of mountain soil erosion threat levels based on digital analysis. BIO Web Conf 173:03003. https://doi.org/10.1051/bioconf/202517303003
25. Sodikova GS, Shamsiddinov TSh, Shadieva NI, Usmonova MI (2025) GIS assessment of soil erosion in Uzbekistan's rainfed mountain lands. BIO Web Conf 173:03024. https://doi.org/10.1051/bioconf/202517303024
26. Shin GJ (1999) The analysis of soil erosion in a watershed using GIS. Dissertation, Kangwon National University
27. Stanchi S, Freppaz M, Godone D, Zanini E (2014) Soil erosion assessment, snow cover dynamics and avalanche release sites: A case study from the Aosta Valley (NW Italy). Nat Hazards Earth Syst Sci 14:1761–1771. https://doi.org/10.5194/nhess-14-1761-2014
28. Terefe B, Melese T, Tsegaye A, Afework A, Yibeltal T, Anagaw A et al (2025) Soil loss estimation in Ethiopia: A comprehensive review of the RUSLE model integrated with geospatial technologies. Discover Sustain 6:334. https://doi.org/10.1007/s43621-025-01037-8
29. Van der Knijff JM, Jones RJA, Montanarella L (2000) Soil erosion risk assessment in Europe. European Soil Bureau, European Commission, Luxembourg
30. Wischmeier WH, Smith DD (1978) Predicting rainfall erosion losses: A guide to conservation planning. USDA Agriculture Handbook No. 537. United States Department of Agriculture, Washington, DC
31. Yesuph AY, Dagnew AB (2019) Soil erosion mapping and severity analysis based on RUSLE model and local perception in the Beshillo Catchment of the Blue Nile Basin, Ethiopia. Environ Syst Res 8:17. https://doi.org/10.1186/s40068-019-0145-1
32. Zhang B, Fang H, Wu S, Li C, Wang Y, Siddique KHM (2024) Soil erosion prediction and spatiotemporal heterogeneity in driving effects of precipitation and vegetation on the northern slope

Figures




Figure 1

Location of the study area in the Tashkent region, Uzbekistan, showing the Parkent district boundaries, elevation gradient (500-1,200 m above sea level), and major geographical features. The map includes a scale bar, north arrow, and coordinate grid (UTM zone 42N, WGS84).




Figure 2

Spatial distribution of rainfall erosivity (R-factor) in the Zarkent massif Parkent district, averaged over the period 2015-2024. Higher values (darker shades) in the northeastern portion of the study area reflect greater orographic precipitation on windward slopes.




Figure 3

K-factor soil erosion map of the Zarkent massif in the Parkent district.




Figure 4

LS-factor map of the Zarkent massif Parkent district, showing the spatial distribution of slope length and steepness effects on erosion potential. Darker shades indicate higher LS values and greater topographic control on erosion. The concentration of high LS values (red tones) in the northeastern mountainous areas is evident.




Figure 5

Mean cover management (C-factor) map for the Zarkent massif Parkent district, 2015–2024, derived from Landsat 8 NDVI time series. Darker shades indicate higher C-values and lower vegetation protection. The highest erosion risk from cover management occurs in cultivated areas (yellow to red tones) in the central and western portions of the district.




Figure 6

Support practice (P-factor) map of the Zarkent massif Parkent district, derived from GlobeLand30 LULC data and field verification. Areas with terracing (P=0.25, dark green) are concentrated in small patches on moderate slopes; the majority of the district has P=0.90-1.00 (yellow to red), indicating minimal conservation interventions.




Figure 7

Erosion risk map of the Zarkent massif Parkent district Uzbekistan, derived from RUSLE analysis over the period 2015-2024. The map illustrates erosion risk zones classified into six degrees: very low (<1 $t \cdot ha^{-1} \cdot yr^{-1}$), low (1–5 $t \cdot ha^{-1} \cdot yr^{-1}$), medium (5-10 $t \cdot ha^{-1} \cdot yr^{-1}$), high (10-20 $t \cdot ha^{-1} \cdot yr^{-1}$), very high (20-40 $t \cdot ha^{-1} \cdot yr^{-1}$), and extremely high (>40 $t \cdot ha^{-1} \cdot yr^{-1}$). Highly eroded soils exceeding 40 $t \cdot ha^{-1} \cdot yr^{-1}$ cover 68.84 hectares as detailed in Table 3. The map is based on Landsat 8 OLI/TIRS remote sensing data and GIS processing (USGS, 2024).




Figure 8

Average amount of eroded soil from 2014-2024