

Maternal hypertension and cardiovascular medications dysregulate placental arterial tone

Teresa Tropea

`teresa.tropea@manchester.ac.uk`

Maternal & Fetal Health Research Centre, Division of Developmental Biology & Medicine, School of Medical Sciences, Faculty of Biology, Medicine and Health, University of Manchester

Elizabeth C. Cottrell

Maternal & Fetal Health Research Centre, Division of Developmental Biology & Medicine, School of Medical Sciences, Faculty of Biology, Medicine and Health, University of Manchester

Raianne Wallworth

Maternal & Fetal Health Research Centre, Division of Developmental Biology & Medicine, School of Medical Sciences, Faculty of Biology, Medicine and Health, University of Manchester

Nardeen Khalil

Maternal & Fetal Health Research Centre, Division of Developmental Biology & Medicine, School of Medical Sciences, Faculty of Biology, Medicine and Health, University of Manchester

Edward D. Johnstone

Maternal & Fetal Health Research Centre, Division of Developmental Biology & Medicine, School of Medical Sciences, Faculty of Biology, Medicine and Health, University of Manchester

Jenny E. Myers

Maternal & Fetal Health Research Centre, Division of Developmental Biology & Medicine, School of Medical Sciences, Faculty of Biology, Medicine and Health, University of Manchester

Paul Brownbill

Maternal & Fetal Health Research Centre, Division of Developmental Biology & Medicine, School of Medical Sciences, Faculty of Biology, Medicine and Health, University of Manchester

Article

Keywords:

Posted Date: June 29th, 2026

DOI: <https://doi.org/10.21203/rs.3.rs-10061694/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Additional Declarations: No competing interests reported.

Version of Record: A version of this preprint was published at Scientific Reports on September 11th, 2026. See the published version at <https://doi.org/10.1038/s41598-026-69028-x>.

Abstract

Cardiovascular medications prescribed to pregnant women, including Ca^{2+} channel blockers (CCBs), α - and β -adrenergic receptor antagonists, and α -adrenergic receptor agonists enter the fetal circulation, with unknown effects on the fetoplacental vasculature. We investigated whether cardiovascular medications alter human fetoplacental vascular tone and placental perfusion.

Chorionic plate arteries (CPAs) were obtained from placentas of normotensive women, women with unmedicated hypertension, and those chronically medicated with amlodipine, nifedipine, labetalol or bisoprolol, or CCB–labetalol combinations. Using wire myography, acute effects of amlodipine, nifedipine, labetalol, methyldopa, doxazosin, bisoprolol and nebivolol were tested in pre-constricted CPAs from normotensive women. Vascular reactivity following chronic hypertension and/or medication exposure was assessed by vasoconstriction to high potassium physiological solution (KPSS) and U46619, and by relaxation to sodium nitroprusside (SNP).

Acute exposure to amlodipine, nifedipine, doxazosin and nebivolol promoted vasorelaxation ($P < 0.05$), while labetalol, methyldopa ($P < 0.05$) and bisoprolol ($P < 0.001$) increased vasoconstriction relative to controls. Chronic hypertension and labetalol-treatment reduced vasoconstriction to KPSS ($P < 0.05$). Chronic hypertension and bisoprolol-treatment attenuated responses to U46619 ($P < 0.01$) and SNP ($P < 0.01$).

Maternal hypertension impairs placental vascular function, and cardiovascular medications may further dysregulate fetoplacental vascular reactivity, with potential implications for maternal and fetal outcomes. Mechanistic studies alongside clinical correlations are essential to guide evidence-based prescribing.

Introduction

Cardiovascular disease is common in pregnant women and is associated with maternal mortality. The spectrum includes cardiac disease and hypertension, both contributing to poor maternal cardiovascular adaptation to pregnancy. Amongst pregnant women who died between 2020–2022, 13% of them were diagnosed with cardiac problems (1).

Pregnancy requires major maternal hemodynamic adaptations to meet fetal metabolic demands, which may challenge the maternal cardiovascular system and contribute to hypertensive disorders (2). Maternal hypertension affects 8–10% of pregnancies worldwide and is diagnosed when blood pressure is greater than 140/90 mmHg (3, 4). Chronic hypertension is defined as occurring before pregnancy or in the first 20 weeks of gestation.

A limited range of antihypertensives, considered safe for the fetus, are prescribed to pregnant women with cardiovascular disease. As the CHIPS and CHAP trials data demonstrated, targeting maternal blood pressure protects the maternal cardiovascular system and prevents the occurrence of severe

hypertension (5, 6). Common antihypertensive medications used in pregnancy include labetalol, methyldopa, and calcium channel blockers (CCBs) such as amlodipine and nifedipine (7). Bisoprolol is one of the most common β_1 -adrenergic receptor antagonists prescribed to pregnant women with cardiac disease, including congenital heart disease, cardiomyopathy and arrhythmia (8, 9). These medications cross the placental barrier (10) and by entering the fetoplacental circulation, may have detrimental effects on fetal growth and placental function (11).

L-type voltage-gated Ca^{2+} channels regulate vascular tone in the smooth muscle layer. Previous studies showed that L-type VGCCs are functionally expressed in the fetoplacental vasculature and inhibition by CCBs, such as nifedipine, promotes vasorelaxation in chorionic plate arteries (CPAs) (12). Medications targeting the adrenergic system are also recommended to treat maternal hypertension and to minimise cardiac workload in women with cardiac disease.

α - and β -adrenergic receptors mediate the effects of the sympathetic nervous system in response to the catecholamines, noradrenaline and adrenaline, and contribute to the regulation of the cardiovascular system (13). In the systemic circulation, α -adrenergic receptors located on the endothelium, as well as β -adrenergic receptors on both endothelial and smooth muscle cells, have been reported to promote vasorelaxation. In contrast, α -adrenergic receptors expressed on smooth muscle cells mediate vasoconstriction (14). Although the placenta lacks sympathetic innervation, evidence suggests that adrenergic receptors are expressed in the fetoplacental circulation, and that their stimulation and differences in receptor abundance between cell types may evoke different vasoactive effects compared to the systemic circulation (14).

High blood flow and low resistance within the placental vasculature enable adequate delivery of oxygen and nutrients to the growing fetus. Antenatal exposure to mainly β -adrenergic receptor antagonists (15–19) and α_2 -adrenergic receptor agonists (20), has been associated with higher risks of fetal growth restriction (FGR), increased rates of small for gestational age (SGA) newborns and preterm birth. It is unclear whether the increased risks of adverse perinatal outcomes observed in women with cardiovascular disease are attributable to dysregulated placental perfusion associated with the underlying maternal disease or to the medications used to manage these conditions.

Currently, clinicians have minimal guidance to determine which antihypertensive treatment most effectively supports healthy outcomes for mothers and their babies (21). In this study, we tested the hypothesis that administration of antihypertensive and cardiac medications to pregnant women would alter placental vascular function. Using the wire myography system, we tested the acute effects of amlodipine and nifedipine (CCBs); labetalol (α/β - adrenergic receptor competitive antagonist) methyldopa (α_2 -adrenergic receptor agonist), doxazosin (α_1 -adrenergic receptor antagonist), bisoprolol and nebivolol (β_1 -adrenergic receptor antagonists), on CPAs isolated from the placenta of healthy women. In addition, we assessed the vascular function of CPAs isolated from women with cardiovascular conditions who were either untreated or chronically exposed *in vivo* to either amlodipine,

or nifedipine, or labetalol therapies, or labetalol in combination with CCBs, or β 1-adrenergic receptor antagonist (bisoprolol) therapy.

Methods

Ethics Statement

The present study was carried out in accordance with institutional guidelines and conformed to the standards set by the Declaration of Helsinki.

All placental tissues were obtained from women who delivered at St. Mary's Hospital, Manchester, United Kingdom, in accordance with approved guidelines at the hospital. Ethical approval was granted by North West - Greater Manchester West Research Ethics Committee (15/NW/0829).

Informed written consent was obtained from study participants prior to delivery.

Data Availability

The data supporting the findings of this study are available from the corresponding author upon reasonable request. The detailed method, including criteria for the study, can be found in the Supplemental Material.

A total of N = 107 study participants was recruited, and the placentas were collected following either elective Caesarean section or normal vaginal delivery.

Isolated CPAs were mounted on the wire myography system (22). The study had two distinct objectives. In the first one, arteries obtained from the placenta of normotensive women were pre-constricted with the thromboxane A₂ mimetic, U46619 (80% maximal effective concentration) and *ex vivo* effects of amlodipine, nifedipine, labetalol, methyldopa, doxazosin, bisoprolol and nebivolol were tested in a concentration-dependent manner (10^{-11} – 10^{-5} M range), based on known (23–29) or estimated (30, 31) fetal venous plasma concentrations (Table S1). In the second objective, vasoconstriction to U46619 (10^{-10} – 2×10^{-6} M), and relaxation (in pre-constricted vessels) to the nitric oxide (NO) donor, sodium nitroprusside (SNP; 10^{-10} – 10^{-5} M), was tested on CPAs isolated from placentas of women with hypertension who either remained untreated throughout their pregnancies or were treated with cardiovascular medications to assess the effect of chronic exposure to hypertension and/or cardiovascular medications on fetoplacental vascular reactivity.

Statistical analysis

Concentration-responses to SNP and medications are expressed as percentage level of vasoconstriction remaining from the pre-constriction achieved with the U46619 EC₈₀ dose. All data are expressed as

mean $\pm$ SEM and differences between groups analysed by 1- or 2-way ANOVA or as otherwise stated. Statistical significance was defined as $P < 0.05$ (for details see Supplemental Material).

Results

Table 1 summarises the clinical characteristics of pregnant women and their pregnancy outcomes (for demographics of pregnant women see Tables S2 Supplemental Material), grouped by maternal disease and/or medication exposure

NP, normal pregnancy. Systolic and diastolic blood pressure (SBP, DBP; mmHg) in the first pregnancy and at the last pre-delivery (SBP_{PD}, DBP_{PD}) measurement, gestational age at last pre-delivery measurement (GA_{PD} >245 days or 35 weeks), pre-delivery umbilical artery pulsatility index (Umb PI_{PD}), gestational age-adjusted centile for umbilical artery Doppler (Umb PI_{PD} centile), gestational age-adjusted centile for estimated fetal weight (EFW) according to the Hadlock formula, soluble fms-like tyrosine kinase-1 to placental growth factor ratio (sFlt-1/PlGF_{PD}), gestational age (days) at delivery (GA_D), birth weight (BW; grams) and sex-adjusted BW centile are presented as median and (interquartile range). Frequency of maternal cardiac disease, systolic function categories, diabetes, gestational diabetes mellitus (GDM; diagnosed before 24 weeks and between 24–30 weeks of gestation), chronic hypertension (HTN, diagnosed pre-pregnancy or before 20 weeks of gestation), pregnancy complications (pre-eclampsia), fetal growth restriction (FGR), small for gestational age (SGA) or other growth-risk, sex of the baby and preterm births (< 259 days) are shown as number n, and percentage (%).

In the total cohort, first pregnancy and pre-delivery blood pressures were higher in the unmedicated and in the medicated hypertensive groups compared to those with no hypertension. 10% of women had cardiac disease, with the highest rate in the bisoprolol group (69%); in the same group, systolic function was moderately and severely impaired in a total of 30% of women. 3% of women had a diagnosis of type 2 and gestational diabetes mellitus (GDM) before 24 weeks, 6% were diagnosed with GDM between 24–30 weeks of pregnancy, and 2% had type 2 diabetes pre-pregnancy. 29% of women had chronic hypertension, with the highest rate in the amlodipine- (100%) and in the labetalol + CCBs-medicated (67%) groups. Pregnancy complications affected 19% of pregnancies in the form of pre-eclampsia (PE), mainly in the nifedipine- (50%) and in the labetalol-medicated (44%) groups.

Umbilical Doppler assessments showed a trend toward higher PI, with median gestation-adjusted umbilical artery PI centiles of 78 (46–90) and 70 (46–91) in the labetalol and bisoprolol groups, respectively. These groups also had the highest rates of SGA (61% and 38%, respectively).

The bisoprolol cohort had the longest exposure time during pregnancy amongst all the medicated groups (Table S3).

Labetalol, whether administered alone or in combination with CCBs, was associated with higher number of preterm births (39% and 33%, respectively).

Acute ex vivo exposure to antihypertensives differentially alters fetoplacental vascular tone

When tested *ex vivo*, the CCBs amlodipine (Fig. 1A) and nifedipine (Figure S1), the α_1 -adrenergic receptor antagonist doxazosin (Fig. 1D), and the third generation β_1 -adrenergic receptor antagonist nebivolol (Fig. 1E), promoted significant vasorelaxation in submaximally pre-constricted CPAs isolated from the placenta of normotensive pregnant women. Compared to controls, amlodipine caused $31.9 \pm 4.3\%$ maximum vasodilatory effect ($E_{\max}$; $P < 0.0001$ overall effect vs control), followed by nebivolol ($P = 0.036$ overall effect vs control) $17.9 \pm 5.6\%$ $E_{\max}$, and doxazosin ($P = 0.0183$ overall effect vs control) $14.8 \pm 4.6\%$ $E_{\max}$.

The α/β -adrenergic receptor competitive antagonist labetalol (Fig. 1B), the α_2 -adrenergic receptor agonist methyldopa (Fig. 1C) and the second generation β_1 -adrenergic receptor antagonist bisoprolol (Fig. 1F) exhibited increased vasoconstriction compared to their relative controls, with labetalol ($P < 0.0469$ overall effect vs control) reaching $15.5 \pm 4.0\%$ $E_{\max}$, and methyldopa ($P = 0.0004$ overall effect vs control) causing $13.0 \pm 3.0\%$ $E_{\max}$. Bisoprolol maintained higher level of pre-constriction over time compared to its relative control group ($94.0 \pm 4.6\%$ vs $82.9 \pm 4.8\%$ at the maximum concentration tested; $P = 0.0005$ overall effect vs control).

Maternal hypertension, but not Ca^{2+} channel blocker therapy, reduces constriction and relaxation responses in ex vivo chorionic plate arteries

Compared to the normotensive group (NP), CPAs in the untreated hypertensive group (HTN) had a significantly reduced response to 120 mmol/L K^+ solution (KPSS vasoconstriction in kPa: NP, 9.2 ± 0.6 ; HTN 6.8 ± 0.6 ; $P < 0.05$ Fig. 2A&D). In comparison with the NP group, CPAs from women treated with CCBs did not have significantly altered KPSS responses (amlodipine, 7.5 ± 0.7 ; nifedipine, 8.0 ± 1.1). Cumulative concentration-dependent vasoconstriction to U46619 was significantly reduced in the HTN and in the CCB groups (U46619 $E_{\max}$ in kPa: NP, 14.8 ± 1 ; HTN, 10.0 ± 0.8 ; amlodipine, 10.9 ± 1.1 ; nifedipine, 11.5 ± 1.4 . $P < 0.0001$ Fig. 2B and 2E). Vasorelaxation was slightly reduced at SNP concentrations of 10^{-7} and 10^{-6} M in the CCB groups, compared to the NP group, but was not different to the underlying HTN effect (Fig. 2C and 2F). Overall, HTN reduced responses to SNP compared with NP (SNP $E_{\max}$ in %: NP, 73.1 ± 4.6 ; HTN, 62.8 ± 4.3 ; amlodipine, 68.5 ± 5.69 ; nifedipine, 62.3 ± 5.7 . $P = 0.005$ Fig. 2C and $P = 0.0006$ 2F). There was no difference between HTN and CCB groups.

Labetalol treatment does not worsen hypertension-associated vascular dysfunction in ex vivo fetoplacental arteries

In women with HTN exposed to labetalol during pregnancy, vasoconstriction of CPAs in response to 120 mmol/L K^+ solution and to U46619 was overall significantly lower compared to the NP group (KPSS vasoconstriction in kPa: 5.9 ± 0.5 $P = 0.0002$ Fig. 3A; U46619 $E_{\max}$ in kPa: 11.1 ± 1.0 . $P < 0.0001$ Fig. 3B). Compared to the NP group, vasorelaxation to SNP was significantly reduced in the labetalol group (SNP

Emax in %: labetalol 68.1 ± 3.4 , $P = 0.0042$ Fig. 3C). However, there was no significant difference between HNT and labetalol groups.

In vivo Ca^{2+} channel blockers combined with labetalol does not further dysregulate the underlying hypertension effect in ex vivo fetoplacental arteries

In CPAs isolated from hypertensive women taking a combination of labetalol and CCBs during pregnancy, vasoconstriction in response to 120 mmol/L K^+ solution was reduced compared to the NP group (KPSS vasoconstriction in kPa: labetalol + CCBs, 8.1 ± 0.8 . $P = 0.0336$ Fig. 4A). However, there was no difference between the HTN and labetalol + CCBs-treated group. A significant attenuation of vasoconstriction in response to U46619 (U46619 Emax in kPa: 11.2 ± 1.0 . $P < 0.0001$ Fig. 4B), and of vasorelaxation to SNP (SNP Emax in %: 68.1 ± 3.3 . $P = 0.0011$ Fig. 4C) was shown in the labetalol + CCBs group compared to NP. At a concentration of SNP 10^{-7}M , comparison between groups showed significant effect of reduced vasorelaxation in both the HTN and the labetalol + CCBs-treated group (Fig. 4C).

In vivo bisoprolol exposure diminishes vasoconstriction and vasorelaxation responses in ex vivo fetoplacental arteries

Women with cardiac dysfunction included in this cohort were not hypertensive, therefore, this group was compared with CPAs from women with a normotensive pregnancy. In women with cardiac dysfunction, treatment with bisoprolol during pregnancy did not affect vasoconstriction of CPAs in response to 120 mmol/L K^+ solution (KPSS vasoconstriction in kPa: bisoprolol, 7.7 ± 1.0 . $P = 0.1847$ Fig. 5A); whereas CPAs from this group exhibited significantly reduced vasoconstriction to U46619 (U46619 Emax in kPa: 12.1 ± 1.3 . $P = 0.0087$ Fig. 5B) compared to NP. Comparison between groups also showed that chronic bisoprolol exposure significantly reduced vasorelaxation to SNP (SNP Emax in %: 54.3 ± 5.1 . $P = 0.0002$ Fig. 5C).

There was insufficient data to compare the effects of *in vivo* nebivolol exposure on fetoplacental vascular responses. Exclusion of CPA samples obtained from PE participants did not affect data analysis (Figure S2-S4).

Discussion

To our knowledge, this is the first study to compare the effects of maternal hypertension, and exposure to antihypertensive and cardioprotective medications commonly prescribed in pregnancy, on human fetoplacental vascular function.

Our *ex vivo* data revealed that acute exposure to CCBs promoted vasorelaxation, whereas medications modulating the adrenergic system elicited contrasting effects on CPAs from normotensive healthy pregnancies. Acute labetalol, methyldopa and bisoprolol exposure induced increased vasoconstriction compared to respective diluent controls. In contrast, doxazosin and nebivolol induced significant

vasorelaxation. Using CPAs from pregnancies complicated by hypertension and cardiac disease, we found that maternal hypertension *per se* reduced CPA reactivity in response to both high potassium solution and to a thromboxane-mimetic vasoconstrictor. Vasorelaxation to exogenous NO was also attenuated in CPAs from women with hypertensive pregnancy. We observed no major differences between CPAs from the untreated hypertensive group and those treated with different medication classes during pregnancy.

In CPAs from normal pregnancies, L-type CCBs amlodipine and nifedipine induced marked vasorelaxation. Our findings are consistent with previous studies on fetoplacental arteries in our group (12), and in those of others conducted in human internal mammary arteries at a comparable concentration-range (32). In contrast to CCBs, medications targeting the adrenergic system, produced more disparate effects. Compelling evidence showed that the human fetoplacental vasculature expresses adrenergic receptors, and their vascular effects vary depending on the activation/deactivation of specific receptor subtype(s) (14). Exposure to β -adrenergic receptor antagonists, including labetalol, reduces fetoplacental blood flow and increases placental vascular resistance in hypertensive pregnant ewes (33). Maternal labetalol administration increases umbilical artery pulsatility index (PI) in human pregnancies, an effect attributed to increased placental vasoconstriction (34), and supported by evidence from *ex vivo* studies using human placentas perfused with labetalol (35). Whilst nifedipine increases placental perfusion *ex vivo* (36), in humans (37) and in animal models (38), in comparison with the oral treatment with labetalol, there was no significant difference in maternal and neonatal outcomes (39). In contrast to our findings of labetalol and methyldopa, which further increased vasoconstriction in arteries pre-constricted with U46619, previous work has shown vasorelaxing properties of labetalol in human forearm radial arteries rings, partly mediated via activation of calcium-activated potassium channels when tested acutely *ex vivo* (10^{-8} – 10^{-4} M range)(40). Labetalol is a non-selective and competitive β -blocker, with an approximate β : α -blockade ratio of 7:1 (41). Given the competitive antagonism of labetalol favouring β -blockade, it may produce a stronger inhibitory effect on the endothelial-dependent vasodilatory actions mediated by β -adrenergic receptor activation. The resulting imbalance in the blockade effect, and the higher expression of α -adrenergic receptors in human term fetoplacental arteries (42), may promote α -mediated vasoconstriction, thereby potentiating the initial level of pre-constriction in response to the thromboxane mimetic in the smooth muscle cells. This observation is consistent with previous findings from *ex vivo* dual perfusion studies of the human placenta, where labetalol administration increased the perfusion pressure induced by endothelin-1 and U46619 (35). The α -adrenergic agonism of methyldopa might have contributed to the observed comparable response in the fetoplacental vasculature.

Bisoprolol did not evoke overt vasoconstriction *ex vivo*, but vessels exhibited a sustained level of tension over time relative to the diluent control. Compared to labetalol, bisoprolol is a cardioselective β -blocker with ~ 100 times higher affinity for β_1 - over β_2 -adrenergic receptors, and with no affinity for α -adrenergic receptors (43). The absence of α -adrenergic agonism might have slightly attenuated the initial pre-constriction, while the β -blocking properties likely prevented vasorelaxation.

In contrast, the α -adrenergic receptor antagonism of doxazosin (44), and the vasodilatory properties of nebivolol (45, 46) promoted relaxation of the fetoplacental vasculature *ex vivo*. Studies have demonstrated the vasodilatory effect of nebivolol in human penile arteries via NO-dependent mechanisms (47) and partly mediated through calcium activated potassium channels in human radial arteries (40). In coronary resistance microarteries of humans and rodents, nebivolol has been shown to target the endothelial β_3 -adrenergic receptors (46), and to release NO via β_2 -adrenergic receptors in mouse thoracic aorta (48). Our study was underpowered to investigate the effect of *in vivo* exposure to nebivolol. However, consistent with our findings of acute exposure causing vasorelaxation, human umbilical vein endothelial cells (HUVECs) exposed to plasma from preeclamptic women showed increased NO production in the presence of nebivolol, an effect that was abolished by NO-synthase and β_3 -adrenergic receptor inhibitors (49).

While endothelial dysfunction and increased production of vasoconstrictors are key factors in the pathophysiology of hypertension (50), vascular smooth muscle cells are the functional effectors of vascular tone modulation (51). In our study, we challenged mechanisms of vasoconstriction and relaxation both converging on vascular smooth muscle cell function. We showed that maternal hypertension was associated with impaired vasoconstriction and reduced endothelium-independent vasorelaxation in CPAs, independent of medication exposure during pregnancy. This suggests that the fetoplacental circulation is altered in pregnancies complicated by maternal hypertension, and that vascular responsiveness to vasoactive agents is different to the systemic circulation potentially due to lower level of sympathetic tone regulation and imbalance between α - and β -adrenergic receptor mediated effects in the fetoplacental vascular bed. Pre-eclampsia developed in 19% of the participants; however, no cases occurred in either the normal or the bisoprolol group. Exclusion of pre-eclampsia cases from our groups did not affect data analysis.

Although current evidence indicates an increased risk of infants being born small for gestational age (SGA) in women with hypertensive disorders of pregnancy, it is uncertain whether this association is exacerbated by antihypertensive medication (52). The Giant PANDA trial, a pragmatic, multicenter, randomized controlled study involving 2,200 women with pregnancy hypertension, aims to generate high-quality evidence to guide the selection of first-line antihypertensive therapy by balancing effective maternal blood pressure management with optimal fetal and neonatal outcomes (21). In our cohorts, we observed no major differences in *ex vivo* vascular reactivity of placental vessels between the unmedicated hypertensive group and those treated with medications during pregnancy, suggesting that *in vivo* changes in placental vascular function had a more significant impact on *ex vivo* vascular reactivity than exposure to maternal antihypertensive agents. These findings were unchanged following removal of pregnancies complicated by pre-eclampsia from the analysis.

In contrast, *in vivo* bisoprolol exposure was associated with a relative reduction of maximal CPA vasorelaxation to SNP of $\sim 26\%$ in comparison to vessels from uncomplicated pregnancies with no medication exposure. The more pronounced impairment of vasorelaxation observed in the bisoprolol cohort may suggest that bisoprolol as a cardioselective β_1 -blocker alters the cell signalling pathway

phenotype leading to altered eNOS expression and/or activation. It is also possible that there are chronic changes in signalling in the smooth muscle cells (e.g. impairments in NO-mediated signalling pathways) induced by *in vivo* bisoprolol exposure. Previous cohort studies support an association between *in vivo* bisoprolol exposure and changes in fetal growth (9) (53) (54) and fetal Dopplers (unpublished data from St Mary's Hospital). However, it is very difficult to demonstrate a casual effect of bisoprolol exposure on placental vascular function as, in the same way that maternal hypertensive disease appears to influence *ex vivo* CPA vasoreactivity, there may be an underlying impact of maternal cardiac disease even though maternal blood pressure has remained normal. The maternal indications for bisoprolol use are also varied, ranging from management of maternal palpitations to significant cardiac impairment. In this small cohort, it has not been possible to disentangle the relative impact of underlying disease severity, bisoprolol duration or dose on *ex vivo* CPA function.

The regulation of the fetoplacental circulation is critical for normal fetal growth and development, as it governs oxygen, nutrient, and waste exchange between the feto-maternal compartments. Within the chorionic plate, CPAs of resistance-artery size are key determinants of fetoplacental vascular resistance and blood flow distribution, thereby influencing downstream perfusion of the villous tree and, consequently, the efficiency of oxygen and nutrient delivery to the fetus (55). Dysregulation of the fetoplacental vascular tone and/or impaired responsiveness to vasoactive stimuli can compromise fetoplacental perfusion and disrupt the exchange process within the terminal villi. This may contribute to complications such as FGR and pre-eclampsia, where placental dysfunction is central to the pathophysiology of the disease (56, 57). A multicentre retrospective cohort study showed that, compared with women not taking β -blockers, those exposed had ~ a 1.4-fold increase in SGA and a 2.3-fold increase in FGR prevalence (9).

One limitation of this study is the concentration ranges chosen for the acute testing, which were based on known (23–29) or estimated (30, 31) fetal venous plasma concentrations. This might have overestimated physiologically relevant responses, and effects observed at pharmacological doses should be interpreted with caution.

Another limitation of this study is the heterogeneity of the participants, especially in terms of pre-delivery gestational age at which *in vivo* measurements of placental function were opportunistically recorded (and placentas obtained for *ex vivo* experiments), as well as medication dosage and timing of treatments. The length of exposure to medications varied from one week to more than one year before birth between groups, which may have had a less or more pronounced effect on CPA reactivity. For instance, the longest exposure to bisoprolol may have been more effective in downregulating endothelial paracrine signalling and/or smooth muscle cell responsivity compared to the shorter exposure to labetalol (Table S3).

Furthermore, in this study it was not possible to infer any associations between pregnancy outcomes and antihypertensive medication exposure due to significant prescriber bias and clinical presentations influencing prescribing decisions.

In conclusion, whilst cardiovascular disease in pregnancy is common the effect of antihypertensive and cardiac medications on placental vascular function remains poorly understood. Our study has confirmed the association between maternal hypertensive disorders and altered placental vascular function but has also identified differential effects of different classes of vasoactive medications in direct exposure experiments, but more importantly in vessels investigated *ex vivo* following exposure during pregnancy. Bisoprolol was associated with the most pronounced effects, justifying the need for further mechanistic studies to better inform the risk benefit discussion around the use of this medication in pregnancy.

Perspectives

Women with hypertension and cardiovascular disease require medication to optimise maternal safety in pregnancy. Although, medications such as labetalol, nifedipine and bisoprolol are recommended to treat maternal hypertension and to minimise cardiac workload in women with cardiac disease, to date there is a lack of robust safety data to support evidence-based prescribing. We have demonstrated a significant association between maternal hypertension and a reduction in vasoreactivity of the fetoplacental vasculature which was not significantly altered by *in vivo* exposure to different antihypertensive medication classes. In addition, fetoplacental arteries isolated from normal pregnancies exposed to vasoactive drugs *ex vivo* demonstrated, that whilst CCBs are vasodilatory within the fetoplacental vasculature, adrenergic receptor agonists and antagonists have markedly different vascular effects. Fetoplacental arteries exposed to bisoprolol both *in vivo* and *ex vivo*, exhibited a vasoconstrictive phenotype. Further studies are justified to investigate the mechanisms of action of vasoactive medications in fetoplacental arteries and to understand how vasoactive mechanisms may differ from the systemic vasculature. Clinical studies are also needed to correlate *ex vivo* effects with maternal and fetal outcomes.

Declarations

Sources of Funding

This work was supported by a British Heart Foundation Project Grant to Paul Brownbill (PG/19/25/34301).

Disclosures

None.

Author Contribution

T.T. and P.B. conceived and designed the experiments. T.T., E.C.C., R.W., N.K., J.E.M. and P.B. acquired, analysed or interpreted experimental data. T.T., E.C.C., E.D.J., J.E.M. and P.B. drafted and/or critically

revised the manuscript for important intellectual content. All authors read and approved the final version of the manuscript.

Acknowledgement

We are grateful to all of the women who donated their placentas to research. We thank the staff in the Maternal and Fetal Health Research Centre for their assistance with patient recruitment, tissue and clinical data collection, particularly Jessica Morecroft.

Data Availability

The data supporting the findings of this study are available from the corresponding author upon reasonable request. The detailed method, including criteria for the study, can be found in the Supplemental Material.

References

1. Saving Lives, M. B. R. R. A. C. E. U. K. Improving Mothers' Care 2024 - Lessons learned to inform maternity care from the UK and Ireland Confidential Enquiries into Maternal Deaths and Morbidity 2020-22 October 2024 [Available from: <https://www.npeu.ox.ac.uk/mbrace-uk/reports/maternal-reports/maternal-report-2020-2022>]
2. Sanghavi, M. & Rutherford, J. D. Cardiovascular physiology of pregnancy. *Circulation* **130** (12), 1003–1008 (2014).
3. Hypertension in pregnancy: diagnosis and management. (2019).
4. Magee, L. A. et al. The 2021 International Society for the Study of Hypertension in Pregnancy classification, diagnosis & management recommendations for international practice. *Pregnancy Hypertens.* **27**, 148–169 (2022).
5. Magee, L. A. et al. Less-tight versus tight control of hypertension in pregnancy. *N Engl. J. Med.* **372** (5), 407–417 (2015).
6. Tita, A. T. et al. Treatment for Mild Chronic Hypertension during Pregnancy. *N Engl. J. Med.* **386** (19), 1781–1792 (2022).
7. Webster, K. et al. Diagnosis and management of hypertension in pregnancy: summary of updated NICE guidance. *BMJ* **366**, l5119 (2019).
8. Ramlakhan, K. P. et al. Perinatal outcomes after in-utero exposure to beta-blockers in women with heart disease: Data from the ESC EORP registry of pregnancy and cardiac disease (ROPAC). *Int. J. Cardiol.* **410**, 132234 (2024).
9. Ormesher, L. et al. Prevalence of pre-eclampsia and adverse pregnancy outcomes in women with pre-existing cardiomyopathy: a multi-centre retrospective cohort study. *Sci. Rep.* **13** (1), 153 (2023).

10. Khedun, S. M., Maharaj, B. & Moodley, J. Effects of antihypertensive drugs on the unborn child: what is known, and how should this influence prescribing? *Paediatr. Drugs.* **2** (6), 419–436 (2000).
11. Meidahl Petersen, K. et al. β -Blocker treatment during pregnancy and adverse pregnancy outcomes: a nationwide population-based cohort study. *BMJ Open.* ;**2**(4). (2012).
12. Cooper, E. J., Wareing, M., Greenwood, S. L. & Baker, P. N. Oxygen tension and normalisation pressure modulate nifedipine-sensitive relaxation of human placental chorionic plate arteries. *Placenta* **27** (4–5), 402–410 (2006).
13. Thomas, G. D. Neural control of the circulation. *Adv. Physiol. Educ.* **35** (1), 28–32 (2011).
14. Tropea, T. et al. Fetoplacental vascular effects of maternal adrenergic antihypertensive and cardioprotective medications in pregnancy. *J. Hypertens.* **41** (11), 1675–1687 (2023).
15. Duan, L., Ng, A., Chen, W., Spencer, H. T. & Lee, M. S. Beta-blocker subtypes and risk of low birth weight in newborns. *J. Clin. Hypertens. (Greenwich).* **20** (11), 1603–1609 (2018).
16. Sibai, B. M., Gonzalez, A. R., Mabie, W. C. & Moretti, M. A comparison of labetalol plus hospitalization versus hospitalization alone in the management of preeclampsia remote from term. *Obstet. Gynecol.* **70** (3 Pt 1), 323–327 (1987).
17. Lydakis, C., Lip, G. Y., Beevers, M. & Beevers, D. G. Atenolol and fetal growth in pregnancies complicated by hypertension. *Am. J. Hypertens.* **12** (6), 541–547 (1999).
18. Ormesher, L., Vause, S. & Higson, S. al. e. Prevalence of pre-eclampsia and adverse pregnancy outcomes in women with pre-existing cardiomyopathy: a multi-centre retrospective cohort study. *Authorea.* (2022).
19. Tanaka, K. et al. Beta-Blockers and Fetal Growth Restriction in Pregnant Women With Cardiovascular Disease. *Circ. J.* **80** (10), 2221–2226 (2016).
20. Hoeltzenbein, M. et al. Pregnancy Outcome After First Trimester Use of Methyldopa: A Prospective Cohort Study. *Hypertension* **70** (1), 201–208 (2017).
21. Ashworth, D. et al. A treatment strategy with nifedipine versus labetalol for women with pregnancy hypertension: study protocol for a randomised controlled trial (Giant PANDA). *Trials* **24** (1), 584 (2023).
22. Tropea, T. et al. Nitrite mediated vasorelaxation in human chorionic plate vessels is enhanced by hypoxia and dependent on the NO-sGC-cGMP pathway. *Nitric Oxide.* **80**, 82–88 (2018).
23. Saotome, T. et al. Labetalol in hypertension during the third trimester of pregnancy: its antihypertensive effect and pharmacokinetic-dynamic analysis. *J. Clin. Pharmacol.* **33** (10), 979–988 (1993).
24. Fischer, J. H. et al. Influence of gestational age and body weight on the pharmacokinetics of labetalol in pregnancy. *Clin. Pharmacokinet.* **53** (4), 373–383 (2014).
25. Filgueira, G. C. O. et al. Effect of type 2 diabetes mellitus on the pharmacokinetics and transplacental transfer of nifedipine in hypertensive pregnant women. *Br. J. Clin. Pharmacol.* **83** (7), 1571–1579 (2017).

26. Silberschmidt, A. L., Kühn-Velten, W. N., Juon, A. M., Zimmermann, R. & von Mandach, U. Nifedipine concentration in maternal and umbilical cord blood after nifedipine gastrointestinal therapeutic system for tocolysis. *Bjog* **115** (4), 480–485 (2008).
27. Morgan, J. L. et al. Pharmacokinetics of amlodipine besylate at delivery and during lactation. *Pregnancy Hypertens.* **11**, 77–80 (2018).
28. Versmissen, J. et al. Doxazosin treatment of phaeochromocytoma during pregnancy: placental transfer and disposition in breast milk. *Br. J. Clin. Pharmacol.* **82** (2), 568–569 (2016).
29. Jones, H. M., Cummings, A. J., Setchell, K. D. & Lawson, A. M. A study of the disposition of alpha-methyldopa in newborn infants following its administration to the mother for the treatment of hypertension during pregnancy. *Br. J. Clin. Pharmacol.* **8** (5), 433–440 (1979).
30. Ulu, S. T. & Aydoğmuş, Z. An HPLC method for the determination of bisoprolol in human plasma and its application to a pharmacokinetic study. *J. Chromatogr. Sci.* **50** (7), 615–619 (2012).
31. Nandania, J. et al. Quantitative determination of nebivolol from human plasma using liquid chromatography-tandem mass spectrometry. *J. Chromatogr. B Analyt Technol. Biomed. Life Sci.* **923–924**, 110–119 (2013).
32. Bai, X. Y. et al. Effect of amlodipine in human internal mammary artery and clinical implications. *Ann. Thorac. Surg.* **90** (6), 1952–1957 (2010).
33. Erkinaro, T. et al. Placental and fetal hemodynamics after labetalol or pindolol in a sheep model of increased placental vascular resistance and maternal hypertension. *Reprod. Sci.* **16** (8), 749–757 (2009).
34. Harper, A. & Murnaghan, G. A. Maternal and fetal haemodynamics in hypertensive pregnancies during maternal treatment with intravenous hydralazine or labetalol. *Br. J. Obstet. Gynaecol.* **98** (5), 453–459 (1991).
35. Petersen, O. B., Skajaa, K., Svane, D., Gregersen, H. & Forman, A. The effects of dihydralazine, labetalol and magnesium sulphate on the isolated, perfused human placental cotyledon. *Br. J. Obstet. Gynaecol.* **101** (10), 871–878 (1994).
36. David, R., Leitch, I. M., Read, M. A., Boura, A. L. & Walters, W. A. Actions of magnesium, nifedipine and clonidine on the fetal vasculature of the human placenta. *Aust N Z. J. Obstet. Gynaecol.* **36** (3), 267–271 (1996).
37. Karahanoglu, E. et al. Nifedipine increases fetoplacental perfusion. *J. Perinat. Med.* **45** (1), 51–55 (2017).
38. Furuhashi, N., Tsujiei, M., Kimura, H., Nagae, H. & Yajima, A. Effects of nifedipine on placental blood flow, placental weight and fetal weight in normotensive and spontaneously hypertensive rats. *Gynecol. Obstet. Invest.* **34** (4), 193–196 (1992).
39. Sanusi, A. A. et al. Pregnancy Outcomes of Nifedipine Compared With Labetalol for Oral Treatment of Mild Chronic Hypertension. *Obstet. Gynecol.* **144** (1), 126–134 (2024).
40. Korkmaz, O. et al. Labetalol, nebivolol, and propranolol relax human radial artery used as coronary bypass graft. *J. Thorac. Cardiovasc. Surg.* **149** (4), 1036–1040 (2015).

41. Conner, C. S. Labetalol: an alpha- and beta-blocker. *Drug Intell. Clin. Pharm.* **17** (7–8), 543–544 (1983).
42. Resch, B. E., Ducza, E., Gáspár, R. & Falkay, G. Role of adrenergic receptor subtypes in the control of human placental blood vessels. *Mol. Reprod. Dev.* **66** (2), 166–171 (2003).
43. Brodde, O. E. Bisoprolol (EMD 33512), a highly selective beta 1-adrenoceptor antagonist: in vitro and in vivo studies. *J. Cardiovasc. Pharmacol.* **8** (Suppl 11), S29–35 (1986).
44. Taylor, S. H. Pharmacotherapeutic stature of doxazosin and its role in coronary risk reduction. *Am. Heart J.* **116** (6 Pt 2), 1735–1747 (1988).
45. Ritter, J. M. Nebivolol: endothelium-mediated vasodilating effect. *J. Cardiovasc. Pharmacol.* **38** (Suppl 3), S13–S16 (2001).
46. Dessy, C. et al. Endothelial beta3-adrenoreceptors mediate nitric oxide-dependent vasorelaxation of coronary microvessels in response to the third-generation beta-blocker nebivolol. *Circulation* **112** (8), 1198–1205 (2005).
47. Angulo, J. et al. Nebivolol dilates human penile arteries and reverses erectile dysfunction in diabetic rats through enhancement of nitric oxide signaling. *J. Sex. Med.* **7** (8), 2681–2697 (2010).
48. Broeders, M. A. et al. Nebivolol: a third-generation beta-blocker that augments vascular nitric oxide release: endothelial beta(2)-adrenergic receptor-mediated nitric oxide production. *Circulation* **102** (6), 677–684 (2000).
49. Bueno-Pereira, T. O., Nunes, P. R., Matheus, M. B., Vieira da Rocha, A. L. & Sandrim, V. C. Nebivolol Increases Nitric Oxide Synthase via $\beta(3)$ Adrenergic Receptor in Endothelial Cells Following Exposure to Plasma from Preeclamptic Patients. *Cells* ;**11**(5). (2022).
50. Versari, D., Daghini, E., Viridis, A., Ghiadoni, L. & Taddei, S. Endothelium-dependent contractions and endothelial dysfunction in human hypertension. *Br. J. Pharmacol.* **157** (4), 527–536 (2009).
51. Touyz, R. M. et al. Vascular smooth muscle contraction in hypertension. *Cardiovasc. Res.* **114** (4), 529–539 (2018).
52. Al Khalaf, S. Y. et al. Impact of Chronic Hypertension and Antihypertensive Treatment on Adverse Perinatal Outcomes: Systematic Review and Meta-Analysis. *J. Am. Heart Assoc.* **10** (9), e018494 (2021).
53. Hoeltzenbein, M. et al. Pregnancy outcome after first trimester exposure to bisoprolol: an observational cohort study. *J. Hypertens.* **36** (10), 2109–2117 (2018).
54. Kayser, A. et al. Neonatal effects of intrauterine metoprolol/bisoprolol exposure during the second and third trimester: a cohort study with two comparison groups. *J. Hypertens.* **38** (2), 354–361 (2020).
55. Chappell, J. et al. A review of feto-placental vasculature flow modelling. *Placenta* **142**, 56–63 (2023).
56. Burton, G. J. & Jauniaux, E. Pathophysiology of placental-derived fetal growth restriction. *Am. J. Obstet. Gynecol.* **218** (2s), S745–s61 (2018).

57. Schneider, H. Placental Dysfunction as a Key Element in the Pathogenesis of Preeclampsia. *Dev. Period Med.* **21** (4), 309–316 (2017).

Tables

Table 1 is available in the Supplementary Files section.

Figures




Figure 1

Contrasting effects of different antihypertensive and cardioprotective medications on chorionic plate arteries *ex vivo*. Cumulative concentration-response curves for (A) amlodipine, (B) labetalol, (C) methyldopa, (D) doxazosin, (E) nebivolol and (F) bisoprolol vs their relative diluent controls on pre-constricted fetoplacental arteries. Concentrations shown within the dotted columns represent the

predicted in vivo dose exposure. Curves are compared by 2-way ANOVA; P values are the overall effect vs control. N=8-10 per group. *P < 0.05, ***P < 0.001, ****P < 0.0001; Šídák's multiple comparison test.




Figure 2

Effects of hypertension and Ca²⁺ channel blocker exposure in women with hypertension on chorionic plate artery vascular responses. Vasoconstriction in response to 120 mM KCl (A, D), and U46619 (B, E); vasorelaxation in response to sodium nitroprusside (SNP; C, F), in chorionic plate arteries isolated from the placenta of women with normal pregnancy (NP), unmedicated hypertension (HTN) and medicated with either amlodipine (A, B, C) or nifedipine (D, E, F). Groups are compared by 1-way ANOVA (A, D) or 2-way ANOVA (B, C, E, F); P values on the graphs are the overall effect. #P < 0.05, ###P < 0.001, ****P < 0.0001, HTN vs NP; *P < 0.05, **P < 0.01, amlodipine or nifedipine vs NP; Tukey's multiple comparison test.




Figure 3

Effect of labetalol exposure in women with hypertension on chorionic plate arteries. Vasoconstriction in response to (A) 120 mM KCl, and (B) U46619; (C) vasorelaxation in response to sodium nitroprusside (SNP), in chorionic plate arteries isolated from the placenta of women with normal pregnancy (NP), hypertension unmedicated (HTN), and medicated with labetalol. Groups are compared by 1-way ANOVA (A) or 2-way ANOVA (B, C); P values on the graphs are the overall effect. #P < 0.05, ###P < 0.001, ####P < 0.0001, HTN vs NP; *P < 0.05, ***P < 0.001, labetalol vs NP; Tukey's multiple comparison test.




Figure 4

Effect of labetalol in combination with Ca²⁺ channel blocker therapy exposure in women with hypertension on chorionic plate arteries. Vasoconstriction in response to (A) 120 mM KCl, and (B) U46619; (C) vasorelaxation in response to sodium nitroprusside (SNP), in chorionic plate arteries isolated from the placenta of women with normal pregnancy (NP), unmedicated hypertension (HTN), and hypertension medicated with labetalol and Ca²⁺ channel blockers (CCBs, amlodipine, nifedipine). Groups are compared by 1-way ANOVA (A) or 2-way ANOVA (B, C); P values on the graphs are the overall effect. #P < 0.05, ###P < 0.001, #####P < 0.0001, HTN vs NP; **P < 0.01, labetalol + CCBs vs NP; Tukey's multiple comparison tests.




Figure 5

Effect of bisoprolol exposure in women with cardiac dysfunction on chorionic plate arteries.

Vasoconstriction in response to (A) 120 mM KCl, and (B) U46619; (C) vasorelaxation in response to sodium nitroprusside (SNP), in chorionic plate arteries isolated from the placenta of women with normal pregnancy (NP) and those with cardiac dysfunction medicated with bisoprolol. Groups are compared by unpaired t-test- (A) or 2-way ANOVA (B, C); P values on the graphs are the overall effect. *P<0.05, **P < 0.01, bisoprolol vs NP; Šídák's multiple comparisons test.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [TropeaetalresearchpaperSupplementalMaterialforsubmissioninSciReports.docx](#)
- [Table1.docx](#)