

Supplementary Information

Ciprofloxacin remediation by photolysis, TiO₂ photocatalysis and high-voltage electrical discharges: kinetics, energy demand and ecotoxicity

Cristèle Delsart, Maryline Pflieger, Urška Lavrenčič Štangar, Nadia Boussetta, Marwa Brahim, Aurélien Trivella

Journal: Clean Technologies and Environmental Policy

Article type: Original research paper

Corresponding author: Cristèle Delsart, c.delsart@istom.fr


Affiliations: UPR ADI-Suds, ISTOM, F-49000 Angers, France; University of Nova Gorica, SI-5001 Nova Gorica, Slovenia; University of Ljubljana, Faculty of Chemistry and Chemical Technology, SI-1000 Ljubljana, Slovenia; Université de Technologie de Compiègne, ESCOM, TIMR, 60203 Compiègne, France; Univ. Bordeaux, CNRS, Bordeaux INP, EPOC, UMR 5805, F-33600 Pessac, France.

Contents

Online Resource 1 Kinetics for CIP degradation as a function of reaction time obtained without light and with light in synthetic freshwater.

Online Resource 2 Bioluminescence inhibition of *Vibrio fischeri* (at 15 min) as a function of treatment time (a) and CIP degradation (b) for different degradation processes.


Online Resource 1




Kinetics for CIP degradation as a function of reaction time obtained without light and with light in synthetic freshwater.

Online Resource 2

(a)



(b)



Bioluminescence inhibition of *Vibrio fischeri* (at 15 min) as a function of treatment time (a) and CIP degradation (b) for different degradation processes.