

Table 3. Treatment conditions as derived from the 14 cases of pediatric Behcet's disease obtained from the literature.

No.	Hormone therapy (methylprednisolone pulse)	Immune suppressants	Biologics	Anticoagulant therapy	Surgical treatment
1#	Y (Y)	CTX, AZA	Remicade	Y	N
2#	Y	CTX, colchicine	N	Y	N
3#	Y	CTX, CsA, MMF, colchicine	daclizumab, anakinra	Y	Mitral valve repair and replacement, aneurysm resection, orthotopic heart transplantation
4#	Y	MMF, colchicine, MTX	N	Y	Surgical thrombectomy
5#	Y	/	/	/	Aortic valvuloplasty with autologous pericardium
6#	Y	N	N	Y	Surgical thrombectomy
7#	Y	Colchicine, AZA, MTX, MMF	N	Y	Surgical resection and coronary artery bypass graft
8#	Y	CTX	Remicade	Y	Surgical resection
9#	Y	CTX	N	Y	N
10#	Y (Y)	CTX, AZA	N	Y	N
11#	Y	CTX	N	Y	Surgical resection
12#	Y (Y)	Colchicine	N	Y	N
13#	Y (Y)	CTX, colchicine	N	Y	N
14#	Y (Unknown)	MMF	N	/	Surgical resection

Notes: CTX: cyclophosphamide; AZA: azathioprine; CsA: cyclosporine; MMF: mycophenolate mofetil; MTX: methotrexate; N:no; Y:yes; /: Condition unknown, statistics unavailable.