

Ecophysiology of soursop seedlings fertilized with calcium silicate under different irrigation frequencies

Patrycia Elen Costa Amorim

Federal Rural University of Semi-Arid

Vander Mendonça

Federal Rural University of Semi-Arid

Thiago dos Reis

Federal Rural University of Semi-Arid

Fagner Nogueira Ferreira

Federal Rural University of Semi-Arid

David Emanuel Gomes da Silva

Federal Rural University of Semi-Arid

João Luíz Lima

Federal Rural University of Semi-Arid

Rodrigo Rafael da Silva

Federal Rural University of Semi-Arid

Matheus Augusto Silva

Federal Rural University of Semi-Arid

Adriana dos Santos Ferreira

Federal Rural University of Semi-Arid

João Everthon da Silva Ribeiro

Federal Rural University of Semi-Arid

Agda Malany Forte de Oliveira

State University of Paraíba - UEPB

Gabriel Cardoso da Silva

Federal Rural University of Semi-Arid

Elaine Cristina Marinho Vieira

Federal Rural University of Semi-Arid

Luilson Pinheiro da Costa

Federal Rural University of Semi-Arid

Antônio Gustavo de Luna Souto

Federal University of Paraíba - UFPB

Francisco Vanies da Silva Sá

vanies_agronomia@hotmail.com

State University of Paraíba - UEPB, Catolé do Rocha

Research Article

Keywords: Annona muricata L, CaSiO₃, irrigation interval, water stress

Posted Date: October 14th, 2026

DOI: <https://doi.org/10.21203/rs.3.rs-10176509/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Additional Declarations: No competing interests reported.

Abstract

Propose: The soursop tree (*Annona muricata* L.) is a tropical fruit tree cultivated in the semi-arid region of Brazil; however, it is sensitive to water deficit. The application of calcium silicate can promote plant growth and development under water stress. Therefore, the objective was to evaluate the ecophysiological and biochemical changes in soursop seedlings fertilized with calcium silicate under different irrigation frequencies.

Methods: The experiment was conducted in a greenhouse using a randomized block design with a 4 × 2 factorial scheme and four replications. The treatments consisted of four irrigation frequencies: TR1 – daily irrigation; TR2 – irrigation every two days; TR3 – irrigation every four days; TR4 – irrigation every six days, combined with the presence or absence of calcium silicate (CaSiO₃), applied via soil, at a dose of 3.5 g plant⁻¹.

Results: Soursop seedlings subjected to TR3 associated with CaSiO₃ showed improved gas exchange, including transpiration, stomatal conductance, and photosynthetic rate. Fertilization with CaSiO₃ regulated proline synthesis, increased leaf biomass and silicon and calcium content in leaf tissues, conferring greater tolerance to water deficit. Treatments TR2 and TR3 were similar to TR1 in plant height, number of leaves, leaf area, biomass accumulation, chlorophyll a fluorescence, chlorophyll content, and total soluble sugars, whereas TR4 significantly reduced these variables, regardless of CaSiO₃ fertilization.

Conclusions: The application of CaSiO₃ under moderate stress (TR3) increases soursop seedlings' tolerance to deficit irrigation regimes.

Highlights

Calcium silicate (CaSiO) alleviates the effects of moderate water deficit in soursop seedlings.

A six-day irrigation interval severely suppresses height, leaf emission, and biomass accumulation in soursop seedlings.

A four-day irrigation frequency combined with (CaSiO) enhances leaf gas exchange parameters in soursop seedlings.

1. Introduction

The soursop tree (*Annona muricata* L.) is a tropical fruit tree native to Central and South America. It is currently cultivated worldwide, including in Africa and Southeast Asia, and particularly in the semi-arid region of Brazil [1, 2]. This crop, formerly neglected or underutilized, has great medicinal potential and contributes to food security due to the high nutritional and caloric value of its fruits [3, 4]. Its fruits, when ripe, are consumed fresh and, after processing, are used to produce juices, jams, ice cream, sweets, nectars, yogurts, syrups, drinks, and shakes [5, 4]. Furthermore, the fruits, leaves, seeds, and stems exhibit important biological and pharmacological activities, including antibacterial, antiviral, antifungal, antitumor, anthelmintic, analgesic, anti-arthritic, hypotensive, anti-inflammatory, antidiabetic, antiparasitic, and pesticidal properties [1, 3, 6].

In Brazil, soursop is widely cultivated in the Northeast region, where it helps supplement family farmers' incomes. In this context, the fruit sector plays a crucial role in the economy of the semi-arid Northeast, both by supplying local and regional markets and by leveraging its export potential. The Northeast region accounts for approximately 27% of national fruit production, positioning it as one of the country's main fruit-growing centers [7]. However, soursops require adequate water availability to grow and sustain good yields, and their growth, development, and productivity are affected under water-deficit conditions [5, 3, 8]. Thus, producing seedlings of the species in the semi-arid areas of Brazil is challenging, as rainfall events are concentrated in a short, irregular period, making it highly relevant to understand this dynamic for agricultural management [9].

In semi-arid conditions, farmers are often compelled to adopt deficit irrigation due to water scarcity. This practice is a potential strategy to help crops withstand mild or moderate water deficits, although it impairs productivity [10, 11]. Water deficit is one of the main abiotic stresses in agriculture, affecting about 30% of the world's cultivated area [12]. This stress can severely affect the morphological, physiological, biochemical, and molecular attributes of the plant, reducing photosynthetic capacity and stomatal conductance, which increases the internal concentration of CO₂ and biomass production [13–16]. Furthermore, it causes excessive production of reactive oxygen species (ROS) [15], a decrease in the plant's water potential and relative water content in tissues, and physiological drought [17]. These effects impair plant growth, development, productivity, and quality [13, 16], thereby compromising agricultural sustainability and causing food shortages [18]. In this context, management strategies to increase plant tolerance to water deficit have been extensively studied [19].

In this context, deficit irrigation emerges not only as a consequence of water scarcity but also as a rational water management strategy, aiming to mitigate the impacts of water stress on crop performance. This practice is a smart and common strategy for reducing water consumption and improving sustainable agricultural management [20, 21]. Experimental evidence indicates that deficit irrigation, a technique that intentionally reduces water supply below the crop's total water needs, has high potential for water savings and is being progressively adopted in regions worldwide [22]. According to Li et al. [23], deficit irrigation is the controlled use of limited water resources to maintain stable productivity, with irrigation frequency being crucial for plant growth and productivity [24]. In recent years, studies have shown that deficit irrigation, defined as reduced irrigation frequency, can offer significant benefits under limited water availability [25, 26, 22].

However, even with rational irrigation management, plants remain subject to water stress, especially in the early stages of development, which drives the search for complementary mitigation strategies. In this context, the use of calcium silicate (CaSiO₃) has proven to be a promising agent for mitigating abiotic stresses, such as water deficit [27, 28, 12]. Silicon (Si) maintains leaf water potential, water content, and photosynthetic pigment content; promotes osmotic adjustments and reduces oxidative stress [29, 30]; and regulates gas exchange, photochemical yields, and growth [31, 14]. Furthermore, Si increases nutrient uptake and biomass production [29, 32].

While calcium (Ca²⁺) plays a fundamental role in signal transduction in response to environmental stress [33], Malyukova et al. [34] indicated that the influx and cytoplasmic levels of Ca²⁺ are important for abscisic acid (ABA) transduction in guard cells. At the same time, ABA can regulate stomatal opening in these cells. Ca²⁺ also plays an essential role in maintaining the stability and integrity of cellular structure [35], thereby favoring photosynthetic capacity, alleviating oxidative damage, and promoting osmolyte accumulation [36, 33].

Previous studies report that the application of calcium silicate increases drought tolerance in various species, including *Oryza sativa* [37], *Saccharum officinarum* [38], *Cucumis sativus* L. [39], and *Zea mays* L. [40]. However, the benefits of Si and Ca depend on soil water availability, environmental conditions, and species-specific physiological characteristics [33, 41].

Therefore, although the beneficial effects of silicon and calcium are known in several species, the interaction between these elements, as demonstrated by the application of calcium silicate to soursop seedlings subjected to varying irrigation frequencies, remains underexplored. A gap remains regarding how this fertilization influences osmotic adjustment and the efficiency of the photosynthetic apparatus when the interval between waterings is extended. Thus, it is hypothesized that the application of CaSiO₃ maintains stomatal function and osmotic balance under moderate water stress, promoting greater physiological efficiency in soursop seedlings. Therefore, the objective was to evaluate the ecophysiological and biochemical changes in soursop seedlings fertilized with calcium silicate under different irrigation frequencies.

2. Materials and Methods

2.1 Location and Characterization of the Experimental Area

The experiment was conducted in a greenhouse in the seedling production sector of the Department of Agronomic and Forestry Sciences at the Federal Rural University of the Semi-Arid Region (UFERSA), in Mossoró, Rio Grande do Norte State, Brazil (5°12'14.2"S and 37°19'29.3"W, with an average altitude of 18 m), from September to December 2023. The greenhouse is protected by an arched tunnel-type roof, measuring 7.0 m wide, 18.0 m long, and 3.0 m high, covered with a transparent, low-density polyethylene sheet, 0.15 mm thick. The structure is made of galvanized steel, and the side and front walls are made of 50% shade cloth.

The region's climate is of the BSwH type (hot and dry), with an average precipitation of 555 mm year⁻¹, an average annual temperature of 27.8°C, and an average relative humidity of 68.9% [42, 43]. The climatic data for temperature and relative humidity during the experiment were recorded using a digital thermometer-hygrometer (SH-122, J-Prolab, Brazil) and are presented in Fig. 1.

2.2 Plant Material, Experimental Design, and Treatments

The experimental design used was a randomized block design, in a 4 × 2 factorial scheme, with four replications and one seedling per replication, totaling 32 experimental units. The first factor consisted of four irrigation frequencies: TR1 – daily irrigation; TR2 – every two days; TR3 – every four days; and TR4 – every six days. The second factor corresponded to the presence or absence of calcium silicate (CaSiO₃) applied to the soil at a dose of 3.5 g plant⁻¹. The irrigation schedules were based on the studies of Fredrick, Bamalo, and Chima [44] and Yisau et al. [3], and the fertilization with CaSiO₃ followed the recommendations of Souza et al. [45].

Seeds of *A. muricata* L. (cultivar Morada), which exhibit high productivity, large fruits, good health, and physical quality, as described by Pinto and Silva [46] and by São José et al. [47], were obtained from mature fruits with good phytosanitary appearance, from trees located in the rural area of Mossoró, Rio Grande do Norte, Brazil. The seeds were manually extracted, washed in running water to remove pulp residues, and dried in the shade on paper for 7 days. Sowing was carried out in 2 dm³ plastic bags filled with sand and Tropstrato® commercial substrate at a 2:1 (v/v) ratio, using three seeds at a depth of 1.5 cm, with thinning performed 20 days after emergence, retaining only one seedling per bag. The commercial substrate Tropstrato®, composed of single superphosphate, potassium nitrate, peat, vermiculite, and pine bark, presents the following chemical attributes: pH in CaCl₂ = 5.75; P = 65.70 mg dm⁻³; K⁺ = 1.60 cmol_c dm⁻³; Ca²⁺ = 23.80 cmol_c dm⁻³; Mg²⁺ = 12.40 cmol_c dm⁻³; Al³⁺ = 0.00 cmol_c dm⁻³; H + Al = 4.20 cmol_c dm⁻³; Sum of bases = 37.80 cmol_c dm⁻³; Cation Exchange Capacity = 42.10 cmol_c dm⁻³ and Base Saturation = 89.8%, according to the manufacturer's data.

One hundred and thirty days after planting, *A. muricata* L. seedlings were transplanted into 5 L plastic pots, spaced 1.0 m apart between rows and 0.5 m between plants. The containers were filled with a substrate composed of sand and Tropstrato® at a 2:1 (v/v) ratio, and the mass of each pot was recorded for standardization. The seedlings were irrigated daily until the water retention capacity (WRC), corresponding to 100% soil moisture, was reached over 30 days of acclimatization before the application of the treatments. The chemical properties of the substrate at the beginning and end of the experiment are detailed in Table 1.

Table 1
Chemical characteristics of the substrate before and at the end of the experiment

Initial characterization													
	pH	EC	O.M	P	K ⁺	Na ⁺	Ca ²⁺	Mg ²⁺	SB	CEC	Si	V	ESP
	(H ₂ O) 1:25	(dS m ⁻¹) 1:25	(g kg ⁻¹)	(mg dm ⁻³)	cmol _c dm ⁻³						(mg kg ⁻¹)	(%).....	
	4.82	1.0	27.84	214.31	0.58	1.09	5.60	2.60	9.87	9,87	83.68	100	11
Characterization after treatments													
TR1 absence CaSiO ₃	5.91	0.70	8.89	177.70	0.58	2.14	3.70	1.30	7.71	7.71	125.30	100	28
TR2 absence CaSiO ₃	5.51	1.08	11.21	245.00	0.55	2.05	4.70	2.00	9.30	9.30	104.60	100	22
TR3 absence CaSiO ₃	5.46	0.81	11.99	256.90	0.58	1.82	4.30	2.20	8.90	8.90	98.90	100	20
TR4 absence CaSiO ₃	5.33	0.83	11.60	284.20	0.71	1.55	4.50	1.30	8.05	8.05	76.75	100	19
TR1 Presence CaSiO ₃	5.45	0.91	7.35	325.93	0.50	2.00	4.60	1.60	8.70	8.70	98.05	100	23
TR2 Presence CaSiO ₃	5.44	0.84	17.01	243.30	0.55	1.91	4.40	1.30	8.16	8.16	89.67	100	23
TR3 Presence CaSiO ₃	5.40	0.80	18.56	293.60	0.63	1.82	4.20	1.60	8.25	8.25	101.60	100	22
TR4 Presence CaSiO ₃	5.48	0.67	17.79	250.95	0.58	1.73	4.20	2.00	8.50	8.50	109.90	100	20
TR1: daily irrigation; TR2: irrigation every two days; TR3: irrigation every four days; and TR4: irrigation every six days. Absence of CaSiO ₃ and presence of CaSiO ₃ . Al ³⁺ , (H + Al) and m: 0, P, K ⁺ , Na ⁺ : Mehlich 1 extractor; Al ³⁺ , Ca ²⁺ , Mg ²⁺ : 1.0 mol L ⁻¹ KCl extractor; EC: electrical conductivity; SB: sum of bases; CEC: cation exchange capacity; O.M.: organic matter; ESP: exchangeable sodium percentage													

2.3 Irrigation and fertilization management

Thirty days after transplanting the seedlings (DAT), irrigation was initiated, according to the irrigation schedule established for each treatment, using the local water supply (Table 2). Irrigation was carried out manually with a graduated cylinder, once a day in the afternoon. During this process, the water retention capacity (WRC) of the substrate was maintained at 100% using the gravimetric method [48].

Table 2
Chemical attributes of the local water supply are used for irrigating soursop seedlings

pH	EC	K ⁺	Na ⁺	Mg ²⁺	Ca ²⁺	Cl ⁻	CO ₃ ²⁻	HCO ₃ ⁻	SAR	Si
H ₂ O	dS m ⁻¹	mmol _c L ⁻¹								mg L ⁻¹
7.50	0.58	0.26	4.59	1.46	0.74	2.40	0.10	3.50	4.4	5.00
pH (H ₂ O): Hydrogen ion potential in water; EC: Electrical conductivity; K ⁺ : potassium; Na ⁺ : sodium; Mg ²⁺ : magnesium; Ca ²⁺ : calcium; Cl: chlorine; CO ₃ ²⁻ : carbonate; HCO ₃ ⁻ : bicarbonate; SAR: sodium adsorption ratio; Si: silicon										

To accurately determine the required water volume, all pots were weighed every 30 days. Subsequently, the pots were placed at maximum saturation and covered with plastic at the top opening to prevent water evaporation. After 24 h, the pots were drained of excess water, indicating that the substrate had reached maximum WRC, and were weighed again. Two reference pots were selected per treatment and served as the basis for calculating the volume of water to be applied during each irrigation event, enabling precise adjustments to maintain the target WRC, following the methodology adapted from Souza et al. [48]

The applied volume (VI) per pot was calculated as the difference between the average weights of the reference pots (PVR1 and PVR2) and their weights before the irrigation event (PVra1 and PVra2), as given by Eq. 1.

$$VI = \left(\frac{PVr1 + PVr2}{2} \right) - \left(\frac{PVra1 + PVra2}{2} \right)$$

1

Where:

VI = Volume of water to be applied in the irrigation event (L);

PVR1 = Weight of reference pot 1 (kg);

PVR2 = Weight of reference pot 2 (kg);

PVra1 = Weight of reference pot before irrigation event 1 (kg);

PVra2 = Weight of reference pot before irrigation event 2 (kg).

The total volume of irrigation applied to the soursop seedlings across treatments was quantified throughout the experimental period (Table 3).

Table 3

Total irrigation volume in soursop seedlings fertilized with calcium silicate at different irrigation frequencies at 120 days

Irrigation volume (L plant ⁻¹)		
Fertilizing		
Irrigation frequencies	Absence CaSiO ₃	Presence CaSiO ₃
TR1	20.258	18.736
TR2	17.602	16.829
TR3	13.433	11.899
TR4	9.624	9.573
TR1: daily irrigation; TR2: irrigation every two days; TR3: irrigation every four days; and TR4: irrigation every six days. Absence of CaSiO ₃ and presence of CaSiO ₃		

Calcium silicate (CaSiO₃), composed of 10–25% calcium oxide (CaO) and 75–90% silicon dioxide (SiO₂), was applied to the soil around the stem of soursop seedlings at 30 DAT, at a dosage of 3.5 g per seedling, as recommended by Souza et al. [45]. Fertilization with nitrogen, potassium, and phosphorus (NPK) was split into 12 applications and supplied weekly via fertigation, corresponding to 50% of the recommended rates of 100 mg N, 300 mg P₂O₅, and 150 mg K₂O per dm⁻³ of soil, as recommended by Novais, Neves, and Barros [49]. For NPK fertilization, 400 mg of urea (45% N), 1,250 mg of monoammonium phosphate (MAP, 12% N and 50% P₂O₅), and 650 mg of potassium chloride (KCl, 60% K₂O) were applied per pot.

2.4 Variables analyzed

Growth and biomass production

120 days after treatment application, the soursop seedlings were evaluated for plant height (PH), stem diameter (SD), root volume (RV), number of leaves (NL), and leaf area (LA). PH was measured from the base to the insertion of the apical bud using a graduated tape measure (cm). SD was obtained at the substrate level using a digital caliper (mm). RV (cm³) was determined by the water displacement method, in which the root system, previously washed and dried, was immersed in a graduated cylinder filled with a known volume of water; the increase in water volume was then recorded, as described by Böhm [50]. NL was determined by direct counting of fully expanded leaves. The LA (cm²) was determined after collecting leaves from each plant, which were digitized using a flatbed scanner (Xpress SL-C480FW, Samsung Electronics, South Korea) at 600 dpi. The images obtained were subsequently processed and analyzed using ImageJ® software.

After growth analysis, the seedlings were collected and separated into leaves, stems, and roots, and washed in running water. The plant material was then placed in Kraft paper bags and heated in a forced-air oven at 65°C until a constant weight was reached after 72 hours. Subsequently, the samples were weighed on an analytical balance with a precision of 0.0001 g to determine the dry mass of each seedling part. The results were expressed in g.

Leaf gas exchange and chlorophyll a fluorescence

115 days after treatment application, leaf gas exchange measurements were performed using an infrared gas analyzer (IRGA; GFS–3000, Walz, Effeltrich, Germany). The equipment was programmed with an impeller speed of 7; a reading area of 8 cm²; a cuvette temperature of 25°C; a CO₂ concentration of 400 ppm; a flux rate of 750 µmol min⁻¹; and a photon flux density of 1,200 µmol m⁻² s⁻¹. Measurements were taken between 7:00 and 11:00 on the third fully expanded leaf, located

in the upper middle third of each plant. The parameters included: CO₂ assimilation rate (A_N - $\mu\text{mol m}^{-2} \text{s}^{-1}$), transpiration (E - $\text{mmol of H}_2\text{O m}^{-2} \text{s}^{-1}$), stomatal conductance (g_s - $\text{mol of H}_2\text{O m}^{-2} \text{s}^{-1}$), internal CO₂ concentration (C_i - $\mu\text{mol CO}_2 \text{mol}^{-1}$ of air), leaf temperature (T_l - °C), and vapor pressure deficit (VPD - kPa). From these data, water-use efficiency ($\text{WUE} = A_N/E$), intrinsic water-use efficiency ($\text{iWUE} = A_N/g_s$), and carboxylation efficiency (A_N/C_i) were calculated.

Chlorophyll *a* fluorescence analysis was performed between 7:00 and 11:00 on the same leaves used for gas exchange measurements. The leaves were subjected to 30 min of pre-adaptation to darkness using leaf tweezers. Subsequently, using a portable fluorometer (OS-30p, Opti-Sciences Chlorophyll Fluorometer, United States), the initial fluorescence (F_0) and maximum fluorescence (F_m) values were obtained. Based on these data, the variable fluorescence (F_v), the potential quantum efficiency of photosystem II (F_v/F_m), the primary maximum efficiency of photosystem II (F_v/F_0), and the basal quantum efficiency of the non-photochemical process of photosystem II (F_0/F_m) were calculated [51].

Photosynthetic pigments

Chlorophyll *a*, *b*, and total indices were determined at 120 DAT (days after transplanting) in the morning, on the middle third of the third fully expanded leaf from the seedling apex, using a portable chlorophyll meter (ClorofiLOG® CFL 1030, Falker, Brazil). The results were expressed as Falker Chlorophyll Index (FCI) [52].

Osmotic adjustment

At 113 days DAT, fresh leaves from the middle third of the seedlings were collected for analysis of total soluble sugars and proline. To prepare the crude extracts, 150 mg of fresh leaves from three replicates in each treatment were placed in Eppendorf tubes containing 1 mL of 80% alcohol for extraction. The tubes were placed in the digester for 2 min to grind the tissue completely, and the supernatant was collected. The supernatant was then centrifuged at 10,000 rpm for 5 min at 10°C to obtain a clear supernatant, stored at -80°C, and subsequently used for biochemical analyses.

Total soluble sugars (TSS) were determined by the anthrone method [53] and expressed in $\text{mg TSS g}^{-1} \text{MF}$. The proline (PRO) content was determined in the supernatant using the method described by Bates, Waldren, and Teare [54], and the results were expressed as $\text{mol PRO g}^{-1} \text{MF}$.

Leaf content of K, Ca, and Si

The dry biomass of the leaves was ground in a Wiley stainless-steel mill and stored in hermetically sealed containers for the determination of potassium (K) and calcium (Ca) concentrations using the sulfuric acid digestion method [55]. For this, 0.4 g of the dry material was transferred to digestion tubes, to which 2 mL of hydrogen peroxide (H_2O_2 - 30%), 4 mL of sulfuric acid (H_2SO_4 - 98%), and 1 g of the salt mixture were added. The samples were then slowly heated to 350°C (approximately 2 h) in the digestion block to obtain a light greenish liquid, and heating was continued until a colorless liquid was achieved (approximately 4 h). After cooling, the mixture was diluted with 50 mL of deionized water in a collection bottle using a graduated cylinder, and the extract was properly stored. The K concentration was determined by flame photometry after diluting 1 mL of the extract with 19 mL of deionized water.

For calcium determination, 1 mL of the diluted extract was transferred to test tubes, to which 19 mL of deionized water was added. Then, 1 mL of this new dilution was transferred to another test tube, to which 4 mL of lanthanum solution was added. The Ca concentration was quantified by atomic absorption spectrophotometry [55]. The Si concentration was determined at the Soil and Plant Analysis Laboratory (LASP) in Petrolina, Pernambuco, Brazil, according to the methodology of Furlani and Gallo [56]. The samples were dried, ground, and subjected to alkaline digestion to solubilize the silicon. Subsequently, the element was quantified using a colorimetric method, with reading in a spectrophotometer.

Seedling quality was assessed based on plant height (PH), stem diameter (SD), total dry mass (TDM), shoot dry mass (SDM), and root dry mass (RDM). Dickson's quality index (DQI) was calculated according to Dickson et al. [57], as shown in

Eq. 2.

$$DQI = \frac{TPM - DM}{SB} \quad (2)$$

Drought tolerance (DT) was evaluated by the total dry mass of plants subjected to water deficit (Absence of CaSiO₃ and Presence of CaSiO₃) during the different irrigation shifts (DMd) and by the total dry mass of the control treatment (TR1Absence of CaSiO₃ or TR1Presence of CaSiO₃) (DMc), and calculated using the formula proposed by Fischer et al. [58], as shown in Eq. 3.

$$DT = \frac{DMc}{DMd} \quad (3)$$

The drought sensitivity index (DSI) was calculated using the equation by Ali and El-Sadek [59], with the same parameters as in Eq. 4.

$$DSI = \frac{DMc - DMd}{DMc} \quad (4)$$

2.5 Statistical analysis

The data obtained were subjected to the Shapiro-Wilk normality test ($p \geq 0.05$) [60]. Subsequently, they were subjected to analysis of variance (ANOVA) using the F test ($p \leq 0.05$). In cases of significance for the isolated factors and their interaction, the Student's t-test ($p \leq 0.05$) was applied to the fertilization factor, and the Tukey test ($p \leq 0.05$) to the irrigation schedule factor, using the statistical software SISVAR®, version 5.3 [61].

3. Results

3.1 Growth

Analysis of variance indicated no significant interaction ($p > 0.05$) between irrigation frequencies and fertilization with calcium silicate (CaSiO₃) for stem diameter (SD), plant height (PH), number of leaves (NL), leaf area (LA), and root volume (RV) of soursop seedlings (Table 4 and Fig. 2). However, irrigation frequency significantly influenced PH, NL, LA, and RV ($p \leq 0.05$), whereas SD did not differ significantly between irrigation schedules ($p > 0.05$) (Table 4). Fertilization with CaSiO₃, in turn, did not show a significant effect on any of the growth variables analyzed ($p > 0.05$; Table 4).

Table 4

F-test, Tukey test, and Student's t-test were performed for the variables: stem diameter (SD), plant height (PH), number of leaves (NL), leaf area (LA), and root volume (RV) in soursop seedlings fertilized with calcium silicate under different irrigation frequencies, 120 days after transplanting

F-test (p-value)					
SV	SD (mm)	PH (cm)	NL	LA (cm ² /planta)	RV (cm ³)
Blocks	0.5754 ^{ns}	0.6648 ^{ns}	0.6766 ^{ns}	0.7472 ^{ns}	0.0969 ^{ns}
TR	0.1810 ^{ns}	0.0026 [*]	0.0060 [*]	0.0000 [*]	0.0001 [*]
CaSiO ₃	0.2663 ^{ns}	0.8391 ^{ns}	0.0891 ^{ns}	0.8950 ^{ns}	0.7111 ^{ns}
TR x CaSiO ₃	0.6678 ^{ns}	0.3993 ^{ns}	0.9580 ^{ns}	0.1335 ^{ns}	0.2094 ^{ns}
TR (days)	Tukey's test (SE, n = 8)				
TR1	2.23 ± 0.19 a	41.25 ± 1.68 a	31.38 ± 1.72 a	1113.37 ± 34.39 a	26.25 ± 1.83 ab
TR2	2.20 ± 0.34 a	38.28 ± 2.30 a	25.88 ± 1.41 ab	961.30 ± 45.81 ab	27.50 ± 2.50 a
TR3	1.85 ± 0.15 a	34.88 ± 1.10 ab	28.88 ± 1.79 ab	904.41 ± 39.14 b	20.00 ± 0.00 bc
TR4	1.56 ± 0.19 a	29.09 ± 2.44 b	22.63 ± 1.24 b	597.28 ± 50.82 c	15.00 ± 1.89 c
Fertilizing	Student's t-test (SE, n = 16)				
Absence CaSiO ₃	1.83 ± 0.15 A	35.66 ± 2.15 A	25.75 ± 1.30 A	891.24 ± 66.69 A	22.50 ± 2.14 A
Presence CaSiO ₃	2.10 ± 0.18 A	36.08 ± 1.29 A	28.63 ± 1.32 A	896.94 ± 44.39 A	21.88 ± 1.36 A
TR1: daily irrigation; TR2: irrigation every two days; TR3: irrigation every four days; and TR4: irrigation every six days. Absence of CaSiO ₃ and presence of CaSiO ₃ . FV = Source of variation. * and ^{ns} = significant ($p \leq 0.05$) and not significant ($p > 0.05$), respectively, according to the F-test. Means followed by the same letter in the column do not differ from each other by Tukey's test and Student's t-test ($p > 0.05$). ***Means ± standard deviation. TR: irrigation frequency; CaSiO ₃ : calcium silicate					

PH, NL, LA, and RV were reduced by 29.48%, 27.88%, 46.35%, and 42.86%, respectively, in treatment TR4 compared with TR1, regardless of CaSiO₃ fertilization (Table 4). There was no significant difference between treatments TR1 and TR2 for the variables PH, NL, and RV; however, no significant difference was observed for LA.

3.2 Leaf gas exchange

There was a significant interaction between irrigation frequencies and fertilization with CaSiO₃ for transpiration (E), stomatal conductance (g_s), and CO₂ assimilation rate (A_N) ($p \leq 0.05$) (Table 5). E , g_s , and A_N in treatment TR3, with the presence of CaSiO₃, increased by 81.05%, 66.48%, and 38.84%, respectively, compared to seedlings without CaSiO₃ (Table 5). The highest E , g_s , and A_N in seedlings with CaSiO₃ occurred in TR3, with increases of 142.25%, 122.31%, and 68.44%, respectively, compared to the lowest performance observed in TR1 (Table 5).

Table 5

F-test, Tukey test, and Student's t-test were performed for the variables: transpiration (E), stomatal conductance (g_s), and CO_2 assimilation rate (A_N) in soursoy seedlings fertilized with calcium silicate under different irrigation frequencies, 120 days after transplanting

F-test (p-value)						
SV	E (mmol de (H ₂ O) m ⁻² s ⁻¹)		g_s (mol de H ₂ O m ⁻² s ⁻¹)		A_N (μmol (CO ₂) m ⁻² s ⁻¹)	
Blocks	0.0671 ^{ns}		0.1045 ^{ns}		0.1145 ^{ns}	
TR	0.0259*		0.0313*		0.1013 ^{ns}	
CaSiO ₃	0.4194 ^{ns}		0.3973 ^{ns}		0.4995 ^{ns}	
TR x CaSiO ₃	0.0115*		0.0218*		0.0445*	
Tukey and Student t-test (SE, n = 4)						
TR (days)	Absence CaSiO ₃	Presence CaSiO ₃	Absence CaSiO ₃	Presence CaSiO ₃	Absence CaSiO ₃	Presence CaSiO ₃
TR1	1.13 ± 0.22 aA	0.71 ± 0.12 bA	50.75 ± 7.69 aA	33.93 ± 3.25 bA	9.28 ± 1.10 aA	6.94 ± 1.32 bA
TR2	1.02 ± 0.13 aA	1.07 ± 0.27 bA	47.97 ± 5.97 aA	50.49 ± 13.38 abA	9.14 ± 1.09 aA	9.33 ± 0.84 abA
TR3	0.95 ± 0.06 aB	1.72 ± 0.22 aA	45.31 ± 5.33 aB	75.43 ± 8.04 aA	8.42 ± 0.82 aB	11.69 ± 0.51 aA
TR4	0.83 ± 0.21 aA	0.81 ± 0.09 bA	39.30 ± 7.65 aA	40.35 ± 2.17 bA	7.65 ± 1.30 aA	8.30 ± 0.39 abA

TR1: daily irrigation; TR2: irrigation every two days; TR3: irrigation every four days; and TR4: irrigation every six days.

Absence of CaSiO_3 and presence of CaSiO_3 . SV = Source of variation. * and ^{ns} = significant ($p \leq 0.05$) and not significant ($p > 0.05$), respectively, according to the F-test. Means followed by the same letter in the column do not differ from each other by Tukey's test and Student's t-test ($p > 0.05$). ***Means ± standard deviation. TR: irrigation frequency; CaSiO_3 : calcium silicate.

Vapor pressure deficit (VPD), internal CO_2 concentration (C_i), intrinsic water use efficiency (iWUE), instantaneous water use efficiency (WUE), instantaneous carboxylation efficiency ($E_i C_i$), and leaf temperature (LT) were not significantly influenced by the interaction between irrigation frequency and CaSiO_3 fertilization, nor individually ($p > 0.05$) (Table 6).

Table 6

F-test, Tukey test, and Student's t-test were performed for the variables: vapor pressure deficit (VPD), internal CO₂ concentration (*C_i*), intrinsic water use efficiency (iWUE), instantaneous water use efficiency (WUE), instantaneous carboxylation efficiency (*E_iC_i*), and leaf temperature (*LT*) in soursop seedlings fertilized with calcium silicate under different irrigation frequencies, 120 days after transplanting

F-test (p-value)						
SV	VPD (kPa)	<i>C_i</i> (μmol CO ₂ mol air ⁻¹)	iWUE (<i>A_N</i> / <i>g_s</i>)	WUE (<i>A_N</i> / <i>E</i>)	<i>E_iC_i</i>	<i>LT</i> (°C)
Blocks	0.1011 ^{ns}	0.6439 ^{ns}	0.6640 ^{ns}	0.1491 ^{ns}	0.3176 ^{ns}	0.0243 [*]
TR	0.4833 ^{ns}	0.2639 ^{ns}	0.2341 ^{ns}	0.0510 ^{ns}	0.3903 ^{ns}	0.2197 ^{ns}
CaSiO ₃	0.7947 ^{ns}	0.8390 ^{ns}	0.8805 ^{ns}	0.7998 ^{ns}	0.1143 ^{ns}	0.9470 ^{ns}
TR x CaSiO ₃	0.6356 ^{ns}	0.4144 ^{ns}	0.3959 ^{ns}	0.2087 ^{ns}	0.4329 ^{ns}	0.4163 ^{ns}
TR (days)	Tukey's test (SE, n = 8)					
TR1	21.20 ± 0.96 a	81.14 ± 17.11 a	0.1927 ± 0.01 a	9.13 ± 0.42 a	0.2683 ± 0.14 a	29.33 ± 0.45 a
TR2	21.44 ± 0.60 a	71.99 ± 18.71 a	0.1979 ± 0.01 a	9.27 ± 0.61 a	0.2285 ± 0.07 a	29.50 ± 0.25 a
TR3	21.95 ± 0.61 a	107.04 ± 13.45 a	0.1731 ± 0.17 a	7.94 ± 0.49 a	0.1030 ± 0.01 a	29.99 ± 0.21 a
TR4	20.35 ± 0.77 a	65.18 ± 6.66 a	0.2013 ± 0.00 a	10.04 ± 0.57 a	0.1320 ± 0.02 a	29.07 ± 0.41 a
Fertilizing	Student's t-test (SE, n = 16)					
Absence CaSiO ₃	21.33 ± 0.55 A	82.91 ± 0.59 A	0.1904 ± 0.00 A	9.03 A ± 0.35	0.1202 ± 0.02 A	29.46 ± 0.24A
Presence CaSiO ₃	21.14 ± 0.52 A	79.76 ± 0.55 A	0.1920 ± 0.01 A	9.16 A ± 0.46	0.2457 ± 0.08 A	29.48 ± 0.26 A
TR1: daily irrigation; TR2: irrigation every two days; TR3: irrigation every four days; and TR4: irrigation every six days. Absence of CaSiO ₃ and presence of CaSiO ₃ . FV = Source of variation. * and ns = significant ($p \leq 0.05$) and not significant ($p > 0.05$), respectively, according to the F-test. Means followed by the same letter in the column do not differ from each other by Tukey's test and Student's t-test ($p > 0.05$). ***Means ± standard deviation. TR: irrigation frequency; CaSiO ₃ : calcium silicate						

3.3 Photosynthetic pigments

There was no significant interaction between irrigation schedules and CaSiO₃ fertilization for chlorophylls *a* and *b* and total content ($p > 0.05$) (Table 7). Chlorophyll *a*, *b*, and total content were affected only by irrigation frequency, regardless of CaSiO₃ fertilization, with reductions of 11.81%, 27.49%, and 17.18%, respectively, in treatment TR4 compared to TR1 (Table 7). The results also show that there was no significant difference in chlorophyll *a*, *b*, and total content among the soursop seedling treatments TR1, TR2, and TR3 (Table 7).

Table 7

F-test, Tukey test, and Student's t-test for the variables: chlorophyll *a*, *b*, and total in soursop seedlings fertilized with calcium silicate under different irrigation frequencies, 120 days after transplanting

F-test (p-value)			
SV	Chlorophyll <i>a</i>	Chlorophyll <i>b</i>	Chlorophyll Total
Blocks	0.1935 ^{ns}	0.2394 ^{ns}	0.2178 ^{ns}
TR	0.0017 [*]	0.0057 [*]	0.0032 [*]
CaSiO ₃	0.7160 ^{ns}	0.8577 ^{ns}	0.7939 ^{ns}
TR x CaSiO ₃	0.8836 ^{ns}	0.9159 ^{ns}	0.9051 ^{ns}
TR (days)	Tukey's test (SE, n = 8)		
TR1	30.31 ± 0.51 a	15.79 ± 0.63 a	46.10 ± 1.12 a
TR2	30.34 ± 0.73 a	15.43 ± 1.07 a	45.77 ± 1.79 a
TR3	29.21 ± 0.49 ab	14.45 ± 0.67 ab	43.66 ± 1.16 ab
TR4	26.73 ± 0.72 b	11.45 ± 0.77 b	38.18 ± 1.49 b
Fertilizing	Student's t-test (SE, n = 16)		
Absence CaSiO ₃	29.03 ± 0.59 A	14.20 ± 0.73 A	43.23 ± 1.31 A
Presence CaSiO ₃	29.26 ± 0.55 A	14.35 ± 0.68 A	43.61 ± 1.22 A
TR1: daily irrigation; TR2: irrigation every two days; TR3: irrigation every four days; and TR4: irrigation every six days. Absence of CaSiO ₃ and presence of CaSiO ₃ . FV = Source of variation. * and ns = significant ($p \leq 0.05$) and not significant ($p > 0.05$), respectively, according to the F-test. Means followed by the same letter in the column do not differ from each other by Tukey's test and Student's t-test ($p > 0.05$). ***Means ± standard deviation. TR: irrigation frequency; CaSiO ₃ : calcium silicate			

3.4 Chlorophyll *a* fluorescence

Statistical analysis revealed no significant interaction between irrigation and CaSiO₃ fertilization schedules ($p > 0.05$), as well as no significant isolated effects ($p > 0.05$) for maximum fluorescence (*F_m*), variable fluorescence (*F_v*), maximum quantum efficiency of photosystem II (PSII) (*F_v/F_m*), maximum primary efficiency of photosystem II (PSII) (*F_v/F₀*), and basal quantum efficiency of the non-photochemical process of photosystem II (PSII) (*F₀/F_m*) (Table 8). For initial fluorescence (*F₀*), there was a significant difference between irrigation frequencies ($p \leq 0.05$, Table 8).

Table 8

F-test, Tukey test, and Student's t-test for the variables: initial fluorescence (F_0), maximum fluorescence (F_m), variable fluorescence (F_v), maximum quantum efficiency of photosystem II (PSII) (F_v/F_m), maximum primary efficiency of photosystem II (PSII) (F_v/F_0), basal quantum efficiency of the non-photochemical process of photosystem II (PSII) (F_0/F_m), in soursop seedlings fertilized with calcium silicate under different irrigation frequencies, 120 days after transplanting

F-test (p-value)						
SV	F_0	F_m	F_v	F_v/F_m	F_v/F_0	F_0/F_m
Blocks	0.0020 ^{ns}	0.0235 [*]	0.0847 ^{ns}	0.4956 ^{ns}	0.6928 ^{ns}	0.7202 ^{ns}
TR	0.0062 [*]	0.1715 ^{ns}	0.4248 ^{ns}	0.2193 ^{ns}	0.5794 ^{ns}	0.5658 ^{ns}
CaSiO ₃	0.9371 ^{ns}	0.8900 ^{ns}	0.8664 ^{ns}	0.8937 ^{ns}	0.8762 ^{ns}	0.8985 ^{ns}
TR x CaSiO ₃	0.3063 ^{ns}	0.6559 ^{ns}	0.8093 ^{ns}	0.6971 ^{ns}	0.9405 ^{ns}	0.9114 ^{ns}
TR (days)	Tukey's test (SE, n = 8)					
TR1	95.75 ± 3.67 b	419.00 ± 16.77 a	323.25 ± 14.57 a	0.7708 ± 0.01 a	3.39 ± 0.13 a	0.2294 ± 0.01 a
TR2	93.87 ± 3.64 b	417.00 ± 15.88 a	323.13 ± 12.74 a	0.7741 ± 0.00 a	3.45 ± 0.08 a	0.2254 ± 0.00 a
TR3	99.88 ± 3.11 ab	423.75 ± 21.41 a	323.88 ± 19.83 a	0.7566 ± 0.01 a	3.24 ± 0.17 a	0.2380 ± 0.01 a
TR4	110.25 ± 5.25 a	461.88 ± 14.31 a	351.63 ± 9.90 a	0.7611 ± 0.01 a	3.21 ± 0.09 a	0.2381 ± 0.01 a
Fertilizing	Student's t-test (SE, n = 16)					
Absence CaSiO ₃	100.06 ± 3.55 A	429.31 ± 12.50 A	329.25 ± 9.81 A	0.7661 ± 0.00 A	3.31 ± 0.08 A	0.2332 ± 0.00 A
Presence CaSiO ₃	99.81 ± 2.77 A	431.50 ± 12.94 A	331.69 ± 11.20 A	0.7652 ± 0.00 A	3.33 ± 0.09 A	0.2323 ± 0.00 A
TR1: daily irrigation; TR2: irrigation every two days; TR3: irrigation every four days; and TR4: irrigation every six days. Absence of CaSiO ₃ and presence of CaSiO ₃ . FV = Source of variation. * and ns = significant ($p \leq 0.05$) and not significant ($p > 0.05$), respectively, according to the F-test. Means followed by the same letter in the column do not differ from each other by Tukey's test and Student's t-test ($p > 0.05$). ***Means ± standard deviation. TR: irrigation frequency; CaSiO ₃ : calcium silicate						

The F_0 in soursop seedlings increased by 15.14% in treatment TR4 compared with TR1, regardless of CaSiO₃ fertilization (Table 8). However, there were no significant differences between TR1, TR2, and TR3, indicating that, regardless of irrigation frequencies, soursop seedlings show similar performance in F_0 , except for TR4.

3.5 Biomass accumulation

There was no significant interaction between irrigation frequencies and CaSiO₃ fertilization for leaf dry mass (LDM), stem dry mass (SDM), shoot dry mass (SDM), root dry mass (RDM), total dry mass (TDM), and the root dry mass/shoot dry mass ratio (RDMR) ($p > 0.05$, Table 9).

For the LSM of soursop seedlings, the isolated factors, irrigation schedules ($p \leq 0.05$) and CaSiO₃ fertilization ($p \leq 0.05$), showed a significant effect (Table 9). For SDM, RDM, and TDM, the significant effect was observed only for irrigation schedules ($p \leq 0.05$; Table 9).

Table 9

F-test, Tukey test, and Student's t-test were performed for the variables: leaf dry mass (LDM), stem dry mass (SDM), root dry mass (RDM), total dry mass (TDM), and root dry mass/shoot dry mass ratio (RDMR) in soursop seedlings fertilized with calcium silicate under different irrigation frequencies, 120 days after transplanting

F-test (p-value)					
SV	LDM (g)	SDM (g)	RDM (g)	TDM (g)	RDMR
Blocks	0.5314 ^{ns}	0.5936 ^{ns}	0.2141 ^{ns}	0.3752 ^{ns}	0.2929 ^{ns}
TR	0.0000*	0.0109*	0.0044*	0.0001*	0.5988 ^{ns}
CaSiO ₃	0.0292*	0.4731 ^{ns}	0.8615 ^{ns}	0.2363 ^{ns}	0.1727 ^{ns}
TR x CaSiO ₃	0.3539 ^{ns}	0.9161 ^{ns}	0.4322 ^{ns}	0.8568 ^{ns}	0.5372 ^{ns}
TR (days)	Tukey's test (SE, n = 8)				
TR1	6.06 ± 0.30 a	6.71 ± 0.39 a	4.85 ± 0.27 a	17.62 ± 0.68 a	0.38 ± 0.01 a
TR2	4.58 ± 0.30 b	5.93 ± 0.56 ab	4.11 ± 0.36 ab	14.62 ± 1.12 ab	0.39 ± 0.02 a
TR3	4.51 ± 0.26 b	5.42 ± 0.39 ab	3.85 ± 0.27 ab	13.78 ± 0.81 bc	0.39 ± 0.02 a
TR4	3.07 ± 0.32 c	4.43 ± 0.20 b	3.16 ± 0.25 b	10.68 ± 0.53 c	0.42 ± 0.03 a
Fertilizing	Student's t-test (SE, n = 16)				
Absence CaSiO ₃	4.23 ± 0.34 B	5.47 ± 0.35 A	3.97 ± 0.27 A	13.67 ± 0.88 A	0.41 ± 0.41 A
Presence CaSiO ₃	4.88 ± 0.31 A	5.78 ± 0.34 A	4.02 ± 0.24 A	14.68 ± 0.80 A	0.38 ± 0.38 A
TR1: daily irrigation; TR2: irrigation every two days; TR3: irrigation every four days; and TR4: irrigation every six days. Absence of CaSiO ₃ and presence of CaSiO ₃ . FV = Source of variation. * and ns = significant ($p \leq 0.05$) and not significant ($p > 0.05$), respectively, according to the F-test. Means followed by the same letter in the column do not differ from each other by Tukey's test and Student's t-test ($p > 0.05$). ***Means ± standard deviation. TR: irrigation frequency; CaSiO ₃ : calcium silicate					

The leaf dry mass (LDM) of soursop irrigated every six days (TR4) decreased by 49.33% compared to seedlings irrigated daily (TR1), regardless of CaSiO₃ supply (Table 9). The presence of CaSiO₃ resulted in a 15.37% increase in LDM compared with seedlings without CaSiO₃, regardless of irrigation frequencies (Table 9).

The stem dry mass (SDM), root dry mass (RDM), and total dry mass (TDM) were reduced by irrigation schedules by 33.98%, 34.85%, and 39.39%, respectively, in TR4 compared to TR1, regardless of CaSiO₃ supply (Table 9). The total dry mass (TDM) showed similar behavior between TR1 and TR2 (Table 9).

3.6 Seedling quality indices and stress tolerance

There was an isolated effect of irrigation frequencies on the Dickson Quality Index (DQI) ($p \leq 0.05$, Table 10). The Dickson Quality Index (DQI) was reduced by 18.03% in treatment TR4 compared to TR1, regardless of fertilization with CaSiO₃ (Table 10). However, treatments TR1, TR2, and TR3 did not differ significantly from one another (Table 10).

Table 10

F-test, Tukey test, and Student's t-test were performed on the Dickson Quality Index (DQI) of soursop seedlings fertilized with calcium silicate under different irrigation frequencies, measured 120 days after transplanting

F-test (p-value)	
SV	DQI
Blocks	0.4574 ^{ns}
TR	0.0416 [*]
CaSiO ₃	0.0739 ^{ns}
TR x CaSiO ₃	0.5530 ^{ns}
TR (days)	Tukey's test (SE, n = 8)
TR1	4.27 ± 0.18 a
TR2	4.06 ± 0.20 ab
TR3	3.91 ± 0.16 ab
TR4	3.50 ± 0.21 b
Fertilizing	Student's t-test (SE, n = 16)
Absence CaSiO ₃	3.76 ± 0.14 A
Presence CaSiO ₃	4.11 ± 0.14 A
TR1: daily irrigation; TR2: irrigation every two days; TR3: irrigation every four days; and TR4: irrigation every six days. Absence of CaSiO ₃ and presence of CaSiO ₃ . SV = Source of variation. * and ^{ns} = Significant ($p \leq 0.05$) and not significant ($p > 0.05$), respectively, according to the F-test. Means followed by the same letter in the column do not differ from each other by Tukey's test and Student's t-test ($p > 0.05$). ***Means ± standard deviation. TR: irrigation frequency; CaSiO ₃ : calcium silicate	

The interaction ($p \leq 0.05$) between irrigation schedules and fertilization with CaSiO₃ was significant for drought tolerance efficiency (DTE) and drought sensitivity index (DSI) in soursop seedlings (Table 11). Water deficit decreased drought tolerance efficiency (DTE) in irrigation frequencies (Table 11). In treatments TR2, TR3, and TR4, the presence of CaSiO₃ resulted in increases of 4.94%, 1.28%, and 20%, respectively, compared to seedlings without CaSiO₃ (Table 11). The lowest DTE in seedlings with CaSiO₃ was observed in TR4, resulting in a 22.35% reduction compared to the highest value in treatment TR2 (Table 11).

Table 11

F-test, Tukey test, and Student's t-test were performed for the variables: drought tolerance efficiency (DTE) and drought sensitivity index (DSI) in sour sop seedlings fertilized with calcium silicate under different irrigation frequencies, 120 days after transplanting

F-test (p-value)				
SV	DTE		DSI	
Blocks	1.0000 ^{ns}		0.0000*	
TR	0.0000*		0.0000*	
CaSiO ₃	0.0000*		0.0000*	
TR x CaSiO ₃	0.0000*		0.0000*	
Tukey and Student t-test (SE, n = 4)				
TR (days)	Absence CaSiO ₃	Presence CaSiO ₃	Absence CaSiO ₃	Presence CaSiO ₃
TR1	-	-	-	-
TR2	0.81 ± 0.00 aB	0.85 ± 0.00 aA	0.19 ± 0.00 aA	0.19 ± 0.00 abA
TR3	0.78 ± 0.00 aB	0.79 ± 0.00 bA	0.22 ± 0.00 bA	0.21 ± 0.00 bB
TR4	0.55 ± 0.00 aB	0.66 ± 0.00 cA	0.45 ± 0.00 cA	0.34 ± 0.00 cB
TR1: daily irrigation; TR2: irrigation every two days; TR3: irrigation every four days; and TR4: irrigation every six days. Absence of CaSiO ₃ and presence of CaSiO ₃ . SV = Source of variation. * and ^{ns} = significant (<i>p</i> ≤ 0.05) and not significant (<i>p</i> > 0.05), respectively, according to the F-test. Means followed by the same letter in the column do not differ from each other by Tukey's test and Student's t-test (<i>p</i> > 0.05). ***Means ± standard deviation. TR: irrigation frequency; CaSiO ₃ : calcium silicate				

The drought sensitivity index (DSI) increased with water deficit, regardless of CaSiO₃ supply (Table 11). The presence of CaSiO₃ reduced the DSI by 4.54% and 24.44% in treatments TR3 and TR4, respectively, compared to seedlings without CaSiO₃ (Table 11). The highest DSI values, both in the absence and presence of CaSiO₃, occurred in TR4, with increases of 136.84% and 78.95%, respectively, compared to the lowest DSI observed in TR2 (Table 11).

3.7 Osmotic adjustment

There was no significant interaction between irrigation schedules and CaSiO₃ fertilization on total soluble sugar (TSS) content ($p > 0.05$) (Table 12). There was only an isolated effect ($p \leq 0.05$) on the irrigation frequency (Table 12). Treatment TR4 resulted in a 27.50% increase in TSS compared to TR1 (Table 12). It is observed that treatment TR2 did not differ from TR1, TR3, and TR4, indicating that seedlings can be irrigated every two days without exhibiting water stress in relation to TSS.

Table 12

F-test, Tukey's test, and Student's t-test were performed on the variable total soluble sugar (TSS) content in leaf tissues of sour sop seedlings fertilized with calcium silicate under different irrigation frequencies, measured 120 days after transplanting

F-test (p-value)	
SV	TSS (mg TSS g FM ⁻¹)
Blocks	0.0253*
TR	0.0053*
CaSiO ₃	0.1073 ^{ns}
TR x CaSiO ₃	0.0585 ^{ns}
TR (days)	Tukey's test (SE, n = 8)
TR1	28.04 ± 1.62 b
TR2	32.57 ± 2.39 ab
TR3	34.02 ± 1.27 a
TR4	35.75 ± 1.48 a
Fertilizing	Student's t-test (SE, n = 16)
Absence CaSiO ₃	31.43 ± 1.19 A
Presence CaSiO ₃	33.77 ± 1.53 A
TR1: daily irrigation; TR2: irrigation every two days; TR3: irrigation every four days; and TR4: irrigation every six days. Absence of CaSiO ₃ and presence of CaSiO ₃ . SV = Source of variation. * and ^{ns} = significant ($p \leq 0.05$) and not significant ($p > 0.05$), respectively, according to the F-test. Means followed by the same letter in the column do not differ from each other by Tukey's test and Student's t-test ($p > 0.05$). ***Means ± standard deviation. TR: irrigation frequency; CaSiO ₃ : calcium silicate	

The interaction between irrigation frequencies and fertilization with CaSiO₃ was significant for proline (PRO) ($p \leq 0.05$) (Table 13). The PRO content increased by 192.30% in seedlings without CaSiO₃ in TR3, compared to seedlings with CaSiO₃ (Table 13). In the absence of CaSiO₃, the greatest accumulation of PRO in seedlings was observed in TR3, with an increase of 329.03% compared to the lowest accumulation recorded in TR2 (Table 13). The presence of CaSiO₃ resulted in the greatest accumulation of PRO in TR3 and TR4, of 50.41% and 192.56%, respectively, compared to the lowest accumulation observed in TR1 (Table 13).

Table 13

F-test, Tukey test, and Student's t-test for the variable proline content (PRO) in the leaf tissues of soursop seedlings fertilized with calcium silicate under different irrigation frequencies, 120 days after transplanting

F-test (p-value)		
SV	PRO (mol PRO g MF ⁻¹)	
Blocks	0.2594	
TR	0.0003*	
CaSiO ₃	0.0310*	
TR x CaSiO ₃	0.0057*	
Tukey's test (SE, n = 4)		
TR (days)	Absence CaSiO ₃	Presence CaSiO ₃
TR1	3.65 ± 0.49 bA	2.42 ± 0.79 bA
TR2	2.48 ± 0.39 bA	2.63 ± 0.58 bA
TR3	10. 64 ± 2.43 aA	3.64 ± 0.85 abB
TR4	6.07 ± 0.12 bA	7.08 ± 1.36 aA
TR1: daily irrigation; TR2: irrigation every two days; TR3: irrigation every four days; and TR4: irrigation every six days. Absence of CaSiO ₃ and presence of CaSiO ₃ . SV = Source of variation. * and ^{ns} = significant (<i>p</i> ≤ 0.05) and not significant (<i>p</i> > 0.05), respectively, according to the F-test. Means followed by the same letter in the column do not differ from each other by Tukey's test and Student's t-test (<i>p</i> > 0.05). ***Means ± standard deviation. TR: irrigation frequency; CaSiO ₃ : calcium silicate		

3.8 Leaf content of K, Ca, and Si

There was no significant effect of the interaction between the factors studied, isolated or in combination, on the potassium (K) content ($p > 0.05$) (Table 14).

Table 14

F-test, Tukey test, and Student's t-test for the variable potassium (K) content in the leaf tissues of soursop seedlings fertilized with calcium silicate under different irrigation frequencies, 120 days after transplanting

F-test (p-value)	
SV	K (g kg ⁻¹)
Blocks	0.2609 ^{ns}
TR	0.0567 ^{ns}
CaSiO ₃	0.9960 ^{ns}
TR x CaSiO ₃	0.7766 ^{ns}
TR (days)	Tukey's test (SE, n = 8)
TR1	13.45 ± 0.36 a
TR2	13.22 ± 0.85 a
TR3	11.85 ± 0.35 a
TR4	11.24 ± 0.68 a
Fertilizing	Student's t-test (SE, n = 16)
Absence CaSiO ₃	12.44 ± 0.42 A
Presence CaSiO ₃	12.44 ± 0.53 A
TR1: daily irrigation; TR2: irrigation every two days; TR3: irrigation every four days; and TR4: irrigation every six days. Absence of CaSiO ₃ and presence of CaSiO ₃ . SV = Source of variation. * and ^{ns} = significant ($p \leq 0.05$) and not significant ($p > 0.05$), respectively, according to the F-test. Means followed by the same letter in the column do not differ from each other by Tukey's test and Student's t-test ($p > 0.05$). ***Means ± standard deviation. TR: irrigation frequency; CaSiO ₃ : calcium silicate	

The interaction between irrigation frequencies and fertilization with CaSiO₃ was significant for silicon (Si) and calcium (Ca) contents ($p \leq 0.05$) (Table 15).

Table 15

F-test, Tukey test, and Student's t-test were performed for the variables silicon (Si) and calcium (Ca) content in leaf tissues of soursop seedlings fertilized with calcium silicate under different irrigation frequencies, 120 days after transplanting

F-test (p-value)				
SV	Si (g kg ⁻¹)		Ca (g kg ⁻¹)	
Blocks	0.4979 ^{ns}		0.0713 ^{ns}	
TR	0.0000*		0.2647 ^{ns}	
CaSiO ₃	0.0000*		0.0006*	
TR x CaSiO ₃	0.0000*		0.0342*	
Tukey's test (SE, n = 4)				
TR (days)	Absence CaSiO ₃	Presence CaSiO ₃	Absence CaSiO ₃	Presence CaSiO ₃
TR1	8.57 ± 0.33 aA	9.01 ± 0.25 bA	1.73 ± 0.20 aA	1.69 ± 0.14 bA
TR2	9.05 ± 0.36 aA	9.76 ± 0.40 bA	1.63 ± 0.16 aA	1.86 ± 0.10 abA
TR3	8.58 ± 0.11 aB	14.73 ± 0.10 aA	1.63 ± 0.14 aB	2.32 ± 0.26 aA
TR4	8.31 ± 0.31 aA	8.73 ± 0.27 bA	1.49 ± 0.12 aB	2.27 ± 0.10 abA
TR1: daily irrigation; TR2: irrigation every two days; TR3: irrigation every four days; and TR4: irrigation every six days. Absence of CaSiO ₃ and presence of CaSiO ₃ . SV = Source of variation. * and ^{ns} = significant (<i>p</i> ≤ 0.05) and not significant (<i>p</i> > 0.05), respectively, according to the F-test. Means followed by the same letter in the column do not differ from each other by Tukey's test and Student's t-test (<i>p</i> > 0.05). ***Means ± standard deviation. TR: irrigation frequency; CaSiO ₃ : calcium silicate				

Silicon (Si) content increased by 71.68% in treatment TR3, in which seedlings were fertilized with CaSiO₃, compared to the absence of CaSiO₃ (Table 15). The highest Si content in seedlings with CaSiO₃ was observed in TR3, which corresponds to an increase of 68.72% compared to the lowest content observed in TR4 (Table 15).

There was an increase in calcium (Ca) content of 42.33% and 52.35% in treatments TR3 and TR4, respectively, in seedlings with CaSiO₃, compared to those without CaSiO₃ (Table 15). In the presence of CaSiO₃, the highest Ca content in seedlings was observed in TR3, with an increase of 37.28% compared to the lowest content recorded in TR1 (Table 15).

4. Discussion

Water stress is one of the main limiting factors in agricultural production, profoundly affecting plant growth, development, and metabolism, especially in TR4, irrigated every 6 days. The impacts are even more severe in perennial species of high economic value, such as soursop (*A. muricata* L.), which, in this study, showed reductions in growth variables, including height, number of leaves, leaf area, and root volume [44, 3, 8, 16]. The application of CaSiO₃ significantly improves the CO₂ assimilation rate, stomatal conductance, transpiration, leaf biomass, modulation of proline synthesis, and leaf silicon and calcium content.

The positive effects observed with the application of CaSiO₃ are mainly because silicon is an important element for some fruit crops, since, when deposited below the cuticle, it modulates several physiological activities, reducing excessive water loss and maintaining stable stomatal opening [12]. In this way, the species maintained active cellular metabolism, even with less water available. Maintaining cellular hydration favors physiological efficiency and plant resistance to various

stresses [62, 12, 63]. In addition, calcium contributes to membrane stability and the maintenance of cell turgor, which enhances photosynthesis and transpiration under water stress [64].

Our results demonstrate that daily irrigation (TR1), every two days (TR2), and every four days (TR3) yield similar growth performance, biomass, initial fluorescence (F_0), total soluble sugars, and seedling quality index, regardless of CaSiO_3 fertilization. This finding indicates that moderate deficit irrigation every four days (TR3) does not compromise soursop seedling development, confirming that irrigation scheduling is an important component of irrigated and sustainable agriculture, aiming to improve water use efficiency by applying the right amount of water at the right time [65].

These results represent an important strategy for the semi-arid region of Brazil, since rainfall irregularity limits agricultural production, including fruit species, and irrigation is an input that maintains the economic and social viability of the production system [66]. However, under severe stress every six days (TR4), seedling growth decreased, since soursop requires adequate water availability [5, 8] and, in very dry soils with an arid moisture regime, growth is severely restricted [4]. Under these conditions, there is a reduction in cell turgor pressure, which causes less cell division and development [67, 16], contributing to the limitation of new leaf expansion, abscission, wilting, senescence, and reduction of leaf area [13, 3, 68].

Under severe stress conditions (TR4), seedlings likely also divert metabolic energy towards osmotic acclimation and the synthesis of compatible solutes, such as total soluble sugars, which accumulate in the leaves, mainly in the cytoplasm, as mechanisms to mitigate the adverse effects of water stress [13, 69, 12]. This results in lower availability of energy resources for growth and biomass production [70], also compromising the quality of soursop seedlings [8].

Water stress also increased F_0 , indicating that, when electron flow through photosystem II (PSII) is obstructed, the plastoquinone acceptor (QA^-) cannot be fully oxidized under stress, thereby damaging PSII and reducing the species' photosynthetic efficiency [71, 51]. Furthermore, the reduction in chlorophyll content indicates that drought is directly associated with chlorophyll degradation, primarily due to excessive formation of reactive oxygen species in chloroplasts [72–74]. These alterations interfere with CO_2 assimilation rates and lead to reductions in growth, biomass, and the quality of soursop seedlings.

On the other hand, the application of CaSiO_3 proved effective in improving leaf biomass and gas exchange, increasing leaf calcium and silicon levels, and regulating proline content. The increase in leaf biomass may be related to the incorporation of silicon into plant structures, especially in leaves, which require less energy than the synthesis of organic compounds because of the high permeability of lipid bilayers [75]. Thus, the increase in silicon levels in plant tissues can partially replace carbon-based components, thereby reinforcing cell walls, conferring greater rigidity and resistance to desiccation, and increasing accumulated biomass [76]. Therefore, the increase in leaf biomass in fertilized seedlings indicates that the application of CaSiO_3 partially reduced the growth inhibition caused by drought. The research results corroborate studies that report that Triticum plants fertilized with silicon increased shoot dry mass when exposed to water stress, compared to the unfertilized control [13]. These results reinforce the role of silicon as a strategic element in mitigating the negative impacts of water limitation on plant development [12].

Regarding the application of CaSiO_3 , our results indicate that under moderate water-deficit conditions (TR3), it significantly improves transpiration (E), stomatal conductance (g_s), and CO_2 assimilation rate (A_N) in soursop seedlings. These results can be attributed to Si deposits, in the form of amorphous silica ($\text{SiO}_2 \cdot n\text{H}_2\text{O}$) in the cell wall, which form a double layer of cuticular silica, providing better osmotic adjustment, maintaining cell turgor, and increasing photosynthetic capacity [77, 78]. The accumulation of Si in the leaf epidermis can help minimize oxidative damage, protecting cell membranes and the photosynthetic apparatus [79].

The effect of Si on transpiration rates varies according to the severity of the stress. Under moderate stress conditions, Si can increase water absorption by the roots, thereby increasing transpiration rates and, consequently, plant growth under

drought stress [29]. However, our research observed continued growth up to TR3. The results corroborate recent research indicating that A_N , g_s , and E reached higher values than those of the control seedlings when treated with Si, which demonstrates the adaptation of the studied species [80, 38, 30]. Calcium also acts as an essential second messenger, amplifying signals and modulating target signaling proteins, culminating in the activation of physiological responses and the potentiation of antioxidant mechanisms, thereby increasing plants' tolerance to drought [80, 81].

The accumulation of proline in stressed plants indicates a defense mechanism to maintain intracellular osmotic pressure [19, 18], as it is an antioxidant and participates in cell signaling [35, 29]. In the present study, the significant reduction in proline content in seedlings treated with calcium silicate under moderate stress (TR3) suggests that calcium silicate may indicate relief from stress-induced damage [83]. This result may be related to foliar gas exchange and the benefits of silicon and calcium in regulating water loss through foliar transpiration, depending on the root water *status*, which would lead to greater water transport to the leaves, reflecting an increase in osmotic potential, reducing plant stress and, consequently, proline concentration [29, 84]. This osmotic adjustment is believed to be critical for protecting cellular structures and maintaining membrane and protein integrity, preventing cell damage and ensuring continuous metabolic activity [30, 85]. Similar results were observed in *Phoenix dactylifera* L. [83].; *Oryza sativa* L. [86]., and *Triticum aestivum* L. [84].

Increased transpiration in TR3 led to higher concentrations of Si and Ca in seedlings fertilized with CaSiO_3 . This result relates to the elements' primary absorption mechanism, which occurs predominantly via mass flow. In this process, Si is absorbed as silicic acid at the root level and, once inserted into the xylem vessels, is transported to the shoots and leaves through the transpiration stream [87]. Calcium, in turn, shows similar absorption, and its increase confirms that CaSiO_3 acts as a direct source of the nutrient, releasing Ca^{2+} into the soil as it solubilizes. Under water-deficit conditions, the cytosolic accumulation of Ca^{2+} plays a central role in signal transduction, activating various metabolic and molecular pathways involved in stress adaptation [88].

Under moderate stress conditions (TR3), the use of CaSiO_3 demonstrates that seedling producers, researchers, or farmers can adopt fertilization as a sustainable strategy in semi-arid regions. This practice allows for longer irrigation intervals without compromising foliar gas exchange performance and mineral nutrition, thereby increasing the drought tolerance of soursop seedlings and reducing their water sensitivity. For the future, we recommend further studies on this species, particularly focused on evaluating different CaSiO_3 dosages, to expand knowledge of its agronomic potential.

Soursop seedlings irrigated daily, every two days, and every four days showed similar results for growth, photosynthetic pigments, biomass accumulation, chlorophyll *a* fluorescence, and total soluble sugars, demonstrating that irrigation management every four days is viable in a water-deficient environment. Therefore, irrigation every six days, which indicates a severe water deficit, restricts seedling growth and development. Applying CaSiO_3 during the 4-day irrigation interval is an effective strategy to mitigate the impacts of deficit irrigation on soursop seedlings, thereby improving gas exchange, leaf biomass, proline accumulation, and silicon and calcium absorption. These effects increased the seedlings' tolerance to water deficits, optimizing water use and promoting more sustainable production.

5. Conclusions

Soursop plants subjected to water stress and silicon application showed physiological and nutritional responses that reinforced the potential of these practices as a sustainable strategy in semi-arid regions. Irrigation every two days kept the seedlings free from water stress, while intervals of up to four days were manageable under water-deficit conditions without compromising plant growth and development. On the other hand, irrigation every six days negatively affected the growth, physiology, and biochemistry of the seedlings, highlighting the limits of drought tolerance. CaSiO_3 increases tolerance to water deficit in soursop seedlings, mainly under moderate stress, by maintaining stomatal regulation and

osmotic balance, favoring physiological performance. Future perspectives indicate the need for further studies at different CaSiO₃ dosages to expand knowledge of its agronomic potential for the species.

Declarations

Acknowledgements

The authors would like to extend their sincere appreciation to the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES, grant number 001) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for awarding grants to researchers.

-Ethical Approval

Not applicable.

-Consent to Participate

Not applicable.

-Consent to Publish

Not applicable.

-Authors Contributions

Conceptualization: PECA, VM, FVSS; Methodology: PECA, VM, FVSS, JESR, AMFO; Formal analysis and investigation: PECA, VM, FVSS, FNF, JESR, AMFO, JLL, DEGS, ASF, TR, MAS, GCS, RRS, LPC, ECMV; Writing – preparing the original draft: PECA, VM, FVSS, AMFO, JESR; Writing – proofreading and editing: PECA, VM, FVSS, AMFO, JESR, AGLS; Obtaining funding: VM, FVSS; Resources: VM, FVSS, ASF; Supervision: VM, FVSS, JESR.

-Funding

There was no funding.

-Availability of data and materials

All data generated or analyzed during this study are included in this published article.

-Declaration of Interest Statement

We have no conflicts of interest to disclose.

References

1. Patil HV, Dhankani MA, Dhankani AR (2023) A review on Marvel fruit: *Annona muricata*. Med Sci Forum 21(1):26. <https://doi.org/10.3390/ECB2023-14355>
2. Sá FVS, Torres SB, Oliveira FDCD, Santos AS, Souza AAT, Pereira KTO, Peixoto TDC, Silva LA, Moreira RCL, Paiva EP, Almeida HA, Melo AS, Neto MF, Fernandes PD, Dias NS (2024) Ecophysiology of Soursop Seedlings Irrigated with Fish Farming Effluent under NPK Doses. Sustainability 16(11):4674. <https://doi.org/10.3390/su16114674>
3. Yisau JA, Abass RA, Emilimor PN, Ojekunle OO, Majomi AA (2023) Influence of watering regimes on early growth of *Annona muricata* seedlings. J Res For Wildl Environ 15(2):105-112.

4. Kome GK, Enang RK, Silatsa FBT, Yerima BPK, Ranst EV (2024) Baseline edaphic requirements of soursop (*Annona muricata* L.). Trop Plants 3(1):e022. <https://doi.org/10.48130/tp-0024-0023>
5. Leal F, Paull RE (2023) The soursop (*Annona muricata*): Botany, horticulture, and utilization. Crop Sci 63(2):362-389. <https://doi.org/10.1002/csc2.20894>
6. Zubaidi SN, Mohd Nani H, Ahmad Kamal MS, Abdul Qayyum T, Maarof S, Afzan A, Mohmad Misnan N, Hamezah HS, Baharum SN, Mediani A (2023) *Annona muricata*: Comprehensive review on the ethnomedicinal, phytochemistry, and pharmacological aspects focusing on antidiabetic properties. Life 13(2):353. <https://doi.org/10.3390/life13020353>
7. Alves SDC, Ferreira IO, Marangon, BB, Amaral Silva A, Castro JDS, Gonçalves PVL, Calijuri, ML (2025) Multicriteria analysis as a tool to support fruit culture in the Brazilian semi-arid. Environ Dev Sustain 27(6):14281-14304. <https://doi.org/10.1007/s10668-024-04521-9>
8. Freitas RS, dos Santos-Júnior JL, dos Santos Luz AF, da Silva, EC (2024) Hydration and dehydration cycles for enhanced soursop (*Annona muricata*) seedling quality and seedling drought stress resistance. New For 55(6):1843-1861. <https://doi.org/10.1007/s11056-024-10062-w>
9. Nogueira DB, da Silva AO, Giroldo AB, da Silva AP, Costa BR (2023) Dry spells in a semi-arid region of Brazil and their influence on maize productivity. J Arid Environ 209:104892. <https://doi.org/10.1016/j.jaridenv.2022.104892>
10. Bassoi LH, de Melo Chaves AR, Teixeira RP (2021) Responses of Syrah grapevine to deficit irrigation in the Brazilian semi-arid region. Agric Water Manage 258:107186. <https://doi.org/10.1016/j.agwat.2021.107186>
11. Salem EMM, Kenaway MK, Saudy HS, Mubarak M (2022) Influence of silicon forms on nutrients accumulation and grain yield of wheat under water deficit conditions. Gesunde Pflanz 74(3):539-548. <https://doi.org/10.1007/s10343-022-00629-y>
12. Rachappanavar V, Kumar M, Negi N, Chowdhury S, Kapoor M, Singh S, Rustagi S, Rai AK, Shreaz S, Negi R, Yadav AN (2024) Silicon derived benefits to combat biotic and abiotic stresses in fruit crops: Current research and future challenges. Plant Physiol Biochem 211:108680. <https://doi.org/10.1016/j.plaphy.2024.108680>
13. Seleiman MF, Al-Suhaibani N, Ali N, Akmal M, Alotaibi M, Refay Y, Dindaroglu T, Abdul-Wajid HH, Battaglia ML (2021) Drought stress impacts on plants and different approaches to alleviate its adverse effects. Plants 10(2):259. <https://doi.org/10.3390/plants10020259>
14. Santos CC, Scalón SPQ, Foresti AC, Reis LC, Dresch DM (2022) The role of silicon in the mitigation of water stress in *Eugenia myrcianthes* Nied. seedlings. Brazilian Journal of Biology 82: e260420. <https://doi.org/10.1590/1519-6984.260420>
15. Silva MS, Scalón SPQ, Santos CC, Silverio JM, Santos JKV, Dresch DM (2022) Does silicon help to alleviate water deficit stress and in the recovery of *Dipteryx alata* seedlings?. Braz J Biol 82:e259016. <https://doi.org/10.1590/1519-6984.259016>
16. Khallizadeh J, Nasrabadi RG, Etesami H (2025) Synergistic effects of silicate and *Streptomyces chartreusis* on water deficit resistance in soybean (*Glycine max* L.). Rhizosphere 34:101064. <https://doi.org/10.1016/j.rhisph.2025.101064>
17. Chen D, Wang S, Yin L, Deng X (2018) How does silicon mediate plant water uptake and loss under water deficiency?. Front Plant Sci 9:281. <https://doi.org/10.3389/fpls.2018.00281>
18. Roy TK, Islam MS, Mahiddin NA, Hossain SA, Biswas T, Antu UB, Serity SA, Miti JF, Akter S, Roy S, Biswas A, Mazrin M, Ismail, Z (2025) Application of nanoparticles (NPs) to ameliorate abiotic stress in economically important crop species: a potential review. J Crop Health 77(1):19. <https://doi.org/10.1007/s10343-024-01069-6>
19. Queiroz LLGD, Mesquita EFD, Sousa CdS, Pereira RF, Diniz JPC, Melo ASD, Alencar RSD, Dias GF, Soares VCDS, Mesquita FDO, Pires JPM, Rodrigues SS, Lins LKS, Alves ADS, Araújo KTA, Ferrz PDSC (2025) Foliar Silicon Alleviates Water Deficit in Cowpea by Enhancing Nutrient Uptake, Proline Accumulation, and Antioxidant Activity. Plants 14(8):1241. <https://doi.org/10.3390/plants14081241>

20. Zhi X, George-Jaeggli B, Han Y, Chen Q, Zhang S, Feng L, Lei Y, Li X, Wang G, Wang Z, Xiong S, Xin M, Jiao Y, Ma Y, Yang, B (2024) Comprehensive analysis on investigating water-saving potentials of irrigated cotton in semi-arid area in China. *Agric Water Manage* 301:108960. <https://doi.org/10.1016/j.agwat.2024.108960>
21. Bounajra A, El Guemmat K, Mansouri K, Akef F (2024) Towards efficient irrigation management at field scale using new technologies: A systematic literature review. *Agric Water Manage* 295:108758. <https://doi.org/10.1016/j.agwat.2024.108758>
22. Dash PK, Waller A, Baltimore T, Chowdhury M (2026) Enhancing Water Conservation in Hydroponically Grown Strawberries using Potassium Silicate. *Silicon* 18:411-425 <https://doi.org/10.1007/s12633-025-03534-x>
23. Li Q, Chen Y, Sun S, Zhu M, Xue J, Gao Z, Zhao J, Tang Y (2022) Research on crop irrigation schedules under deficit irrigation—A meta-analysis. *Water Resources Management* 36(12): 4799-4817. <https://doi.org/10.1007/s11269-022-03278-y>
24. Yuan M, Xu L, Dou J, Tang Y, Tan X, Xu W, Yu G, Zhuo B (2025) Optimizing irrigation regimes for foxtail millet (*Setaria italica* L.) in Northwest China: Effects of irrigation quota and frequency on yield and crop water productivity. *Agric Water Manage* 322:110022. <https://doi.org/10.1016/j.agwat.2025.110022>
25. Silva LDA, Soares LADA, Lima GSD, Roque IA, Fátima RTD, Lima AS (2023) Morphophysiology and water relations of *Spondias* rootstocks under different irrigation frequencies. *Rev Caatinga* 36(4): 865-874. <http://dx.doi.org/10.1590/1983-21252023v36n414rc>
26. Costa FH, Sousa GGD, Lima JMDP, Almeida MDS, Sousa HC, Gomes SP, Cruz Filho EMD, Azevedo BMD (2024) Frequencies of irrigation in millet crop under salt stress. *Rev Bras Eng Agríc Ambient* 28:e272197. <https://doi.org/10.1590/1807-1929/agriambi.v28n3e272197>
27. Etesami H, Jeong BR, Rizwan M (2020) The use of silicon in stressed agriculture management: action mechanisms and future prospects. In: Deshmukh R, Tripathi DK, Guerriero G (eds) *Metalloids in plants: advances and future prospects*. John Wiley & Sons Ltd., New Jersey
28. Souri Z, Khanna K, Karimi N, Ahmad P (2021) Silicon and plants: current knowledge and future prospects. *J Plant Growth Regulat* 40(3):906-925. <https://doi.org/10.1007/s00344-020-10172-7>
29. Wang M, Wang R, Mur LAJ, Ruan J, Shen Q, Guo S (2021) Functions of silicon in plant drought stress responses. *Horticul Res* 8:254. <https://doi.org/10.1038/s41438-021-00681-1>
30. Foresti AC, de Paula Quintão Scalón S, Santos CC, Reis LC, Linné JA (2024) Does silicon in *Eugenia myrcianthes* seedlings under water stress contribute in the tolerance and recovery?. *J Soil Sci Plant Nutr* 24(2):2208-2220. <https://doi.org/10.1007/s42729-024-01664-6>
31. Alam P, Balawi TH, Altalayan FH, Hatamleh AA, Ashraf M, Ahmad P (2021) Silicon attenuates the negative effects of chromium stress in tomato plants by modifying antioxidant enzyme activities, ascorbate–glutathione cycle and glyoxalase system. *Acta Physiol Plant* 43(7):110. <https://doi.org/10.1007/s11738-021-03276-4>
32. Alizadeh R, Jalali M, Valizadeh-Rad K, Etesami H (2025) Enhancing water deficit tolerance in canola (*Brassica napus* L.) through the synergistic application of nano-silicon and sulfur. *BMC Plant Biol* 25(1):486. <https://doi.org/10.1186/s12870-025-06535-5>
33. Zhao X, Lin S, Yu S, Zhang Y, Su L, Geng L, Cheng C, Jiang X (2024) Exogenous calcium enhances the physiological status and photosynthetic capacity of rose under drought stress. *Hortic Plant J* 10(3):853-865. <https://doi.org/10.1016/j.hpj.2023.01.010>
34. Malyukova LS, Koninskaya NG, Orlov YL, Samarina LS (2022) Effects of exogenous calcium on the drought response of the tea plant (*Camellia sinensis* (L.) Kuntze). *PeerJ* 10:e13997. <https://doi.org/10.7717/peerj.13997>
35. Ramírez-Builes VH, Küsters J, de Souza TR, Simmes C (2020) Calcium nutrition in coffee and its influence on growth, stress tolerance, cations uptake, and productivity. *Front Agron* 2:590892. <https://doi.org/10.3389/fagro.2020.590892>

36. Abbas M, Abdel-Lattif H, Shahba M (2021) Ameliorative effects of calcium sprays on yield and grain nutritional composition of maize (*Zea mays* L.) cultivars under drought stress. *Agriculture* 11(4):285. <https://doi.org/10.3390/agriculture11040285>
37. Yang Y, Guo Y (2018) Elucidating the molecular mechanisms mediating plant salt-stress responses. *New phytol* 217(2):523-539. <https://doi.org/10.1111/nph.14920>
38. Verma KK, Song XP, Verma CL, Chen ZL, Rajput VD, Wu KC, Liao F, Chen GL, Li YR (2021) Functional relationship between photosynthetic leaf gas exchange in response to silicon application and water stress mitigation in sugarcane. *Bio Res* 54:15. <https://doi.org/10.1186/s40659-021-00338-2>
39. Faroutine G, Arteaga-Ramírez R, Pineda-Pineda J, Vázquez-Peña MA (2023) Effect of calcium silicate and moisture content of the substrate on the growth and productivity parameters of cucumber. *Chilean J Agric Res* 83(3):334-346. <http://dx.doi.org/10.4067/S0718-58392023000300334>
40. Marques DJ, Bianchini HC, Maciel GM, de Mendonça TFN, Silva MFE (2022) Morphophysiological changes resulting from the application of silicon in corn plants under water stress. *J Plant Growth Regulat* 41(2):569-584. <https://doi.org/10.1007/s00344-021-10322-5>
41. Da Silva BD, Vieira NCDS, Paixão-Daruichi AP, Furlani Junior E (2025) Application of silicon and nitrogen in gas exchange of Arabic coffee plants as a function of water availability. *J Plant Nutr* 48(20):3684-3694. <https://doi.org/10.1080/01904167.2025.2519840>
42. Alvares CA, Stape JL, Sentelhas PC, Gonçalves JDM, Sparovek G (2013) Köppen's climate classification map for Brazil. *Meteorol Z* 22(6):711-728. <https://doi.org/10.1127/0941-2948/2013/0507>
43. Climate-Data (2024) Climate Mossoró - RN. <https://pt.climate-data.org/america-do-sul/brasil/rio-grande-do-norte/mossoro-4448/>. Accessed on January 15, 2025.
44. Fredrick C, Bamalo UM, Chima UD (2018) Effect of watering regimes on the germination and early seedling growth of *Annona muricata* Linn. *Prod Agric Technol* 14(1):66-72.
45. Souza TMAD, Mendonca V, Sá FVDS, Silva MJD, Dourado CST (2020) Calcium silicate as salt stress attenuator in seedlings of yellow passion fruit cv. BRS GA1. *Rev Caatinga* 33(2):509-517. <https://doi.org/10.1590/1983-21252020v33n223rc>
46. Pinto ACDQ, Silva EMD (1995) A cultura da graviola. Embrapa-SPI, Brasília
47. São José AR, Pires MDM, Freitas ALGED, Ribeiro DP, Perez, LAA (2014) Atualidades e perspectivas das Anonáceas no mundo. *Rev Bras de Frutic* 36:86-93. <https://doi.org/10.1590/S0100-29452014000500010>
48. Souza CCD, Oliveira FAD, Silva IDFD, Amorim Neto MDS (2000) Avaliação de métodos de determinação de água disponível e manejo da irrigação em terra roxa sob cultivo de algodoeiro herbáceo. *Rev Bras Eng Agríc Ambient* 4(3):338-342. <https://doi.org/10.1590/S1415-43662000000300006>
49. Novais RF, Neves JCL, Barros NF (1991) Ensaio em ambiente controlado. In: Oliveira AJ, Garrido WE, Araujo JD, Lourenco S. Métodos de pesquisa em fertilidade do solo. Embrapa-SEA, Brasília
50. Böhm W (1979) Root parameters and their measurement. In: Böhm W (ed) *Methods of studying root systems*. Springer, Berlin
51. Fernandes CDS, Ferreira-Neto M, Dias NDS, Reges LBL, Silva LDA, Moreira RCL, da Silva AA, Paiva EPD, Fernandes PD, Sá FVDS (2022) The appropriate source of nitrogen for Italian Zucchini under salt stress conditions. *J Soil Sci Plant Nutr* 22(1):560-570. <https://doi.org/10.1007/s42729-021-00668-w>
52. Falker (2008) Falker Automação Agrícola Ltda. Manual do medidor eletrônico de teor de clorofila (ClorofiLOG/CFL 1030). Falker Automação Agrícola, Porto Alegre
53. Yemm EW, Willis AJ (1954) The estimation of carbohydrates in plants extracts by antrone. *Biochem J* 57(3):508-514. <https://doi.org/10.1042/bj0570508>

54. Bates LS, Waldren RPA, Teare ID (1973) Rapid determination of free proline for water-stress studies. *Plant Soil*, 39(1):205-207. <https://doi.org/10.1007/BF00018060>
55. Silva FC (2009) *Manual de Análises Químicas de Solos, Plantas e Fertilizantes*. Embrapa Solos, Brasília
56. Furlani PR, Gallo JR (1978) Determinação de silício em material vegetal, pelo método colorimétrico do " azul-de-molibdênio". *Bragantia* 37(1):5-11. <https://doi.org/10.1590/S0006-87051978000100018>
57. Dickson A, Leaf AL, Hosner JF (1960) Quality appraisal of white spruce and white pine seedling stock in nurseries. *For Chron* 36(1):10-13. <https://doi.org/10.5558/tfc36010-1>
58. Fischer KS, Johnson EC, Edmeades GO (1982) Breeding and selection for drought resistance in tropical maize. CIMMYT, Mexico DF
59. Ali MB, El-Sadek AN (2016) Evaluation of drought tolerance indices for wheat (*Triticum aestivum* L.) under irrigated and rainfed conditions. *Commun Biom Crop Sci* 11(1):77-89.
60. Shapiro SS, Wilk MB (1965) Analysis of variance test for normality. *Biometrika* 52(3-4):591-611. <https://doi.org/10.2307/2333709>
61. Ferreira DF (2019) SISVAR: A computer analysis system to fixed effects split plot type designs. *Braz J Biom* 37(4):529-535. <https://doi.org/10.28951/rbb.v37i4.450>
62. Jesus EGD, Fatima RTD, Guerrero AC, Araújo JLD, Brito ME (2018) Growth and gas exchanges of arugula plants under silicon fertilization and water restriction. *Rev Bras Eng Agríc Ambient* 22(2):119-124. <https://doi.org/10.1590/1807-1929/agriambi.v22n2p119-124>
63. Ali AM, Bijay-Singh (2025) Silicon: a crucial element for enhancing plant resilience in challenging environments. *J Plant Nutr* 48(3): 486-521. <https://doi.org/10.1080/01904167.2024.2406479>.
64. Gupta S, Kaur N, Kant K, Jindal P, Ali A, Naeem M (2023) Calcium: A master regulator of stress tolerance in plants. *S Afr J Bot* 163:580-594. <https://doi.org/10.1016/j.sajb.2023.10.047>
65. Marković M, Krizmanić G, Brkić A, Atilgan A, Japundžić-Palenić B, Petrović D, Barać Ž (2021) Sustainable management of water resources in supplementary irrigation management. *Appl Sci* 11(6):2451. <https://doi.org/10.3390/app11062451>
66. Cavalcante LF, Silva-Matos RRS, Souto AGDL, Lima Neto AJD, Nunes JC, Mesquita FDO (2021) Soursop production under supplementary irrigation and mulching in the semiarid region of Brazil. *Rev Bras Eng Agríc Ambient* 25(8):522-528. <https://doi.org/10.1590/1807-1929/agriambi.v25n8p522-528>
67. Eghlima G, Mohammadi M, Ranjabr ME, Nezamdoost D, Mammadov A (2024) Foliar application of nano-silicon enhances drought tolerance rate of pot marigold (*Calendula officinalis* L.) by regulation of abscisic acid signaling. *BMC Plant Biol* 24(1):1220. <https://doi.org/10.1186/s12870-024-05986-6>
68. Bonnin M, Diop K, Cavelier G, Crastes M, Groenewald R, Nguyen HT, Morillon R, Pontvianne F (2025). Physiologic, Genetic and Epigenetic Determinants of Water Deficit Tolerance in Fruit Trees. *Plants* 14(12):1769. <https://doi.org/10.3390/plants14121769>
69. Wahab A, Abdi G, Saleem MH, Ali B, Ullah S, Shah W, Mumtaz S, Yasin G, Muresan CC, Marc RA (2022) Plants' physio-biochemical and phyto-hormonal responses to alleviate the adverse effects of drought stress: A comprehensive review. *Plants* 11(13):1620. <https://doi.org/10.3390/plants11131620>
70. Nour MM, Aljabi HR, AL-Huqail AA, Horneburg B, Mohammed AE, Alotaibi MO (2024) Drought responses and adaptation in plants differing in life-form. *Front Ecol Evol* 12:1452427. <https://doi.org/10.3389/fevo.2024.1452427>
71. Zhuang J, Wang Y, Chi Y, Zhou L, Chen J, Zhou W, Song J, Zhao N, Ding J (2020) Drought stress strengthens the link between chlorophyll fluorescence parameters and photosynthetic traits. *PeerJ* 8:e10046. <http://doi.org/10.7717/peerj.10046>
72. Yang X, Lu M, Wang Y, Wang Y, Liu Z, Chen, S (2021) Response mechanism of plants to drought stress. *Horticulturae* 7(3):50. <https://doi.org/10.3390/horticulturae7030050>

73. Rady MM, Boriek SHK, Abd El-Mageed TA, Seif El-Yazal MA, Ali EF, Hassan FAS, Abdelkhalik A (2021) Exogenous gibberellic acid or dilute bee honey boosts drought stress tolerance in *Vicia faba* by rebalancing osmoprotectants, antioxidants, nutrients, and phytohormones. *Plants* 10(4):748. <https://doi.org/10.3390/plants10040748>
74. Hussain I, Ayub A, Nayab A, Ashraf MA, Ashraf MA, Hussain S, Siddiqui MH, Sabir MA, Zulfiqar U, Khan TH (2024) Exogenous application of silicon and zinc attenuates drought tolerance in *Eruca sativa* L. through increasing chlorophyll pigments, osmoprotectants, and modulating defense mechanisms. *J Plant Growth Regulat* 43(9):3221-3237. <https://doi.org/10.1007/s00344-023-11116-7>
75. Hao Q, Yang S, Song Z, Li Z, Ding F, Yu C, Hu G, Liu, H (2020) Silicon affects plant stoichiometry and accumulation of C, N, and P in grasslands. *Front Plant Sci* 11:1304. <https://doi.org/10.3389/fpls.2020.01304>
76. Liu L, Song Z, Yu C, Yu G, Ellam RM, Liu H, Singh BP, Wang H (2020) Silicon effects on biomass carbon and phytolith-occluded carbon in grasslands under high-salinity conditions. *Front Plant Sci* 11:657. <https://doi.org/10.3389/fpls.2020.00657>
77. Verma KK, Song X-P, Li D-M, Singh M, Rajput VD, Malviya MK, Minkina T, Singh RK, Singh P, Li Y-R (2020) Interactive role of silicon and plant-rhizobacteria mitigating abiotic stresses: A new approach for sustainable agriculture and climate change. *Plants* 9(9):1055. <https://doi.org/10.3390/plants9091055>
78. Pereira S, Monteiro A, Moutinho-Pereira J, Dinis LT (2024) Silicon, an emergent strategy to lighten the effects of (a) biotic stresses on crops: A review. *J Agron Crop Sci* 210(6):e12762. <https://doi.org/10.1111/jac.12762>
79. Zhu Y-X, Gong H-J, Yin J-L (2019) Role of silicon in mediating salt tolerance in plants: a review. *Plants* 8(6):147. <https://doi.org/10.3390/plants8060147>
80. Neves JM, Aquino LAD, Berger PG, Neves JC, Rocha GC, Barbosa EA (2019) Silicon and boron mitigate the effects of water deficit on sunflower. *Rev Bras Eng Agríc Ambient* 23:175-182. <https://doi.org/10.1590/1807-1929/agriambi.v23n3p175-182>
81. Li H, Huo Y, Weng X, Zhou Y, Sun Y, Zhang G, Songzhu Z, Liu L, Pei J (2022) Regulation of the growth of Mongolian pine (*Pinus sylvestris* var. *mongolica*) by calcium-water coupling in a semiarid region. *Ecol Indic* 137:108736. <https://doi.org/10.1016/j.ecolind.2022.108736>
82. Li Y, Liu Y, Jin L, Peng R (2022) Crosstalk between Ca^{2+} and other regulators assists plants in responding to abiotic stress. *Plants* 11(10):1351. <https://doi.org/10.3390/plants11101351>
83. Al-Mayahi AMW (2016) Effect of silicon (Si) application on *Phoenix dactylifera* L. growth under drought stress induced by polyethylene glycol (PEG) in vitro. *Am J Plant Sci* 7(13):1711-1728. <https://doi.org/10.4236/ajps.2016.713161>
84. Ning D, Zhang Y, Li X, Qin A, Huang C, Fu Y, Gao Y, Duan A (2023) The effects of foliar supplementation of silicon on physiological and biochemical responses of winter wheat to drought stress during different growth stages. *Plants* 12(12):2386. <https://doi.org/10.3390/plants12122386>
85. Renzetti M, Funck D, Trovato M (2024) Proline and ROS: A unified mechanism in plant development and stress response?. *Plants* 14(1):2. <https://doi.org/10.3390/plants14010002>
86. Yan G, Fan X, Peng M, Yin C, Xiao Z, Liang Y (2020) Silicon improves rice salinity resistance by alleviating ionic toxicity and osmotic constraint in an organ-specific pattern. *Front Plant Sci* 11:260. <https://doi.org/10.3389/fpls.2020.00260>
87. Andreotti C, Roupheal Y, Colla G, Basile B (2022) Rate and timing of application of biostimulant substances to enhance fruit tree tolerance toward environmental stresses and fruit quality. *Agronomy* 12(3):603. <https://doi.org/10.3390/agronomy12030603>
88. Feng D, Wang X, Gao J, Zhang C, Liu H, Liu P, Sun X (2023) Exogenous calcium: Its mechanisms and research advances involved in plant stress tolerance. *Front Plant Sci* 14:1143963. <https://doi.org/10.3389/fpls.2023.1143963>

Figures




Figure 1

Temperature and relative humidity: maximum (max.), minimum (min.), and average during the experimental period




Figure 2

Soursop seedlings are fertilized with calcium silicate under different irrigation frequencies. TR1: daily irrigation; TR2: irrigation every two days; TR3: irrigation every four days; and TR4: irrigation every six days. Absence of CaSiO_3 and presence of CaSiO_3