

Supplemental Figure
for

**SARM1 deficiency preferentially protects against autonomic over
somatic neuropathy in type 2 diabetic db/db mice by preserving
mitochondrial integrity**

Yingshu Liu^{1,2}, Keya Meyers¹, Baohan Pan¹, Zola Zhou³, Aysel Fisgin³, Daniel Tsottles¹, Michael J. Polydefkis¹, Ahmet Hoke³#, Ying Liu¹ #

The Johns Hopkins University, Department of Neurology, Baltimore, MD

1. Department of Neurology and the Cutaneous Nerve Laboratory, Johns Hopkins University School of Medicine, Baltimore, Maryland.
2. Department of Endocrinology, Central Hospital of Dalian University of Technology, Dalian, Liaoning, China.
3. Department of Neurology and the Division of Neuromuscular, Johns Hopkins University School of Medicine, Baltimore, Maryland.


Corresponding Authors:

Ying Liu, Department of Neurology and the Cutaneous Nerve Laboratory, Johns Hopkins University School of Medicine, 885 North Wolfe St., Baltimore, MD 21205;
E-mail: yliu41@jhmi.edu.

Ahmet Hoke, Department of Neurology and the Division of Neuromuscular, Johns Hopkins University School of Medicine, 885 North Wolfe St., Baltimore, MD 21205;
E-mail: ahoke@jhmi.edu.

Michael J. Polydefkis, Department of Neurology and the Cutaneous Nerve Laboratory, Johns Hopkins University School of Medicine, 885 North Wolfe St., Baltimore, MD 21205;
E-mail: mpolyde@jhmi.edu.

Supplement Figure. Genetic deletion of *Sarm1* does not affect body weight and blood glucose in db/db mice



Supplement Figure. Genetic deletion of *Sarm1* does not affect body weight and blood glucose in db/db mice

(A) demonstrates that blood glucose levels were significantly elevated in both female and male 24-week-old db/db mice compared to db+/wt and db+/Sarm1-KO mice ($p < 0.0001$ for both sexes). (B) shows a corresponding significant increase in body weight in both female and male db/db mice relative to the db+/wt controls ($p < 0.0001$ for both sexes).