

Sestrin-mediated leucine sensing by mTORC1 is essential for detecting and adapting to a low leucine diet in *Drosophila*

David Sabatini (✉ sabatini@wi.mit.edu)

Whitehead Institute for Biomedical Research <https://orcid.org/0000-0002-1446-7256>

Norbert Perrimon

Harvard Medical School, Blavatnik Institute & Howard Hughes Medical Institute <https://orcid.org/0000-0001-7542-472X>

Xin Gu

Whitehead Institute for Biomedical Research / MIT biology department

Patrick Jouandin

Harvard Medical School

Max Valenstein

Whitehead Institute

Richard Binari

Harvard Medical School

Jason Locasale

Department of Pharmacology and Cancer Biology, Duke University School of Medicine

Annamarie Allen

Department of Pharmacology and Cancer Biology, Duke University School of Medicine

Michael Reid

Department of Pharmacology and Cancer Biology, Duke University School of Medicine

Keywords:

Posted Date: November 13th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-103234/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Nature on July 20th, 2022. See the published version at <https://doi.org/10.1038/s41586-022-04960-2>.

Abstract

The authors have requested that this preprint be removed from Research Square.