

Selective Antiproliferative Activity of an Oleandrogenin-Derived Androstane in Prostate Cancer Cells

Martina Metz^{a,b}, Jana Steigerová^{a,b*}, Marie Kvasnicová^{c,d}, Zlata Rychnovská^a, Tereza Kašparová^a, Tereza Štenclová^c, Lucie Rárová^d, Karol Michalak^e, Jerzy Wicha^e, Miroslav Strnad^d

^aDepartment of Clinical and Molecular Pathology, Faculty of Medicine and Dentistry, Palacký University, Hněvotínská 3, 775 15 Olomouc, Czech Republic

^bInstitute of Molecular and Translation Medicine, Faculty of Medicine and Dentistry, Palacký University and Faculty Hospital in Olomouc, Puškinova 6, 775 20 Olomouc, Czech Republic


***Corresponding author.**

Jana Steigerova, Department of Clinical and Molecular Pathology, Faculty of Medicine and Dentistry, Palacký University and University Hospital, Hněvotínská 3, 77515 Olomouc Czech Republic, Tel.: +420 585 639 541; email: jana.steigerova@upol.cz

Supplementary Information – Uncropped Western blots

Molecular weight markers (kDa) are indicated on the left of each uncropped blot.

Fig. 3A






Fig. 4A




Fig. 5A




Fig. 6A

