

Supplemental Table 2. Zone diameters and antimicrobial resistance rates of blood culture.

Species (numbers of isolates)		Antimicrobial	Amikacin	Colistin
		Break point (susceptible/resistant, mm)	$\geq 17, \leq 14^a$	n.d.
<i>A. baumannii</i> (n = 11)	Mean ZD $\pm$ SD (mm)	16.5 $\pm$ 7.2	11 $\pm$ 2.8	
	ZD range (mm)	6 to 24	7 to 15	
	Percentage of drug-resistant isolates (%)	36.4	n.d.	
			Ampicillin	Cefepime
		Break point (mm)	$\geq 17, \leq 13$	$\geq 25, \leq 18$
<i>E. coli</i> (n = 13)	Mean ZD $\pm$ SD (mm)	7.4 $\pm$ 5.0	15.8 $\pm$ 8.3	
	ZD range (mm)	6 to 24	6 to 30	
	Percentage of drug-resistant isolates (%)	92.3	69.2	
			Cefepime	Cefotaxime
		Break point (mm)	$\geq 25, \leq 18$	$\geq 18, \leq 14$
<i>K. pneumoniae</i> (n = 85)	Mean ZD $\pm$ SD (mm)	11.2 $\pm$ 6.1	7.8 $\pm$ 5.4	
	ZD range (mm)	6 to 31	6 to 31	
	Percentage of drug-resistant isolates (%)	91.8	93	
			Cefoxitin	
		Break point (mm)	$\geq 22, \leq 21$	
<i>S. aureus</i> (n = 26)	Mean ZD $\pm$ SD (mm)	22.6 $\pm$ 5.2		
	ZD range (mm)	13 to 28		
	Percentage of drug-resistant isolates (%)	30.8		
			Cefotaxime	Ceftazidime
		Break point (mm)	$\geq 26, \leq 22$	$\geq 22, \leq 19$
<i>S. Typhi</i> (n = 25)	Mean ZD $\pm$ SD (mm)	31.7 $\pm$ 4.4	29.4 $\pm$ 3.8	
	ZD range (mm)	25 to 40	23 to 37	
	Percentage of drug-resistant isolates (%)	0	0	

MIC: minimum inhibitory concentration; n.d.: not determined; ZD: zone diameter

^aBreakpoints for microbial resistance were determined following the guidelines of Clinical and Laboratory Standards Institute for which European Society of Clinical Microbiology and Infectious Diseases was used.

^bCotrimoxazole: Trimethoprim:sulfamethoxazole in the ratio 1:5

Gentamicin	Meropenem	Tigecycline			
$\geq 15, \leq 12$	$\geq 18, \leq 14$	n.d.			
14 ± 6.3	16.9 ± 9.0	22.7 ± 2.8			
6 to 20	6 to 26	17 to 26			
45.5	36.4	n.d.			
Cefotaxime	Ceftazidime	Ciprofloxacin	Colistin	Co-trimoxazole	Meropenem
$\geq 18, \leq 14$	$\geq 21, \leq 17$	$\geq 26, \leq 21$	n.d.	n.d.	$\geq 23, \leq 19$
11.3 ± 8.8	12.9 ± 8.1	12.3 ± 10.2	11.8 ± 2.0	11.1 ± 9.7	21.9 ± 9.1
6 to 34	6 to 27	6 to 34	6 to 14	6 to 30	6 to 30
92.3	69.2	76.9	n.d.	76.9	30.8
Ceftazidime	Ciprofloxacin	Colistin	Co-trimoxazole	Meropenem	
$\geq 21, \leq 17$	$\geq 26, \leq 21$	n.d.	n.d.	$\geq 23, \leq 19$	
8.5 ± 5.2	11.7 ± 7.2	13.1 ± 1.0	10.0 ± 6.7	14.1 ± 7.6	
6 to 28	6 to 28	8 to 15	6 to 25	6 to 28	
88.5	84.1	n.d.	67.8	67.0	

Ciprofloxacin	Meropenem
$\geq 31, \leq 20$	$\geq 23, \leq 19$
7.7 ± 3.4	32.4 ± 3.4
6 to 18	27 to 40
96.0	0

oratory Standards Institute for all antimicrobial agents except for ceftazidime in *S. Typhi*,