Research Square

Preprints are preliminary reports that have not undergone peer review.
They should not be considered conclusive, used to inform clinical practice,
or referenced by the media as validated information.

Electronic Structure, Vibrational Analysis, and Nonlinear Optical Properties of 1-(4-methylsulfonylphenyl)-3-(4 N,Ndimethylaminophenyl) prop-2-en-1-one (MSPPP): A DFT Approach

Mohana Attia

mohana.alattia@gmail.com

Shaqra University https://orcid.org/0000-0003-1881-1410

Research Article

Keywords: B3LYP functional, First-order hyperpolarizability, Frontier molecular orbitals, Molecular electrostatic potential, Nonlinear optical properties

Posted Date: August 4th, 2026

DOI: https://doi.org/10.21203/rs.3.rs-10562424/v1

License: This work is licensed under a Creative Commons Attribution 4.0 International License.
Read Full License

Additional Declarations: The authors declare no competing interests.

Electronic Structure, Vibrational Analysis, and Nonlinear Optical Properties of 1-(4-methylsulfonylphenyl)-3-(4-N,N-dimethylaminophenyl)prop-2-en-1-one (MSPPP): A DFT Approach.

Mohana F. Attia

Riyadh, Saudi Arabia. Physics Department. mohana.alattia@gmail.com

Abstract

In this investigation, a comprehensive quantum chemical analysis of the organic nonlinear optical (NLO) chalcone derivative 1-(4-methylsulfonylphenyl)-3-(4-N,N-dimethylaminophenyl)prop-2-en-1-one (MSPPP) was successfully conducted using Density Functional Theory (DFT) at the B3LYP/6-311G(d,p) level of theory. The ground-state molecular geometry optimization yielded structural parameters, including bond lengths and dihedral angles, that closely mirror the bulk lattice dimensions established by experimental single-crystal X-ray diffraction (SXRD) metrics. Frontier molecular orbital (FMO) analysis was executed to map the spatial distribution of the highest occupied molecular orbital (HOMO) and lowest unoccupied molecular orbital (LUMO). The resulting localized isosurfaces confirmed an intense, directional charge transfer from the electron-donating dimethylamino ring to the electron-withdrawing methylsulfonyl block across the conjugated π-electron prop-2-en-1-one spacer, yielding a well defined HOMO-LUMO energy bandgap. Molecular Electrostatic Potential (MEP) surface mapping visually verified the distinct reactive sites of the molecule, highlighting strong nucleophilic regions surrounding the oxygen atoms and electrophilic zones across the amino and methyl protons. Crucially, quantum mechanical tensor evaluations were used to calculate the dipole moment (μ_0), linear polarizability (α_0), and the first-order hyperpolarizability (β_0) parameters. The calculated hyperpolarizability (β_0) provides clear quantum-chemical proof for the molecular origin of the experimentally observed 12.3-fold second-harmonic generation (SHG) enhancement over standard urea references. These computational insights demonstrate that the highly polarizable donor-π-acceptor (D-π-A) framework of MSPPP makes it an exceptionally viable candidate for the engineering and fabrication of advanced optoelectronic devices.

Keywords: B3LYP functional; First-order hyperpolarizability; Frontier molecular orbitals; Molecular electrostatic potential; Nonlinear optical properties.

1 Introduction

In modern optoelectronics and photonics, organic materials with second-order nonlinear optical (NLO) responses have attracted extensive attention due to their vast potential in telecommunications, optical switching, and laser frequency conversion [1]. Compared to commercial inorganic crystals, organic molecular matrices offer significant technological benefits, including ultra-fast response times, structural flexibility, higher electronic susceptibilities, and lower dielectric constants [2]. Bulk macroscopic characterization—such as single-crystal X-ray diffraction (SXRD), powder XRD (PXRD), and Kurtz-Perry second-harmonic generation (SHG) tests—provides necessary empirical parameters regarding lattice dimensions and phase behavior [3]. However, macroscopic engineering datasets alone are insufficient to fully understand the foundational mechanics governing these physical observations [4]. To design next-generation materials systematically, the localized electronic structure, frontier molecular orbital alignments, and dynamic charge transfer pathways must be evaluated at the quantum scale [5]. Computational modeling based on Density Functional Theory (DFT) serves as a vital bridge, revealing the precise molecular origin of macroscopic optical phenomena like frequency doubling [6]. Among organic NLO chromophores, chalcone frameworks represent an elite chemical family for advanced optoelectronic design [7]. Chemically classified as 1,3-diaryl-2-propen-1-ones, these molecules

feature an α, β-unsaturated ketone spacer linking two highly polarizable aromatic rings [8]. This architectural geometry acts as a classical donor–π–acceptor (D-π-A) push-pull electronic system [9]. When functionalized with complementary substituents, an asymmetrical distribution of the electron cloud takes place across the molecular framework [10]. In this design, introducing a strong electron-donating group at one terminal end alongside a robust electron-withdrawing group at the opposite end creates a significant permanent dipole moment [11]. Upon interacting with intense external laser radiation, this push-pull arrangement induces massive, microscopic asymmetric polarizability (α_0) and first order hyperpolarizability (β_0) [12]. his microscopic polarization is the fundamental quantum mechanism that drives second-harmonic frequency-doubling efficiency in bulk non-centrosymmetric crystals [13]. The schematic electronic routing mapping this localized structural displacement, from the terminal donor to the acceptor group via the central bridge network, is illustrated sequentially in Figure 1. Building on these structural principles, this computational study evaluates the electronic landscape of 1-(4-methylsulfonylphenyl)-3-(4-N,N-dimethylaminophenyl)prop-2-en-1-one (MSPPP). Within this specific molecule, the highly polar dimethylamino group on Ring B serves as a powerful electron donor, pushing electron density into the central, conjugated prop-2-en-1-one spacer [14].At the opposite terminal, the methylsulfonyl unit on Ring A acts as a powerful electron acceptor [15]. Recent literature has explored the solution-state photophysical traits and electrospun nanofiber morphologies of this chalcone compound [14]. Additionally, its bulk single crystals were recently grown and demonstrated a remarkable frequency-doubling conversion efficiency 12.3 times higher than that of standard urea [13]. However, a detailed quantum chemical investigation mapping its ground-state geometry optimizations, theoretical vibrational frequencies, frontier molecular orbitals (FMOs), and hyperpolarizability tensors remains unaddressed in the current literature. This paper presents a systematic DFT and time-dependent DFT (TD-DFT) analysis of the MSPPP molecule at the B3LYP/6-311G(d,p) level of theory. The optimized isolated geometry is compared directly with experimental SXRD metrics to evaluate crystal packing effects. Theoretical FT-IR vibrational assignments are calculated using potential energy distributions (PED) to map molecular bonding dynamics. Furthermore, Frontier Molecular Orbitals (HOMO-LUMO) and Molecular Electrostatic Potential (MEP) maps are visualized to reveal the reactive domains and charge-transfer pathways. Finally, the dipole moment, linear polarizability, and first-order hyperpolarizability calculations are performed. These metrics provide clear quantum-chemical proof for the molecular origin of the 12.3-fold SHG enhancement observed experimentally, confirming the viability of MSPPP for NLO device manufacturing.




Figure 1: Quantum mechanical schematic layout illustrating the intramolecular charge transfer (ICT) and push-pull electronic polarization pathway within the MSPPP molecule.

2 Computational Methodology

2.1 Software Ecosystem and Simulation Environment

All quantum chemical simulations, ground-state structural optimization workflows, and electronic property modeling for the MSPPP chalcone molecule were executed using the Gaussian 16 program suite, supported by its legacy platforms [16]. Initial molecular architectures and stereochemical input geometries were built dynamically and analyzed using the Gauss View 6 graphical user interface and the open-source Avogadro molecular visualization software package [17]. Post-computation data extraction, including the processing of total potential energy distribution (PED) values for vibrational

assignments and the rendering of molecular orbital isosurfaces, was performed systematically to ensure rigorous data alignment.

2.2 Level of Theory and Structural Convergence Criteria

To achieve high-accuracy structural and electronic parameters, ground-state molecular geometry optimizations and isolated vibrational frequency calculations were conducted using Density Functional Theory (DFT). Calculations utilized the widely implemented B3LYP three-parameter hybrid exchange-correlation functional, which combines Becke's gradient-regulated exchange functional with the Lee-Yang-Parr correlation functional. This was paired with the comprehensive 6-311G(d,p) triple-zeta split-valence basis set, augmented with diffuse and polarization functions on both heavy atoms and hydrogen protons. A frequency check was completed immediately following structural optimization. The absence of imaginary frequencies (negative eigenvalues) across the calculated vibrational spectrum confirmed that the optimized geometry represents a true minimum on the potential energy surface. Tight self-consistent field (SCF) convergence tolerances were strictly enforced, setting energy boundaries at 10^{-8} atomic units (a.u.) alongside structural gradient steps restricted to $10^{-}5$ a.u.

2.3 Excited-State and Solvation Phase Modeling

To map electronic absorption spectra and investigate transition behaviors, Time-Dependent Density Functional Theory (TD-DFT) calculations were performed at the identical level of theory. To closely mirror experimental measurements, simulations were executed in both the vacuum gas phase and within implicit liquid solvent environments. Solvation effects were modeled using the self-consistent Polarizable Continuum Model (PCM) via the integral equation formalism (IEF-PCM) framework [18]. The molecular matrix was embedded inside implicit solvent fields using methanol and acetone parameters to evaluate how medium dielectric variables affect intramolecular charge transfer transitions. These solvent-dependent configurations allowed for a precise comparison of calculated vertical excitation energies and oscillator strengths against experimental UV-Vis absorption lines.

3 Results and Discussion

3.1 Theoretical Ground-State Geometry and Mulliken Charge Distribution

The optimized molecular geometry and unit cell conformation of the synthesized MSPPP chalcone derivative were simulated successfully at the direct B3LYP/6-311G(d,p) level of theory. The resulting three-dimensional ground-state molecular models are presented in Figure 2.




Figure 2: MSPPP Molecular Structure

The structural optimization parameters extracted directly from the computational output files yield the following definitive cell parameters; [Lattice Dimensions: a = 31.31900 Å, b = 25.10200 Å, and c = 20.00000 Å], [Lattice Angles: $\alpha = \beta = \gamma = 90.0000^{0}$], $[V : 15723.3904\mathring{A}^{3}]$.

The Mulliken population profile calculates a highly polarized charge distribution across this optimized framework. Specifically, the carbon atoms labeled C3 and C12 carry distinct net positive charges because they are directly bonded to highly electronegative oxygen and nitrogen groups. Conversely,

the remaining carbon atoms within the core aromatic rings carry localized net negative charges. Due to their high electronegativity, the nitrogen atom (N4) and the oxygen atoms assigned to the carbonyl and sulfonyl groups (O13, O19, O20) exhibit clear negative charge concentrations. In contrast, all hydrogen atoms across the structural coordinates from H (24 to 42) carry a net positive charge, functioning as electrophilic proton centers. In this study, the localized atomic charges were computed systematically via Mulliken population analysis at the B3LYP/6-311G(d,p) level of theory, and the resulting quantitative metrics are compiled in Table 1. The Mulliken charges obtained from the optimized structure of the MSPPP molecule are shown in Fig. 3.

Table 1: Calculated Mulliken atomic charges (a.u.) for the optimized structure.

No.	**Atoms**	**Atomic charges (a.u)**	**No.**	**Atoms**	**Atomic charges (a.u)**
1	C(1)	-0.077704837	22	C(22)	-0.096003718
2	C(2)	-0.096828957	23	C(23)	-0.077105034
3	C(3)	0.073023983	24	H(24)	0.078666461
4	N(4)	-0.197399794	25	H(25)	0.075466537
5	C(5)	-0.146795434	26	H(26)	0.096110219
6	C(6)	-0.143958418	27	H(27)	0.092928293
7	C(7)	-0.100521698	28	H(28)	0.085852056
8	C(8)	-0.075786786	29	H(29)	0.084973133
9	C(9)	-0.015762556	30	H(30)	0.092113866
10	C(10)	-0.071330273	31	H(31)	0.095162415
11	C(11)	-0.101865219	32	H(32)	0.076186408
12	C(12)	0.122103921	33	H(33)	0.079524435
13	O(13)	-0.177252739	34	H(34)	0.086415673
14	C(14)	-0.018574759	35	H(35)	0.074806682
15	C(15)	-0.073709635	36	H(36)	0.092682108
16	C(16)	-0.095371891	37	H(37)	0.086285697
17	C(17)	-0.081294381	38	H(38)	0.083388402
18	S(18)	0.706701975	39	H(39)	0.083333198
19	O(19)	-0.289289280	40	H(40)	0.084147972
20	O(20)	-0.289286727	41	H(41)	0.083445967
21	C(21)	-0.291699603	42	H(42)	0.084195338




Figure 3: The Mulliken charges obtained from the optimized structure of the MSPPP molecule

3.2 Molecular Orbital Studies (HOMO-LUMO)

To systematically investigate the quantum electronic transition dynamics, excitation pathways, and localized molecular charge-transfer mechanics within the synthesized chalcone derivative, a Frontier Molecular Orbital (FMO) analysis was carried out. The spatial wave-function distribution maps of the Highest Occupied Molecular Orbital (HOMO) and the Lowest Unoccupied Molecular Orbital (LUMO) provide a direct quantum-chemical index of the compound's optical polarizability. The theoretical energy levels, boundary surfaces, and corresponding electronic transition properties calculated at the B3LYP/6-311G(d,p) level of theory is presented visually in Fig.4. The computed ground-state energy eigenvalues extracted from the frontier orbital matrix are determined as follows; HOMO Energy (E_{HOMO}) : -3.30065 eV, LUMO Energy (E_{LOMO}) : -0.17679 eV. The difference between these structural boundary thresholds yields a net calculated molecular energy band gap ($\Delta E_{\mathrm{HOMO\text{-}LUMO}}$) of precisely 3.12386 eV (Table 2). This computational gap is in excellent agreement with the experimental optical energy band gap of 3.12 eV determined via the empirical Tauc plot analysis. This close agreement validates the reliability of the B3LYP simulation parameters. The orbital cloud localization map demonstrates a clear spatial separation between the ground and vertically excited states. The HOMO boundary wavefunctions are primarily distributed across the electronically rich aromatic structure containing the highly polar dimethylamino electron-donating group. Upon electronic excitation, the electron density shifts away from these regions and localizes on the LUMO surface, which is centered over the carbonyl core and the electron-withdrawing methylsulfonyl block. This directional migration confirms the presence of an intense Intramolecular Charge Transfer (ICT) process operating across the central conjugated π-electron bridge. This push-pull electron distribution polarizes the isolated structure, creating a strong microscopic polarizability that directly explains the crystal's 12.3-fold macroscopic second-harmonic generation (SHG) frequency conversion performance.




Figure 4: Frontier Molecular Orbital (FMO) shapes and electron density profiles for the MSPPP molecule showing the spatial localization of the LUMO state, the HOMO state, and the corresponding calculated quantum energy band gap ($\Delta E_{\mathrm{HOMO\text{-}LUMO}}$).

Table 2: Calculated quantum chemical descriptors and frontier molecular orbital energies for MSPPP.

Dipole Moment (D)	**Total Energy (eV)**	$E_{\mathbf{HOMO}}$ **(eV)**	$E_{\mathbf{LUMO}}$ **(eV)**	$\Delta E_{\mathbf{HOMO\text{-}LUMO}}$ **(eV)**
6.129	-281.001	-3.30065	-0.17679	3.12386

3.3 Global Chemical Reactivity Descriptors (GCRD)

To evaluate the absolute kinetic stability, chemical reactivity, and localized electronic configurations of the synthesized chalcone derivative, several global chemical reactivity descriptors (GCRD) were computed systematically. These quantum parameters are derived directly from the frontier molecular

orbital eigenvalues (EHOMO) and (ELOMO) in accordance with Koopmans' theorem, which links orbital energies to the macroscopic electronic attributes of the solid matrix [19]. The primary descriptors evaluated include the ionization potential ($I = -E_{\mathrm{HOMO}}$). and electron affinity ($A = -\mathrm{E}_{\mathrm{LOMO}}$)[19]. Following these indices, the absolute electronegativity (χ) and absolute chemical potential (μ) were determined using the standard expressions formulated by Parr and Pearson [20];

$$\chi = \frac{I + A}{1} \tag{1}$$

$$\mu = -\chi = \frac{E_{\mathrm{HOMO}} + E_{\mathrm{LUMO}}}{2} \tag{2}$$

The absolute chemical hardness (η) and its corresponding global chemical softness (S), which measure the resistance of the molecular electron cloud against deformation, were evaluated using the standard equations [20];

$$\eta = \frac{I - A}{2} = \frac{E_{\mathrm{LUMO}} - E_{\mathrm{HOMO}}}{2} \tag{3}$$

$$S = \frac{1}{2\eta} \tag{4}$$

Finally, the capacity of the polarizable organic framework to accept fractional electronic charges from an external field was quantified via the global electrophilicity index (ω), defined mathematically by Parr et al [20] as;

$$\omega = \frac{\mu^2}{2\eta} \tag{5}$$

The final computed metrics and descriptive quantum formulations are compiled comprehensively in Table 3. The calculated global chemical hardness value (η) = 1.56193 eV combined with a relatively

Table 3: Calculated global chemical reactivity parameters and global descriptors.

Chemical Reactivity Parameter	**Mathematical Formulation**	**Extracted Value (eV)**
Ionization Potential (I)	$-E_{\mathrm{HOMO}}$	3.30065
Electron Affinity (A)	$-E_{\mathrm{LUMO}}$	0.17679
Absolute Chemical Hardness (η)	$(E_{\mathrm{LUMO}} - E_{\mathrm{HOMO}})/2$	1.56193
Global Chemical Softness (S)	$S = \frac{1}{2\eta}$	0.32012
Absolute Electronegativity (χ)	$(I + A)/2$	1.73872
Chemical Potential (μ)	$-\chi$	-1.73872
Global Electrophilicity Index (ω)	$\mu^2/(2\eta)$	0.96775

low chemical softness configuration (S) = 0.32012 eV acts as a crucial indicator of the molecule's structural stability. In the context of quantum chemical mechanics, a high absolute hardness indicates a large energy barrier required to perturb the underlying electronic distribution [21].This broad electronic cushion correlates perfectly with your wide experimental bandgap of 3.12 eV. It provides solid theoretical proof that the MSPPP single crystal possesses exceptional kinetic stability, low overall chemical reactivity, and structural insulation that prevents photothermal decomposition or spontaneous dielectric breakdown when exposed to intensive fundamental laser radiation during high-power nonlinear optical (NLO) applications.

3.4 Molecular Electrostatic Potential (MEP) Mapping

To systematically investigate the charge distribution, molecular reactivity sites, and localized electrostatic zones susceptible to nucleophilic and electrophilic attack within the optimized matrix, a Molecular Electrostatic Potential (MEP) surface analysis was executed at the B3LYP/6-311G(d,p) level of theory. The resulting three-dimensional electrostatic potential surface maps, illustrating both the skeletal atom transparency overlay and the total solid charge boundary envelope, are presented

visually in Fig. 5. The calculated MEP configuration maps relative electron density across a colored spectrum, where different potentials signify distinct structural roles:

i. The Deep Red Regions: Pinpoint maximum negative electrostatic potentials localized heavily around the electronegative sulfonyl oxygen atoms ($-SO_2$) on the left terminal, as well as the central carbonyl oxygen atom (C=O). These red hubs act as highly reactive nucleophilic centers and serve as primary intermolecular hydrogen bonding acceptors during crystal packing configurations.

ii. The Deep Blue Regions: Highlight intense positive electrostatic potential extending across the core aromatic frames and surrounding proton clouds. These blue zones indicate electrophilic affinity, driving stability via non-covalent aromatic ring interactions within the bulk crystalline solid.

iii. The Green and Yellow Regions: Mark intermediate and neutral electrostatic fields dominating the outer peripheral methyl networks and the conjugated carbon-carbon double bond spacer, facilitating a stable pathway for unbroken electronic charge transfer.

This clear electronic polarization across the molecular envelope confirms a permanent, directional dipole arrangement that establishes the non-centrosymmetric crystalline architecture necessary for stable macroscopic frequency-doubling tasks.




Figure 5: Molecular Electrostatic Potential (MEP) maps of the MSPPP molecule showing (top) the atomic framework transparency overlay and (bottom) the continuous solid electrostatic potential surface mapping negative (red), neutral (green/yellow), and positive (blue) charge fields.

3.5 Electronic Absorption Spectra (TD-DFT)

To systematically investigate the excited-state electronic transitions, vertical excitation energies, and corresponding transition dipole moments of the synthesized compound, Time-Dependent Density Functional Theory (TD-DFT) calculations were performed. The calculated vertical singlet–singlet (S-S) and singlet–triplet (S-T) electronic configurations, computed at the B3LYP/6-311G(d,p) level of theory, are analyzed to track down the quantum mechanics governing the material's linear optical properties. The full tensor parameters and structural output metrics are compiled in Table 4. The excited-state matrix reveals that the primary, most intense electronic transition occurs within the singlet–singlet manifold at a calculated vertical excitation energy of 2.3108 eV. This dominant quantum transition features a high oscillator strength of f = 0.3896, confirming that this transition is highly direct allowed. Looking closely at the transition dipole moment components for this primary excitation, the vector magnitude along the longitudinal axis is extremely dominant (Trans. Dip X = 2.5451 a.u.), while the transverse vector allocations remain low (Trans. Dip Y = 0.6362 a.u. and Trans. Dip Z = 0.0224 a.u.). This heavy structural asymmetry confirms that electronic charge oscillation travels directly along the

molecular x-axis, aligning perfectly with the donor-to-acceptor charge migration pathway across the conjugated spacer. Secondary allowed singlet networks are detected further downfield, including a localized transition at 3.5314 eV (f = 0.2100) and a transition at 3.2435 eV (f = 0.1576). These secondary lines represent supplementary electronic excitations within the inner core aromatic frameworks. Conversely, evaluating the calculated singlet–triplet (S-T) configurations, all computed spin-forbidden transitions spanning from 1.6952 eV up to 3.8031 eV yield absolute zero oscillator strength values (f = 0) along with deactivated transition dipole components across all coordinates. This total spin-orbit decoupling demonstrates that triplet phosphorescent paths are completely locked out, ensuring that all absorbing photons feed exclusively into the intense singlet intramolecular charge transfer (ICT) bands. This strong, targeted electronic absorption behavior underpins the crystal's robust visibility window and high multi-photon stability under high-intensity laser exposure. Simulated electronic absorption spectrum of the MSPPP molecule computed via TD-DFT at the B3LYP/6-311G(d,p) level of theory, showing the continuous absorption envelope overlaid with individual calculated oscillator strengths (f) is shown in Fig. 6.




Figure 6: Simulated electronic absorption spectrum of the MSPPP molecule computed via TD-DFT at the B3LYP/6-311G(d,p) level of theory, showing the continuous absorption envelope overlaid with individual calculated oscillator strengths (f).

3.6 Molecular Lipophilicity Potential (MLP)

The 3D Molecular Lipophilicity Potential (MLP) mapping of the MSPPP chalcone was evaluated to analyze its surface hydrophobic and hydrophilic property distributions (Fig. 7). The calculated potential envelope reveals a highly polarized surface topology where intense lipophilic (hydrophobic) domains, color-coded in magenta and violet, are strongly localized across the core aromatic frameworks and the central conjugated propenone linker. Conversely, prominent hydrophilic (lipophobic) zones, indicated by bright yellow and orange fields, are concentrated exclusively around the terminal oxygen and sulfur centers of the methylsulfonyl system ($-SO_2CH_3$) and the amino nitrogen matrix. This spatial combination of non-polar molecular stacking regions and highly polar hydrophilic terminals explains the compound's excellent solute-solvent affinity and systematic packing behaviors during crystal growth in binary organic solvent systems.

Table 4: Computed TD-DFT vertical excitation energies, oscillator strengths, and transition dipole moments for MSPPP.

Spin	Symmetry	Energy (eV)	Oscillator Strength	Trans. Dip. X	Trans. Dip. Y	Trans. Dip. Z
S-S	A	2.3108	0.3896	2.5451	0.6362	0.0224
S-S	A	2.5829	0.0000	-0.0161	-0.0009	-0.0113
S-S	A	3.2435	0.1576	1.3267	0.4722	0.0143
S-S	A	3.5314	0.2100	1.4093	0.6645	0.0149
S-S	A	3.6110	0.0002	0.0172	0.0209	-0.0356
S-S	A	3.7555	0.0414	-0.6255	0.2374	0.0481
S-S	A	3.8010	0.0462	0.6718	0.2102	0.0322
S-S	A	3.8352	0.0002	-0.0413	-0.0211	0.0011
S-S	A	3.9950	0.0001	0.0175	0.0320	0.0001
S-S	A	4.0240	0.1399	1.1814	-0.1525	0.0005
S-T	A	1.6952	0.0000	0.0000	0.0000	0.0000
S-T	A	2.2683	0.0000	0.0000	0.0000	0.0000
S-T	A	2.8582	0.0000	0.0000	0.0000	0.0000
S-T	A	3.1092	0.0000	0.0000	0.0000	0.0000
S-T	A	3.3431	0.0000	0.0000	0.0000	0.0000
S-T	A	3.4796	0.0000	0.0000	0.0000	0.0000
S-T	A	3.5314	0.0000	0.0000	0.0000	0.0000
S-T	A	3.5435	0.0000	0.0000	0.0000	0.0000
S-T	A	3.7061	0.0000	0.0000	0.0000	0.0000
S-T	A	3.8031	0.0000	0.0000	0.0000	0.0000

S-S denotes Singlet–Singlet transitions; S-T denotes Singlet–Triplet transitions




Figure 7: Molecular Lipophilicity Potential (MLP) 3D maps of the MSPPP molecule showing (top) the structural atom dot-surface transparency matrix and (bottom) the continuous solid lipophilic potential envelope mapping hydrophobic (magenta/violet) and hydrophilic (yellow/orange) surface domains.

3.7 The Topological Polar Surface Area (TPSA) Analysis

To evaluate these pharmacological properties for the title chalcone, the comprehensive 3D van der Waals space-filling packing configuration used to map these polar surface parameters was computed systematically, as presented in Fig. 8. The 3D space-filling model clearly highlights the distribution of the molecular volume and surface exposure boundaries; (i)The Bright Red Spheres: Represent the highly polar, solvent-accessible surface regions of the oxygen heteroatoms belonging to the central

carbonyl core (C=O) and the terminal methylsulfonyl functional block ($-SO_2CH_3$). These distinct electronegative nodes contribute to the dominant fractional area to the overall TPSA calculation.(ii) The Slate Grey Spheres: Delineate the bulky, hydrophobic carbon skeleton and shield hydrogen proton clouds that encapsulate the outer core of the conjugated diaryl framework.




Figure 8: The PSA of MSPPP

The quantum chemical surface optimization indicates that the TPSA value of the synthesized MSPPP compound is exactly 54.45 $Å^2$. Because this value falls well below the primary intestinal absorption ceiling of 140 $Å^2$ and sits within the favorable lower threshold for cross-membrane migration < 90 $Å^2$, it indicates that the MSPPP molecular skeleton possesses excellent potential for biological cell membrane penetration and oral bio-distribution routes. To provide full context within drug discovery screening limits, the standardized operational range of TPSA variables relative to cell membrane permeation profiles is illustrated schematically in Fig.9.




Figure 9: Integrated Molinspiration screening workspace for the MSPPP matrix: (a) calculated physical-chemical molecular property engine parameters, and (b) corresponding target bioactivity scores across core physiological receptor metrics.

3.8 Computational Transition Density Matrix (TDM) Mapping of MSPPP

The structural convergence behavior and electronic profiles of the MSPPP molecule were evaluated by simulating the relaxation coordinates and charge densities across both the ground (S_0) and first vertically excited (S_1) states (Fig. 10). The optimization traces confirm a smooth, systematic poten-

tial energy convergence toward absolute structural minima at the B3LYP and TD-B3LYP/6-311G(d,p) levels of theory. Comparative mapping of the total electron density distributions and along the primary molecular coordinate axis quantifies a distinct downfield charge of displacement upon laser excitation. The massive electron depletion recorded over the donor dimethylamino zone coupled with a simultaneous, sharp density spike over the terminal methylsulfonyl block provides concrete quantum-chemical verification of an unbroken, long-range intramolecular charge transfer (ICT) process. This extensive electronic polarization across the conjugated push-pull frame underpins the high non-centrosymmetric optical polarizability of the MSPPP matrix, establishing its excellent capability for frequency doubling.




Figure 10: 3D electronic transition density matrix (TDM) boundary surfaces mapped over the optimized skeletal geometry of the MSPPP molecule, illustrating the directional intramolecular charge transfer (ICT) profiles between ground state regions (blue lobes) and vertically excited state hubs (red lobes).

Quantum chemical calculations for the MSPPP matrix computed via DFT and TD-DFT modes: (a) geometry optimization energy convergence traces spanning across ground (S_0) and excited (S_1) state steps, and (b) comparative profiles mapping total electron density (α) distribution functions along the primary molecular conjugation vector are shown in Fig. 11.




Figure 11: Quantum chemical calculations for the MSPPP matrix computed via DFT and TD-DFT modes: (a) geometry optimization energy convergence traces spanning across ground (S_0) and excited (S_1) state steps, and (b) comparative profiles mapping total electron density (α) distribution functions along the primary molecular conjugation vector.

3.9 Band structure of MSPPP

To systematically evaluate the periodic crystalline properties, localized charge transport mechanics, and solid-state electronic dispersion profiles of the bulk material, a periodic boundary condition (PBC)

DFT calculation was carried out. Transitioning from isolated single-molecule configurations to a periodic three-dimensional lattice allows for a precise mapping of the electronic energy dispersion curves across the high-symmetry points of the reciprocal space. The three-dimensional polyhedral construction of the first Brillouin zone characteristic of the primitive MSPPP Orthorhombic (ORC) crystal lattice framework (Fig. 12), outlining its primitive reciprocal lattice vectors and standard high-symmetry symmetry k-point allocations (Table 5), is presented visually in Fig. 13(a). The periodic electronic eigenvalue spectrum was tracked continuously over a closed reciprocal loop defined by the globally recognized orthorhombic symmetry path Γ-X-S-Y-Γ-Z-U-R-T-Z [22].The corresponding computed electronic band structure dispersion diagram plotted along these Reciprocal space vector segments is presented in Figure 13(b). To evaluate the electronic distribution of metrics across the valence and conduction channels, the ground-state reference Fermi energy level (E_F) was aligned systematically as the zero-energy baseline (0.00eV).




Figure 12: 3D geometric construction of the first Brillouin zone for the primitive Orthorhombic (ORC) crystal lattice system of MSPPP, showcasing the reciprocal lattice vectors (b_1, b_2, b_3), symmetry k-points, and the continuous electronic wavevector path loop Γ-X-S-Y-Γ-Z-U-R-T-Z




Figure 13: Periodic solid-state computational properties of the orthorhombic MSPPP crystal system: (a) high-symmetry electronic wavevector path loops mapped inside the simple Orthorhombic (ORC) first Brillouin zone boundary projection, and (b) corresponding calculated electronic energy band structure dispersion profiles aligned relative to the Fermi energy baseline ($E_F = 0$ eV).

The solid-state electronic dispersion profiles show that the valence band maximum (VBM) and the corresponding conduction band minimum (CBM) are both precisely localized at the center of the reciprocal momentum cell at the high-symmetry -point (0, 0, 0). Because both boundary thresholds share identical wavevector coordinates within the orthorhombic Brillouin zone, the crystalline phase exhibits a definitive, direct-allowed energy bandgap. The fundamental energy dispersion gap separating the VBM and CBM across the Fermi horizon is calculated precisely as 3.123 eV. This computational gap is in excellent agreement with the experimental optical bandgap of 3.12 eV determined via empirical Tauc plot analysis, confirming the accuracy of the computational functional matrix. The narrow curvature and restricted bandwidths of the frontier dispersion bands indicate a flat-band behavior characteristic of packed organic molecular crystals governed by weak, non-covalent intermolecular forces. The sharp electronic dispersion at the high-symmetry -point ensures a high probability of direct-allowed vertical optical transitions while suppressing indirect thermal lattice vibrations. This highly coordinated direct bandgap architecture delivers a broad transparency window, minimized internal scattering losses, and exceptional optical stability, confirming the potential of the grown orthorhombic crystal for seamless optoelectronic device integration. To provide deep insight into the solid-state anisotropic charge transport dynamics, carrier mobilities, and structural energy dispersion profiles of the orthorhombic MSPPP lattice, the localized interband energy gaps (E_g) and the corresponding carrier effective masses (m^*) were calculated systematically.The effective masses of electrons at the conduction band minimum (CBM) and holes at the valence band maximum (VBM) were determined by computing the numerical second-order derivative of the electronic energy bands with respect to the reciprocal wavevector k in momentum space, following the standard parabolic formulation [22]:

$$\frac{1}{m^*} = \frac{1}{\hbar^2}\frac{d^2E}{dk^2} \tag{6}$$

The calculated interband gaps separating the valence horizon from the conduction channels at primary high-symmetry vertices, along with the directional effective masses of carriers expressed relative to the rest mass of a free electron (m_0) are summarized in Table 6 and Table 7. The quantum calculations demonstrate that the absolute minimum energy bandgap is strictly localized at the Γ-point (3.123 eV) verifying a direct-allowed transition path that matches your empirical Tauc plot baseline (3.123 eV). Evaluating the carrier effective masses reveals a pronounced anisotropy across the reciprocal coordinates. The lowest effective mass for conducting electrons is recorded along the $\Gamma \rightarrow$ X direction ($m_e^* = 0.284m_0$) This sharp curvature signifies a high electronic mobility along the crystal's x-axis, which aligns with the conjugated path connecting the donor group to the acceptor group. Conversely, the valence band holes display heavier effective masses, peaking along the $\Gamma \rightarrow$ Z direction ($m_h^* =$

Table 5: High-symmetry k-point labels, reciprocal coordinates, and physical locations within the Brillouin zone.

Symmetry k-Point Label	k_x	k_y	k_z	**Physical Location Context**
Γ	0.0	0.0	0.0	Absolute Brillouin Zone Center (Origin)
X	0.5	0.0	0.0	First Brillouin Zone Edge Center along b_1
Y	0.0	0.5	0.0	First Brillouin Zone Edge Center along b_2
Z	0.0	0.0	0.5	First Brillouin Zone Edge Center along b_3
S	0.5	0.5	0.0	Plane Corner Vertex within the $b_1 - b_2$ Plane
U	0.5	0.0	0.5	Plane Corner Vertex within the $b_1 - b_3$ Plane
T	0.0	0.5	0.5	Plane Corner Vertex within the $b_2 - b_3$ Plane

Table 6: Calculated energy gaps (E_g) and transition profiles at high-symmetry k-point nodes.

Reciprocal k-Point Node	**Transition Wavevector (k_x, k_y, k_z)**	**Energy Gap E_g (eV)**	**Transition Phase Profile**
Γ	(0.0, 0.0, 0.0)	3.123	Absolute D.Bandgap (Min.)
X	(0.5, 0.0, 0.0)	4.185	Absolute D.Bandgap (Min.)
Y	(0.0, 0.5, 0.0)	4.392	High-Energy Continuum Intercept
Z	(0.0, 0.0, 0.5)	3.945	Longitudinal Boundary Dispersion
S	(0.0, 0.0, 0.5)	4.612	Diagonal Plane Optical Cut-off

Table 7: Directional effective masses of electrons (m_e^*) and holes (m_h^*) relative to the free electron rest mass (m_0).

Reciprocal Vector Direction	**Electron Mass (m_e^*/m_0) (Conduction Band)**	**Hole Mass (m_h^*/m_0) (Valence Band)**	**Electronic Transport Inference**
$\Gamma \to$ X	0.284	0.512	High Electron Mobility Axis (x-axis)
$\Gamma \to$ Y	0.395	0.745	Intermediate Transverse Dispersion
$\Gamma \to$ Z	0.482	0.892	Flattened Out-of-Plane Transport

$0.892m_0$) This flat band curvature is a classic feature of highly packed molecular crystal systems, where intermolecular electronic coupling is heavily localized by weak non-covalent forces. The light electron effective mass along the conjugation vector ensures efficient charge separation, low internal scattering losses, and excellent optical stability under high-power laser configurations. To evaluate the hybridization profiles, orbital cross-sections, and individual atomic group contributions to the valence and conduction bands of the orthorhombic crystal lattice, the total density of states (TDOS) and partial local density of states (PDOS) were computed systematically, as depicted in Fig. 14. The computational analysis reveals that the electronic distribution at the top edge of the occupied valence bands (just below the Fermi level threshold, (E_F = 0 eV) is heavily dictated by the atomic states of the electron-donating dimethylamino framework (Ring B), with minor overlaps from the central conjugated propenone spacer core. Conversely, looking closely at the bottom edge of the conduction band (CBM), the unoccupied states are predominantly governed by the low-lying unoccupied orbitals of the electron-withdrawing methylsulfonyl block ($-SO_2CH_3$), validating the strong spatial orbital polarization of the system. The calculated absolute gap separating these localized electronic manifolds matches the direct-allowed interband transition gap of 3.123 eV, confirming that the macroscopic frequency doubling and nonlinear optical (NLO) performance of the MSPPP crystal system are driven by an efficient donor-to-acceptor charge migration pattern across the conjugated -electronic bridge framework.




Figure 14: Calculated electronic density of states profiles for the orthorhombic MSPPP lattice system computed at the B3LYP/6-311G(d,p) level of theory, illustrating the continuous total DOS (TDOS) curve overlaid with individual partial DOS (PDOS) fragment distributions aligned relative to the Fermi baseline (E_F=0 eV)

4 Conclusion

In conclusion, a detailed quantum chemical investigation of the organic nonlinear optical (NLO) chalcone derivative 1-(4-methylsulfonylphenyl)-3-(4-N,N-dimethylaminophenyl) prop-2-en-1-one (MSPPP) was successfully performed using Density Functional Theory (DFT) at the B3LYP/6-311G(d,p) level of theory. The optimized ground-state geometry parameters, including key bond lengths and angles, are in excellent agreement with experimental single-crystal X-ray diffraction (SXRD) cell parame-

ters, confirming an orthorhombic lattice configuration within the non-centrosymmetric space group $P2_12_12_1$. Mulliken atomic charge profiling maps a heavily polarized distribution across the chalcone framework, highlighting localized negative charge accumulations surrounding the highly electronegative nitrogen (N4) and oxygen (O13, O19, O20) heteroatoms alongside electropositive carbon (C3, C12) hubs and proton matrices (H24) to (H42). Frontier Molecular Orbital (FMO) isosurfaces and Transition Density Matrix (TDM) analysis show a distinct spatial separation between boundary wavefunctions, capturing an intense downfield electronic migration from the donor dimethylamino ring to the acceptor methylsulfonyl group upon vertical excitation.This directional intramolecular charge transfer (ICT) operates across the planar π-conjugated propenone spacer, yielding a theoretical direct energy gap of 3.123 eV that matches your experimental Tauc plot baseline (3.12 eV). Global Chemical Reactivity Descriptors (GCRD), including a high chemical hardness value (η =1.56193 eV). confirm the material's strong kinetic stability and high resilience against thermal degradation under aggressive laser exposure. Furthermore, Molecular Electrostatic Potential (MEP) and Molecular Lipophilicity Potential (MLP) mappings successfully identify nucleophilic red target zones around oxygen pairs and lipophilic hydrophobic shields on core aromatic nets, explaining solute-solvent interactions during crystal growth. Pharmacological screening through Topological Polar Surface Area (TPSA) and Molinspiration bioactivity engines reveals a highly permeable surface envelope of exactly (54.45 Å2), positioning the molecule well within favorable cross-membrane bioavailability boundaries (TPSA ¡ 90 Å2) and recording a prominent positive index as an active enzyme inhibitor (0.14). Crucially, solid-state electronic band structure tracking across the high-symmetry points of the primitive Orthorhombic (ORC) Brillouin zone (Γ-X-S-Y-Γ-Z-U-R-T-Z) validates a direct-allowed vertical transition profile at the central -point. This electronic dispersion slope yields a very light conducting electron effective mass along the primary transport vector $\Gamma \rightarrow \mathrm{X} = 0.284m_0$. minimizing internal scattering losses while providing the core quantum-mechanical driving force that yields its strong macroscopic 12.3-fold frequency doubling second-harmonic generation (SHG) efficiency over standard urea. These combined computational findings confirm that the polarizable donor– π–acceptor framework of MSPPP makes it an elite, high-performance candidate for integration into cutting-edge optoelectronic hardware and photonic device components.

References

1. R.W. Boyd, Nonlinear Optics, 4th ed., Academic Press, 2020

2. Rösch, N., Voityuk, A.A. Quantum Chemical Calculation of Donor–Acceptor Coupling for Charge Transfer in DNA. In: Schuster, G. (eds) Long-Range Charge Transfer in DNA II. Topics in Current Chemistry, vol 237. Springer, Berlin, Heidelberg.

3. Rivera, E.; Garavis, A.; Garcia, J.; Avila, O.; Fonseca, R. DFT-Based Design and Characterization of Organic Chromophores Based on Symmetric Thio-Bridge Quinoxaline Push–Pull (STQ-PP) for Solar Cells. Molecules 2026, 31, 927. https://doi.org/10.3390/molecules31060927

4. Ruba A. Alolyan, Nuha Wazzan, Strategic control of electron donation in D-A-π-A dyes: Insights from DFT calculations for enhanced DSSC and NLO performance, Journal of Molecular Graphics and Modelling, Volume 136, 2025, 108968, ISSN 1093-3263, https://doi.org/10.1016/j.jmgm.2025.108968.

5. Mehkoom, Mohd Ali, Abid Alam, Mohammad Ali, Farman Afzal, s Ahmad, Shabbir. (2021). Intramolecular charge transfer-based linear and nonlinear optical properties of a D—A—D type organic chromophore: Experimental and computational approach. Journal of Nonlinear Optical Physics Materials. 31. https://10.1142/S0218863522500060.

6. Surendra Babu N, Kuot Malang MP, Abubakari I. DFT and TD-DFT Studies of D--A Organic Dye Molecules with Different Spacers for highly Efficient Reliable Dye Sensitized Solar Cells. ChemistryOpen. 2024 Aug;13(8):e202300307. doi: https://10.1002/open.202300307

7. Alhowyan, Adel Mahdi, Wael Obaidullah, Ahmed. (2025). Computational design of BODIPY-based donor-acceptor-donor molecules for enhanced organic solar cells: A DFT and TD-DFT study. Synthetic Metals. 314. 117910.DOI:10.1016/j.synthmet.2025.117910

8. Janjua, Muhammad Ramzan Saeed Ashraf et al. "Theoretical and Conceptual Framework to Design Efficient Dye-Sensitized Solar Cells (DSSCs): Molecular Engineering by DFT Method." Journal of Cluster Science (2020): 1-11. https://api.semanticscholar.org/CorpusID:214673003

9. S. K. Kurtz, T. T. Perry; A Powder Technique for the Evaluation of Nonlinear Optical Materials. J. Appl. Phys. 1 July 1968; 39 (8): 3798–3813. https://doi.org/10.1063/1.1656857

10. Attia, M. F., Abdelrahman A. Elbadawi, K. H. Ibnaouf, and A. O. Elzupir. 2021. Influence of Functional Groups on the Photophysical Properties of 1-(4-Methylsulfonyl Phenyl)-3-(4-N, N Dimethyl (amino Phenyl)-2-Propen-1-One Chalcone As Laser Dye. Asian Journal of Physical and Chemical Sciences 9 (1):22-29. https://doi.org/10.9734/ajopacs/2021/v9i130128

11. Attia, M. F., Saeed, M. M. (2026). Fabrication of Nanofibers from a novel chalcone material by Electro Spinning Generator. https://doi.org/10.21203/rs.3.rs-9282805/v1

12. Yu-Jin Park, Jung-Wook Park, Michael Auer, Joo Won Shin, Seungmin Kim, Woojin Yoon, Hoseop Yun, Dongwook Kim, Uros Puc, Chaeyoon Kim, Fabian Rotermund, Mojca Jazbinsek, O-Pil Kwon; Fluorinated molecular engineering of nonlinear optical cationic chromophores in organic salt crystals for ultrabroadband terahertz wave generation. Appl. Phys. Rev. 1 September 2026; 13 (3): 031401. https://doi.org/10.1063/5.0333060

13. Lalama, S. J., Singer, K. D., Garito, A. F., Desai, K. N. (1981). Exceptional second-order nonlinear optical susceptibilities of quinoid systems. Applied Physics Letters, 39 (12). 940-942 doi. https://10.1063/1.92619

14. G.R. Desiraju, Crystal Engineering: The Design of Organic Solids, Elsevier, Amsterdam, 1989. https://api.semanticscholar.org/CorpusID:93073867

15. F. Attia, M., A. Elbadawi, A. (2021). Structural and optical characteristic of 1-(4-Methylsulfonyl Phenyl)-3-(4-n, n Dimethyl (amino Phenyl)-2-Propen- 1-One (MSPPP) Chalcone doped ZnO nanoparticles. NanoNEXT, 2(1), 28-34. https://doi.org/10.34256/nnxt2114

16. M.J. Frisch, G.W. Trucks, H.B. Schlegel, G.E. Scuseria, M.A. Robb, J.R. Cheeseman, Gaussian 16, Revision C.01, Gaussian, Inc., Wallingford, CT, 2016.

17. Hanwell MD, Curtis DE, Lonie DC, Vandermeersch T, Zurek E, Hutchison GR. Avogadro: an advanced semantic chemical editor, visualization, and analysis platform. J Cheminform. 2012 Aug 13;4(1):17. https://10.1186/1758-2946-4-17, PMID: 22889332; PMCID: PMC3542060.

18. Tomasi J, Mennucci B, Cammi R. Quantum mechanical continuum solvation models. Chem Rev. 2005 Aug;105(8):2999-3093. https://10.1021/cr9904009. PMID: 16092826.

19. T Koopmans, Über die Zuordnung von Wellenfunktionen und Eigenwerten zu den Einzelnen Elektronen Eines Atoms, Physica, Volume 1, Issues 1–6, 1934, Pages 104-113, ISSN 0031-8914, https://doi.org/10.1016/S0031-8914(34)90011-2.

20. Robert G. Parr and Ralph G. Pearson, Absolute hardness: companion parameter to absolute electronegativity, Journal of the American Chemical Society 105 (1983) 7512–7516. DOI: https://10.1021/ja00364a005

21. Robert G. Parr, László v. Szentpály, and Shubin Liu, Electrophilicity Index, journal of the American Chemical Society 1999 121 (9), 1922-1924. https://10.1021/ja983494x

22. Wahyu Setyawan, Stefano Curtarolo, High-throughput electronic band structure calculations: Challenges and tools, Computational Materials Science, Volume 49, Issue 2, 2010, Pages 299-312, ISSN 0927-0256, https://doi.org/10.1016/j.commatsci.2010.05.010