

Fecal Microbiota Transplantation as an Adjunctive Therapy for Schizophrenia: A Multicenter, Randomized, Double-Blind, Placebo-Controlled Trial Protocol

Cailing Wei

First Affiliated Hospital of Xi'an Jiaotong University

Min Jia

First Affiliated Hospital of Xi'an Jiaotong University

Yijun Li

First Affiliated Hospital of Xi'an Jiaotong University

Yanmei Ju

Shaan Probiomicros Co.,Ltd

Jiaming Liang

First Affiliated Hospital of Xi'an Jiaotong University

Xinyue Ma

Xi'an Jiaotong University

Beilin Hou

Xi'an Jiaotong University

Yudan Zhang

First Affiliated Hospital of Xi'an Jiaotong University

Ruijin Guo

BGI Research

Xueqin Song

First Affiliated Hospital of Zhengzhou University

Xiancang Ma

First Affiliated Hospital of Xi'an Jiaotong University

Feng Zhu

zhufeng1982@xjtu.edu.cn

First Affiliated Hospital of Xi'an Jiaotong University

Keywords: Fecal microbiota transplantation, Schizophrenia, Microbiota-gut-brain axis, Randomized controlled trial, Adjunctive therapy, Clinical trial protocol

Posted Date: October 23rd, 2026

DOI: <https://doi.org/10.21203/rs.3.rs-10696294/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Additional Declarations: No competing interests reported.

**Fecal Microbiota Transplantation as an Adjunctive Therapy for
Schizophrenia: A Multicenter, Randomized, Double-Blind,
Placebo-Controlled Trial Protocol**

Cailing Wei¹, Min Jia¹, Yijun Li¹, Yanmei Ju³, Jiaming Liang¹, Xinyue Ma², Beilin Hou², Yudan Zhang¹, Ruijin Guo⁴, Xueqin Song⁵, Xiancang Ma^{1*}, Feng Zhu^{1*}

¹Department of Psychiatry, The First Affiliated Hospital of Xi'an Jiaotong University, 277 Yanta West Road, Xi'an 710061, China

²College of Medicine and Forensics, Xi'an Jiaotong University, 277 Yanta West Road, Xi'an 710061, China

³Shaan Probiomicros Co.,Ltd. Xi'an 710199, China

⁴BGI Research, Shenzhen 518083, China

⁵Department of Psychiatry, The First Affiliated Hospital of Zhengzhou University, Zhengzhou 450052, China

*Corresponding authors:

Feng Zhu, The First Affiliated Hospital of Xi'an Jiaotong University, Xi'an, China.

E-mail: zhufeng1982@xjtu.edu.cn

Xiancang Ma, The First Affiliated Hospital of Xi'an Jiaotong University, Xi'an, China.

E-mail: maxiancang@163.com

Abstract

Background: Schizophrenia is a chronic, severe mental disorder with a substantial

23 global disease burden. Current antipsychotic medications demonstrate limited efficacy
24 on negative symptoms and cognitive impairment and are frequently associated with
25 significant gastrointestinal adverse effects. Emerging evidence implicates gut
26 microbiota dysbiosis in the pathophysiology of schizophrenia via the
27 microbiota-gut-brain axis, suggesting that fecal microbiota transplantation (FMT)
28 may offer a novel adjunctive therapeutic strategy. This trial aims to evaluate the
29 efficacy and safety of FMT as an adjunctive therapy in patients with
30 non-treatment-resistant schizophrenia.

31 **Methods:** This is a multicenter, randomized, double-blind, placebo-controlled trial. A
32 total of 80 patients with schizophrenia, aged 18–65 years, with a disease duration ≤ 10
33 years and a Positive and Negative Syndrome Scale (PANSS) total score ≥ 60 , will be
34 enrolled from 11 centers across China and randomized in a 1:1 ratio to either the FMT
35 group (receiving standard antipsychotic treatment in combination with FMT capsules)
36 or the placebo group (receiving standard antipsychotic treatment in combination with
37 placebo capsules). The FMT capsules are prepared from healthy donor feces, with
38 each capsule containing 1.6×10^{11} colony-forming units of active microbiota. The
39 intervention regimen consists of 10 capsules daily for 6 days, followed by 5 capsules
40 daily for 12 days, totaling 120 capsules over 18 days. The primary outcome is the
41 change in PANSS total score during the 24-week follow-up. Secondary outcomes
42 include cognitive function assessed by the MATRICS Consensus Cognitive Battery
43 (MCCB), safety assessments using the Treatment Emergent Symptom Scale (TESS),
44 Clinical Global Impression (CGI), Gastrointestinal Symptom Rating Scale (GSRS),

Bristol Stool Form Scale (BSFS), and adverse event/serious adverse event reporting, as well as laboratory parameters (complete blood count, liver and renal function, lipid profile, fasting insulin) and gut microbiota diversity. Assessments are scheduled at baseline, day7, day 18, week 4, week 8, week 12, and week 24.

Discussion: This trial will provide high-level evidence on the efficacy and safety of FMT as an adjunctive therapy for non-treatment-resistant schizophrenia and may elucidate mechanistic pathways involving the microbiota-immune-cognition axis. Regardless of the direction of results, the findings will inform the design of future studies and contribute to the evidence base for microbiome-based interventions in psychiatric disorders.

Trial Registration: *ClinicalTrials.gov* (NCT05850585). Registered on 26 May 2023.

Keywords: Fecal microbiota transplantation; Schizophrenia; Microbiota-gut-brain axis; Randomized controlled trial; Adjunctive therapy; Clinical trial protocol

Contributions to the literature

➤ FMT has been proposed as a potential adjunctive therapy for schizophrenia, but completed evidence lags behind registered trial protocols. Existing protocols focus on treatment-resistant cases; this trial targets the much larger non-treatment-resistant population, a group not yet addressed in any published RCT protocol.

- This multicenter RCT is the first to evaluate FMT specifically for non-treatment-resistant schizophrenia, using a rigorous design with comprehensive clinical and biological measurements.
- If completed, this trial will provide the definitive evidence to establish whether FMT has a role in schizophrenia treatment. Regardless of outcome, it will transform the field by moving FMT from registered protocols to actionable answers for clinicians and patients.

1. Introduction

Schizophrenia is a chronic, severe, and lifelong disabling mental disorder that typically emerges in late adolescence or early adulthood, characterized by significant abnormalities in cognition, thought, emotion, and volition. According to the Global Burden of Disease Study 2021, approximately 23.18 million people worldwide were affected by schizophrenia in 2021, with a global point prevalence of approximately 0.32%(1, 2). Schizophrenia accounted for an estimated 14.82 million disability-adjusted life years (DALYs), ranking among the top 20 causes of disability worldwide. Alarming, a systematic review and meta-analysis found that only approximately 1 in 7 (about 14%) of treated patients achieve recovery defined by improvements in both clinical and social domains, and this proportion has not increased despite major advances in treatment options over recent decades(3). Schizophrenia has thus become a globally recognized public health priority.

Current first-line antipsychotic medications, including atypical antipsychotics such

89 as olanzapine, quetiapine, risperidone, aripiprazole, amisulpride, and paliperidone,
90 demonstrate efficacy in alleviating positive symptoms but show limited improvement
91 in negative symptoms and cognitive impairment. Meta-analyses have confirmed that
92 while atypical antipsychotics offer advantages in overall symptom improvement, they
93 show limited and time-dependent efficacy on negative symptoms and cognitive
94 impairment(4). Regarding side effects, atypical antipsychotics are commonly
95 associated with anticholinergic adverse effects including dry mouth, constipation,
96 blurred vision, and urinary retention(5). Constipation, in particular, has a prevalence
97 rate of 28.1% to 36.3% among antipsychotic-treated patients(6). It not only
98 compromises medication adherence but may also progress to serious complications
99 such as paralytic ileus, bowel ischemia, intestinal perforation, and even
100 life-threatening conditions if left untreated(7). Therefore, there is an urgent need to
101 identify adjunctive therapeutic strategies that can enhance antipsychotic efficacy
102 while alleviating gastrointestinal adverse effects.

103 In recent years, the role of gut microbiota in modulating brain function and
104 behavior through the “microbiota-gut-brain” axis has garnered increasing attention(8,
105 9). Accumulating evidence implicates gut microbiota dysbiosis in the pathophysiology
106 of schizophrenia(10). Our previous research demonstrated that gut microbiota from
107 patients with schizophrenia can modulate the tryptophan-kynurenine metabolic
108 pathway in mice and induce schizophrenia-related behaviors(11). Furthermore, 16S
109 rRNA high-throughput sequencing analyses have revealed significant alterations in
110 gut microbiota composition in patients with schizophrenia(12), suggesting that gut

microbial features may serve as auxiliary biomarkers for clinical diagnosis.

Fecal microbiota transplantation involves the transfer of functional microbiota from rigorously screened healthy donors into the patient's gastrointestinal tract via specific medical techniques (e.g., oral capsules, enema, endoscopy), with the aim of reconstructing a healthy intestinal microecology. Compared to probiotics, FMT offers higher species diversity, including bacteria, viruses, fungi, and their metabolites, and can re-establish the intestinal barrier and regulate immunity through the transplantation of a complete donor-derived ecosystem(13). FMT has already been recommended for recurrent *Clostridium difficile* infection and has demonstrated positive effects in improving both gastrointestinal and behavioral symptoms in patients with autism spectrum disorder(14).

A systematic search of *ClinicalTrials.gov* (search date: July 28, 2026; filters: condition “Schizophrenia”, study type “Interventional”) yielded 3,547 registered studies, while a parallel search of the Chinese Clinical Trial Registry (ChiCTR) yielded 308 studies. Despite this substantial pool, those specifically investigating FMT in schizophrenia constitute only five discrete investigator-initiated trials (IITs) to date (Error! Unknown switch argument.).

Table 1 The currently identifiable FMT IITs in schizophrenia

Trial Identifier	Title	Population	Design	Sample Size	Status
NCT05850585	A Study of the	Non-tr	Double-bli	80	Recruiting

	Efficacy of Fecal Microbiota Transplantation (FMT) in the Treatment of Schizophrenia	eatmen t-resist ant schizo phrenia , duratio n ≤10 years, PANS S ≥60 Schizo phrenia with comor bid treatme nt-resis tant depress ion	nd, Randomiz ed Controlled Trial, oral FMT capsules-p lacebo Single Group Assignme nt, Open-labe l, oral capsules		
NCT04001439	Fecal Microbiota Transplantation for Major Depression in Schizophrenia			3088	Unknown status
ChiCTR24000 92799	Efficacy and safety of Fecal Microbiota Transplantation (FMT) by gastroscopy in the treatment of patients	Treatm ent-resi stant schizo phrenia (TRS)	Double-bli nd, three-arm: gastroscop y FMT vs oral	120	Protocol published

	with refractory		capsule		
	schizophrenia: a		FMT vs		
	randomized,		placebo		
	double-blind,				
	placebo-controlled				
	trial				
	A Randomized	Treatm	Blind,		
	Clinical Trial of	ent-resi	Randomiz		
ChiCTR21000	Fecal Microbiota	stant	ed	40	Recruiting
45243	Transplant for	schizo	Controlled		
	Treatment-Resistant	phrenia	Trial,		
	Schizophrenia	(TRS)	parallel		
	Clinical Efficacy				
	and Mechanism of	Psycho			
	Fecal Microbiota	tropic	Random		
ChiCTR26001	Transplantation in	Drug-	parallel	100	Not yet
21876	the Treatment of	Relate	control,		recruiting
	Psychotropic	d	blind		
	Drug-Related	Consti			
	Constipation	pation			

130

131 This scarcity underscores the nascent stage of clinical translation for
132 microbiome-based interventions in schizophrenia, with the majority of published
133 work remaining at the preclinical stage. Among these five IITs, only one full trial
134 protocol, the TRS-FMT protocol (ChiCTR2400092799), has been published in a

peer-reviewed journal, and it specifically targets treatment-resistant schizophrenia (TRS)(15). However, to date, no published randomized controlled trial protocol has specifically addressed the substantially larger population of non-treatment-resistant schizophrenia patients(16). The present study is therefore positioned as the only publicly available, peer-reviewed protocol targeting this broader group, offering potential advantages for earlier intervention before treatment resistance consolidates.

While the TRS-FMT protocol provides valuable insights into a difficult-to-treat subgroup, the present study distinguishes itself through its focus on non-TRS patients, a two-arm design optimized for statistical power on the primary endpoint, exclusive use of oral capsules for scalability, systematic cognitive assessment via MCCB, comprehensive metabolic monitoring, and a 24-week follow-up to assess durability (Error! Unknown switch argument.).

Table 2 Comparative Analysis: NCT05850585 vs. Published TRS-FMT Protocol
(ChiCTR2400092799)

Dimension	NCT05850585 (Present Study)	TRS-FMT Protocol (ChiCTR2400092799)
Population	Non-treatment-resistant schizophrenia	Treatment-resistant schizophrenia (TRS)
Disease duration	≤ 10 years	≤ 15 years
Rationale	Broader patient population with greater generalizability	Higher unmet need, but narrower applicability
Design	Two-arm (FMT vs placebo)	Three-arm: gastroscopy FMT vs oral capsule FMT vs double-placebo control (1:1:1 ratio)
Sample size	80 (40 per arm)	120 (40 per arm)
Route	Oral capsules only	Gastroscopy (intraduodenal) vs oral capsules
Dose	1.6×10^{11} CFU/capsule, 10 capsules/day $\times$ 6 days, then 5 capsules/day $\times$ 12 days	Not specified in available information
Standardization	Single standardized oral regimen	Exploratory comparison of two routes
Blinding	Double-blind (capsules)	Double-blind (double-placebo design: gastroscopy + capsules)
Multi-omics	Metagenomics, metatranscriptomics, metabolomics, immunomics	Microbiomics and metabolomics

Primary outcome	PANSS total score change at 24 weeks	PANSS total score change from baseline to 12 weeks
Assessment points	7 (baseline, day7, day18, week4, week8, week12, week24)	4 (baseline, week4, week8, week12)
The longest follow-up period	24 weeks	12 weeks

Note: CFU, Colony-Forming Units.

Against this background, we designed this multicenter, randomized, double-blind, placebo-controlled trial to investigate whether FMT as an adjunctive therapy can synergistically improve psychopathological symptoms, restore gut microbial homeostasis, modulate immune function, and mitigate antipsychotic-induced metabolic disturbances in patients with non-treatment-resistant schizophrenia receiving conventional antipsychotic treatment.

2. Objectives

2.1 Primary Objective

To evaluate the efficacy of FMT as an adjunctive therapy on psychopathological symptoms in patients with schizophrenia, as measured by the change in PANSS total score, and to comprehensively assess its safety profile through systematic adverse event monitoring and laboratory evaluations.

2.2 Secondary Objectives

(1) To investigate whether FMT adjunctive therapy restores gut microbial ecosystem homeostasis, as assessed by metagenomic species/strain composition, metatranscriptomic functional activity, and metabolomic profiles (SCFAs, bile acids, tryptophan metabolites);

(2) To explore whether FMT-induced microbial remodeling is associated with systemic immune rebalancing, evaluated through high-dimensional immunophenotyping (spectral flow cytometry), inflammatory proteomics, and PBMC functional assays;

(3) To evaluate whether FMT adjunctive therapy mitigates antipsychotic-induced glucose-lipid metabolic disturbances, as determined by longitudinal changes in fasting glucose, fasting insulin, Homeostatic Model Assessment of Insulin Resistance (HOMA-IR), and full lipid profiles.

3. Methods and Design

3.1 Study Design and center

This multicenter, randomized, double-blind, placebo-controlled trial aims to evaluate the efficacy and safety of FMT as an adjunctive therapy in patients with non-treatment-resistant schizophrenia. The lead center is the First Affiliated Hospital of Xi'an Jiaotong University. Participating centers include: The First Affiliated Hospital of Zhengzhou University, West China Hospital of Sichuan University, The Second Xiangya Hospital of Central South University, Suzhou Guangji Hospital, Guangzhou Brain Hospital, Ankang Hanbin District Third Hospital, Ankang Central

188 Hospital, Xunyang Disabled Persons' Federation Bo'ai Hospital, Xianyang Central
189 Hospital, Ankang Jinzhou Mental Health Rehabilitation Hospital, and Ankang Antai
190 Hospital. The overall study design, including participant allocation, intervention, and
191 assessment schedule, is summarized in **Figure 1**.

192




Figure 1. CONSORT flow diagram of the trial design.

Eligible patients with non-treatment-resistant schizophrenia (PANSS total score ≥ 60 , disease duration ≤ 10 years) will be randomized 1:1 to receive either FMT capsules or placebo capsules (identical in appearance, weight, and odor, containing lyophilized protectant only)

adjunctive to standard antipsychotic treatment, with both groups following the same administration schedule. Follow-up assessments are scheduled at baseline (V1), day 7 (V2), day 18 (V3, end of intervention), week 4 (V4), week 8 (V5), week 12 (V6), and week 24 (V7, endpoint). Assessments include clinical scales (PANSS, MCCB, TESS, CGI, GSRS, BSFS), laboratory examinations (blood counts, liver/renal function, lipid profile, fasting insulin), and biospecimen collection (blood and feces) for multi-omics profiling. The analytical framework comprises four layers: Layer 1 evaluates the clinical efficacy of FMT versus placebo (primary and secondary outcomes). Layer 2 assesses capsule fidelity by comparing the finished capsule microbiota to the original donor stool. Layer 3 quantifies donor strain engraftment, identifies its determinants, and examines its association with clinical outcomes. Layer 4 integrates multi-omics data (microbiome, metabolome, immune, and host transcriptome) to explore mechanistic pathways. Clinical scale abbreviations: FMT, fecal microbiota transplantation; PANSS, Positive and Negative Syndrome Scale; MCCB, MATRICS Consensus Cognitive Battery; TESS, Treatment Emergent Symptom Scale; CGI, Clinical Global Impression; GSRS, Gastrointestinal Symptom Rating Scale; BSFS, Bristol Stool Form Scale. Statistical and analytical abbreviations are defined in the main text abbreviations table.

3.2 Sample Size

A total of 80 participants will be enrolled, with 40 in the FMT group and 40 in the placebo group. The sample size was determined based on an estimated effect size for PANSS score reduction derived from preliminary studies, with a power of 80% and a two-sided alpha of 0.05, accounting for a 15% dropout rate.

3.3 Participants

Eligibility criteria, including inclusion, exclusion, and withdrawal criteria, are summarized in Error! Unknown switch argument..

Table 3 Eligibility Criteria

Category	Criteria
Inclusion Criteria	➤ Age 18–65 years, both sexes; ➤ DSM-5 diagnosis of schizophrenia, disease duration ≤ 10 years; ➤ PANSS total score ≥ 60 (at least 3 positive or 3 negative items ≥ 3); ➤ Junior high school education or above; ➤ Signed informed consent
Exclusion Criteria	➤ Pregnancy or lactation; ➤ Severe somatic diseases (hepatic/renal failure, cancer, active infection, inflammatory bowel disease); ➤ Refractory infections or immunocompromised status; antibiotic or immunosuppressant use within the past month; ➤ Other neuropsychiatric disorders; ➤ Fixed alcohol consumption; ➤ BMI outside 18.5–28.0
Withdrawal Criteria	➤ Severe adverse events; ➤ Symptom deterioration requiring antipsychotic class change; ➤ Voluntary withdrawal; ➤ Concurrent severe infection or use of anti-inflammatory drugs during follow-up;

➤ Poor adherence (< 80 capsules consumed)

225

226 All participating centers are tertiary or specialized psychiatric hospitals with
227 established departments of psychiatry and experience in conducting clinical trials in
228 schizophrenia. Each site has access to appropriate facilities for participant recruitment,
229 clinical assessment, laboratory sample processing, and long-term follow-up. Site
230 investigators are licensed psychiatrists with a minimum of 5 years of clinical
231 experience in schizophrenia treatment and management. All clinical assessors
232 responsible for administering PANSS, MCCB, CGI, TESS, GSRS, and BSFS are
233 required to complete centralized training and achieve an intraclass correlation
234 coefficient (ICC) ≥ 0.85 for PANSS total score and MCCB domain scores prior to
235 study initiation. FMT capsules are prepared by a single third-party manufacturer
236 (Xi'an Shanjun Biotechnology Co., Ltd.) and delivered to all sites in a standardized
237 manner. No site-specific variation in intervention preparation or administration is
238 permitted.

239 Baseline demographic and clinical characteristics will be collected for all enrolled
240 participants prior to randomization, including: age, sex, ethnicity, years of education,
241 marital status, employment status, living arrangements, socioeconomic status, height,
242 weight, body mass index (BMI), age at first onset of schizophrenia, total disease
243 duration, number of hospitalizations, family history of psychiatric disorders, comorbid
244 physical illnesses, substance use history (tobacco, alcohol), and medication history
245 (current and previous antipsychotic regimens, doses, and treatment adherence).

Baseline clinical assessments will include PANSS total and subscale scores, MCCB cognitive domain scores, CGI severity, TESS, GSRS, and BSFS. Laboratory parameters (CBC, liver and renal function tests, lipid profile, fasting glucose, fasting insulin, hs-CRP) and 12-lead ECG will also be recorded at baseline. Baseline characteristics will be summarized by treatment group using descriptive statistics (means with standard deviations for continuous variables; frequencies and percentages for categorical variables) to ensure comparability between groups.

Recruitment is anticipated to commence in November 2025 and continue for approximately 18 months, with a target enrollment of 80 participants. The final follow-up visit (Week 24) for the last enrolled participant is expected to be completed by February 2027. Recruitment periods and follow-up dates will be recorded for each participant and reported in the final study results.

3.4 Randomization, Blinding and Unblinding

The allocation sequence will be generated by an independent biostatistician not involved in participant enrollment or outcome assessment, using a block randomization method (block size of 4) stratified by center with a computer-generated algorithm (SAS PROC PLAN). Participant enrollment and group assignment will be performed by site investigators who have no access to the allocation sequence. Once a participant is deemed eligible and provides written informed consent, the site investigator will contact the central randomization center via the Interactive Web Response System (IWRS) to obtain the next sequential treatment allocation.

Participants, outcome assessors, and data analysts will remain blinded throughout

the study. FMT and placebo capsules are identical in appearance, weight, and odor, and will be uniformly prepared and coded by a third-party company (Xi'an Shanjun Biotechnology Co., Ltd.). Capsules will be administered orally by trained study nurses at each participating center; participants will be instructed to take capsules with water in the morning under observation for the first dose of each day. Adherence will be calculated as the number of capsules consumed divided by the number prescribed ($\geq 80\%$ defined as adherent; $< 80\%$ triggers withdrawal). To ensure intervention fidelity, batch-to-batch consistency will be verified by 16S rRNA sequencing and viable counting, with formal fidelity assessment detailed in Section 4.2 (capsule–stool Aitchison distance). The integrity of blinding will be assessed at the final visit using Bang's Blinding Index, and any deviations from the intended intervention will be recorded with reasons.

Consistent with ICH E6 (R2) Section 5.13.4 and CONSORT 2025 Item 20, a two-stage unblinding procedure will be implemented. The first unblinding (after database lock and blind data review) will reveal only group labels (e.g., Group A vs. Group B) without identifying active treatment versus placebo, allowing the statistician to perform the primary analysis while remaining blinded to treatment identity. The second unblinding (after the statistical analysis plan is executed and the primary analysis is completed) will fully identify the active treatment group. Emergency unblinding may be performed by the Investigator of Record (IoR) through the IWRS in the event of a medical emergency where knowledge of treatment assignment is necessary for clinical management, with documentation requirements (date, time,

reason, and procedures) recorded in the eCRF; once emergency unblinding is performed, the participant will be treated as a dropout. No interim analyses are planned unless required by the Data Monitoring Committee for safety review.

3.5 Interventions

3.5.1 Group Assignments

Participants receive standard antipsychotic treatment in combination with either FMT capsules or visually identical placebo capsules. Detailed intervention protocols are summarized in **Table 4**.

Table 4 Intervention Protocol Comparison

Item	FMT Group (n = 40)	Placebo Group (n = 40)
Antipsychotic treatment	Standard antipsychotic treatment	Standard antipsychotic treatment (same as FMT group)
Capsule content	Active microbiota (1.6×10^{11} CFU/capsule)	Lyophilized protectant only (no active microbiota)
Appearance/Weight/Odor	—	Identical to FMT capsules
Dosing regimen	10 capsules/day $\times$ 6 days, then 5 capsules/day $\times$ 12 days	Same as FMT group
Total capsules	120 capsules over 18 days	120 capsules over 18 days

3.5.2 Bowel Preparation

To enhance colonization efficiency, the following bowel preparation protocol will be performed before the first transplantation: (1) Oral administration of 2L polyethylene glycol 24 hours prior to transplantation for bowel cleansing; (2) Fasting for 4 hours before and 1 hour after transplantation; (3) Morning fasting transplantation, with food and antipsychotic medication permitted 1 hour after transplantation; (4) Antibiotics and immunosuppressants are prohibited during the intervention and follow-up periods.

3.5.3 Concomitant Medication Regulations

During the treatment period, only second-generation antipsychotics (olanzapine, quetiapine, risperidone, paliperidone, aripiprazole, amisulpride, ziprasidone) are permitted as the standard antipsychotic backbone, but clozapine is prohibited. Antipsychotic doses should be titrated to therapeutic levels within 2 weeks of enrollment. Thereafter, the drug class should not be changed, though dose adjustments are allowed. Combination of two antipsychotics is permitted. Benzodiazepines, eszopiclone, zolpidem, and medications for managing side effects (trihexyphenidyl, propranolol, metoprolol) and non-probiotic laxatives are allowed. Transcranial magnetic stimulation and biofeedback therapy are permitted. Antidepressants, mood stabilizers, medications affecting body weight or glucose-lipid metabolism, and modified electroconvulsive therapy are prohibited. Any use of prohibited medications will be recorded as a protocol deviation.

All concomitant treatments, including prescription and over-the-counter

medications, non-pharmacological therapies (e.g., psychotherapy, physical therapy), and dietary supplements, will be systematically documented at each visit (V1–V7) using a standardized case report form. For each concomitant treatment, start and end dates, dosage, frequency, and indication will be recorded. Concomitant care will be summarized by treatment group in the final analysis to assess whether any between-group differences could have confounded the primary or secondary outcomes.

3.6 Outcome Measures and Assessment Schedule

A total of 7 assessment time points are scheduled: baseline (V1), day 7 (V2), day 18 (V3), week 4 (V4), week 8 (V5), week 12 (V6), and week 24 (V7).

3.6.1 Primary Outcome

PANSS total score change: Assessed at baseline, week 4, week 8, week 12, and week 24.

3.6.2 Secondary Outcomes

Cognitive function: MATRICS Consensus Cognitive Battery (MCCB) at baseline, week 12, and week 24.

3.6.3 Safety assessments

Safety will be assessed using the Treatment Emergent Symptom Scale (TESS) for adverse treatment effects, the Clinical Global Impression (CGI) for overall illness severity and improvement, the Gastrointestinal Symptom Rating Scale (GSRS) for gastrointestinal symptoms, and the Bristol Stool Form Scale (BSFS) for stool

consistency. Adverse Event and Serious Adverse Event (AE/SAE) reporting will be conducted at all visits using Common Terminology Criteria for Adverse Events (CTCAE) criteria.

3.6.4 Adverse Event Monitoring and Management

Patients are instructed to monitor for common transient adverse events, including nausea, bloating, and mild abdominal cramping, which typically resolve within 24 hours. If symptoms persist beyond 24 hours or if severe adverse events occur (e.g., fever, severe diarrhea, signs of infection), patients are advised to contact the study site immediately. The following events should be systematically recorded in the case report form (CRF) during the first month post-FMT: infectious events (fever, bacteremia, pneumonia), psychiatric symptom exacerbation (PANSS increase $\geq 20\%$ from baseline), and severe gastrointestinal adverse events (intestinal obstruction, perforation, severe diarrhea). Patients should maintain a bowel movement diary for at least eight weeks post-FMT to monitor intestinal transit and treatment response

3.6.5 Laboratory parameters

Complete blood count, liver and renal function tests, lipid profile, fasting insulin, and 12-lead electrocardiogram at baseline, week 4, week 8, week 12, and week 24. The complete assessment schedule, including all outcome measures and biospecimen collection across the seven time points, is summarized in **Table 5**.

Table 5 Assessment Schedule and Outcome Measures

Phase	Assessment	V1 (Baseline)	V2 (Day 7)	V3 (Day 18)	V4 (Week 4)	V5 (Week 8)	V6 (Week 12)	V7 (Week 24)
Screening	Informed consent & eligibility screening	✓						
	Demographics	✓						
	Dietary and lifestyle information	✓						
Clinical Assessments	Physical examination & vital signs	✓			✓	✓	✓	✓
	12-lead ECG	✓			✓	✓	✓	✓
	Laboratory tests (CBC, liver/renal, lipid, fasting insulin)	✓			✓	✓	✓	✓
	Concomitant medications & adherence	✓	✓	✓	✓	✓	✓	✓

		V1	V2	V3	V4	V5	V6	V7
Phase	Assessment	(Baseline)	(Day 7)	(Day 18)	(Week 4)	(Week 8)	(Week 12)	(Week 24)
Primary Outcome	PANSS total score change	✓			✓	✓	✓	✓
Secondary Outcomes	MCCB cognitive function	✓					✓	✓
	CGI	✓	✓		✓	✓	✓	✓
	TESS	✓	✓		✓	✓	✓	✓
	GSRS	✓	✓		✓	✓	✓	✓
	BSFS	✓	✓		✓	✓	✓	✓
Safety Monitoring	AE / SAE reporting	✓	✓		✓	✓	✓	✓
Biospecimen Collection	Blood	✓		✓	✓	✓	✓	✓
	Feces	✓		✓	✓	✓	✓	✓

366 Note: CBC, complete blood count; ECG, electrocardiogram; PANSS, Positive and Negative

367 Syndrome Scale; MCCB, MATRICS Consensus Cognitive Battery; CGI, Clinical Global

368 Impression; TESS, Treatment Emergent Symptom Scale; GSRS, Gastrointestinal Symptom

Rating Scale; BSFS, Bristol Stool Form Scale; AE, adverse event; SAE, serious adverse event.

3.6.6 Outcome Measurement Quality Assurance

To ensure the quality and reliability of outcome measurements, the following procedures will be implemented. All clinical assessors (psychiatrists and psychologists) responsible for administering PANSS, MCCB, CGI, TESS, GSRS, and BSFS will undergo centralized training and certification prior to study initiation. The training program will include didactic instruction on scale administration, video-based case vignettes, and supervised practice with standardized patients. Inter-rater reliability will be established across all participating sites, with an intraclass correlation coefficient (ICC) ≥ 0.85 required for PANSS total score and MCCB domain scores before study commencement; site investigators failing to meet this threshold will receive additional training until the criterion is achieved. To maintain reliability throughout the trial, joint rating sessions will be conducted quarterly via videoconference, with random double-scoring of assessments (at least 10% of all visits) to monitor drift. For MCCB cognitive assessments, all test administrators will be trained to the same standardized procedures, with strict adherence to the MATRICS standardized administration manual. For adverse event reporting, site investigators will receive training on CTCAE grading criteria and SAE reporting timelines. Assessment windows will be maintained within ± 3 days of the scheduled visit dates, with deviations recorded and reported. All outcome data will be entered into the electronic case report form (eCRF) by trained research coordinators, with

source document verification and range checks for data consistency.

3.7 Biospecimen Collection and Laboratory Assays

Fecal and blood samples will be collected at multiple timepoints as specified in **Table 5**. Fecal samples will be aliquoted into three fractions for (1) metagenomics (shotgun sequencing for species/strain profiling and MAG reconstruction), (2) metatranscriptomics (RNA sequencing for active functional expression), and (3) targeted/untargeted metabolomics (UHPLC-HRMS and LC-MS/MS for SCFAs, bile acids, and tryptophan metabolites). Culturomics will be performed from raw fecal aliquots for live strain isolation.

Blood samples will be separated into plasma, serum, and PBMCs. Plasma will be analyzed by Olink/Luminex inflammatory proteomics and metabolomics; serum by clinical biochemistry and immunoglobulin profiling; PBMCs by spectral flow cytometry (≥ 40 markers), scRNA-seq (10x), bulk RNA-seq, and functional assays (PMA/ionomycin stimulation). A comprehensive summary of biospecimen types, preservation methods, analytical platforms, and targets is provided in **Table 6**. Detailed protocols for each assay are described in the **Supplementary Materials 1**.

Table 6 Biospecimen Types and Analytical Methods

Specimen	Preservation	Assay	Target
Feces (RNA)	RNA stabilization	Metatranscriptomics (Illumina NovaSeq)	Active functional expression (KEGG/MetaCyc/GO)

	buffer		
Feces (DNA)	DNA stabilization buffer	Shotgun metagenomics (Illumina NovaSeq)	Species/strain composition, functional genes, MAG reconstruction, StrainPhlAn4 tracking
Feces (raw)	No additive	Targeted + untargeted metabolomics (UHPLC-HRMS, LC-MS/MS)	SCFAs, bile acids (>20 species), tryptophan metabolites
Feces (raw)	No additive	Culturomics (MALDI-TOF / WGS)	Live bacterial isolation and identification
Plasma	-80°C	Olink/Luminex inflammatory proteomics	Inflammatory cytokines (IL-6, TNF- α , IFN- γ , etc.)
Plasma	-80°C	Metabolomics (LC-MS/MS, UHPLC-HRMS)	SCFAs, bile acids, tryptophan metabolites
Serum	-80°C	ELISA / multiplex assays	IgG/IgA/IgM subclasses, ANA, anti-dsDNA
Serum	-80°C	Automated biochemistry	Liver/renal function, glucose-lipid profile, hs-CRP
PBMC	-80°C	Spectral flow cytometry (≥ 40 markers)	T/NK/monocyte/B cell subsets, activation markers

PBMC	−80°C	scRNA-seq (10x Genomics)	Single-cell transcriptomics
PBMC	−80°C	Bulk RNA-seq	Global gene expression
PBMC	−80°C	Functional assays (PMA/ionomycin)	Intracellular cytokines (IFN- γ , TNF- α , IL-2, IL-17A)

Note: All biospecimens are stored at -80°C unless otherwise specified. Preservation methods listed refer to the conditions applied immediately after collection prior to storage. SCFAs, short-chain fatty acids; MAG, metagenome-assembled genome; WGS, whole-genome sequencing; PBMC, peripheral blood mononuclear cells; ANA, antinuclear antibodies; hs-CRP, high-sensitivity C-reactive protein.

4. Data Analysis Plan

4.1 Statistical analysis of clinical outcomes

Primary analysis: The primary endpoint is the change in PANSS total score during follow-up. Intention-to-treat (ITT) analysis with the last observation carried forward (LOCF) method will handle missing data. Analysis of covariance (ANCOVA) will compare group differences at each time point, adjusting for baseline values, BMI, age, sex, disease duration, and other covariates. The primary contrast will be the between-arm difference in PANSS total score change from baseline to week 24, reported with 95% confidence interval and Cohen's d. The intention-to-treat (ITT) set will serve as the primary analysis population, with per-protocol (PP) analysis as

sensitivity. And then the clinical outcomes will be analyzed using the same linear mixed model framework: PANSS subscales (positive, negative, general), response rate ($\geq 30\%$ PANSS reduction, modified Poisson regression), and remission rate (Andreasen criteria, Fisher's exact test). Time course of PANSS will be visualized as arm $\times$ visit interaction plots with 95% CIs.

Secondary analyses: (1) Mixed-effects models for repeated measures (MMRM) will assess changes in MCCB scores and microbiota diversity over time; (2) Pearson or Spearman correlation analyses will examine associations between microbiota diversity and PANSS/MCCB scores; (3) Chi-square or Fisher's exact tests will be used for safety outcomes(17). Safety outcomes (adverse events by CTCAE grade and system organ class) will be compared between arms using Fisher's exact test. Laboratory parameters will be analyzed using the same LMM as the primary analysis. All statistical analyses will be performed using R 4.3 (lme4 package) or SPSS 26.0. No interim analyses are planned unless required by the Data Monitoring Committee for safety review.

To characterize FMT-induced alterations in gut microbial ecology, shotgun metagenomic sequencing data will be processed using MetaPhlAn 4 for taxonomic profiling. Alpha diversity will be assessed using the Shannon diversity index, observed species richness, and Pielou's evenness, with between-group comparisons performed using Wilcoxon rank-sum tests and longitudinal changes evaluated by linear mixed-effects models. Beta diversity will be calculated using Bray-Curtis dissimilarity and Aitchison distance, with principal coordinate analysis (PCoA) for

visualization and PERMANOVA (adonis2) for statistical testing of community structure differences between groups. Differential abundance analysis of species, gene families, and metabolic pathways will be performed using DESeq2 or MaAsLin2, adjusting for baseline covariates and multiple testing (false discovery rate < 0.2). Temporal dynamics of specific taxa of interest (e.g., short-chain fatty acid producers, Bifidobacterium species) will be examined using generalized estimating equations to account for within-subject correlations across repeated measures.

4.2 Capsule Fidelity

To evaluate, at the manufacturing-batch level, the degree to which the FMT capsule retains the original donor stool microbiota, capsule–stool Aitchison distance will be calculated. This is a quality-control step, not an engraftment analysis. Fidelity assessment provides context for interpreting apparent engraftment failures, taxa absent post-FMT may have been lost during capsule preparation rather than failing to colonize. Capsule–stool Aitchison distance is included as a candidate predictor in the engraftment determinants model, under the hypothesis that lower manufacturing fidelity predicts lower donor strain engraftment.

4.3 Engraftment and Statistical Analysis Plan

4.3.1 Microbiome Engraftment and Strain-Level Dynamics

To evaluate donor microbiota colonization efficiency post-FMT, strain-level profiling will be performed using StrainPhlAn 4, which infers strain identity through phylogenetic analysis of species-specific marker genes. Species-level metagenomic profiling will be conducted using MetaPhlAn 4(18). The definitions, formulas, and

470 applications of all engraftment metrics are summarized in **Table 7**. Detailed contents
471 for each assay are described in the **Supplementary Materials 2**. The phylogenetic
472 distance threshold for strain assignment will be determined using Youden's index(19,
473 20).

Table 7 Engraftment Metrics Summary

Metric	Definition	Formula	Notes	Application
Bray-Curtis (BC) Similarity	Community similarity based on relative abundances	$BC_{IJ} = \frac{\sum_{k=1}^p x_{ik} - x_{jk} }{\sum_{k=1}^p (x_{ik} + x_{jk})}$ <i>Bray – Curtis Similarity</i> = $1 - BC_{ij}$	x_{ik} and x_{jk} are the relative abundances of species k in samples i and j , and P is the total number of species	Assess compositional convergence between donor and recipient
Donor Similarity Index (DSI)	Normalized similarity to donor relative to baseline	$DSI_t = \frac{Sim(R_t, D) - Sim(R_0, D)}{1 - Sim(R_0, D)}$	R_0 is the recipient's baseline microbiota, R_t is the recipient's microbiota at post-FMT timepoint t , D is the donor's microbiota, and $Sim(.)$ is any chosen similarity metric (e.g., Bray-Curtis Similarity)	Track longitudinal shift toward donor profile

Jaccard Index (J)	Presence/absence-based species overlap	$J(A, B) = \frac{ A \cap B }{ A \cup B }$	A and B are the sets of species (or strains) detected in two samples (e.g., donor and post-FMT recipient)	Assess species-level donor strain engraftment
Aitchison Distance (AD)	CLR-transformed functional distance	$AD(x, y) = \sqrt{\sum_{k=1}^p \left[\ln\left(\frac{x_k}{g(x)}\right) - \ln\left(\frac{y_k}{g(y)}\right) \right]^2}$	x and y are composition vectors (e.g., functional profiles), and $g(x) = (\prod_{k=1}^p x_k)^{1/p}$ is the geometric mean of the composition	Evaluate functional vs taxonomic engraftment
Microbiota Compatibility Index (MCI)	Donor–recipient compatibility score (0– 100)	$MCI = \alpha \cdot P + \beta \cdot F + \gamma \cdot M$	P represents phylogenetic overlap between donor and recipient, F represents functional gene repertoire similarity, and M represents metabolic pathway complementarity.	Integrate phylogenetic, functional, and metabolic alignment

Strain Engraftment Rate (SER)	Proportion of donor strains detected post-FMT	$SER = \frac{N_{\text{donor strains detected post-FMT}}}{N_{\text{donor strains profiled}}}$	The weights α, β, γ are empirically determined, with $\alpha + \beta + \gamma = 1$ $N_{\text{donor strains profiled}}$ is the total number of donor strains for which strain-level profiling is feasible, and $N_{\text{donor strains detected post-FMT}}$ is the number of these donor strains successfully identified in the recipient after transplantation	Primary measure of donor colonization efficiency
Strain-Sharing Rate (SSR)	Shared strains between two samples	$SSR(A, B) = \frac{N_{\text{shared strains between A and B}}}{N_{\text{species profiled in common between A and B}}}$	$N_{\text{shared strains}}$ is the number of strains detected in both samples A and B, and $N_{\text{species profiled in common}}$ is the number of species for which	Compare donor–post, pre–post, and donor–pre pairs

			StrainPhlAn provided strain-level information in both samples	
Fraction of Donor Strains (FDS)	Donor-derived proportion of post-FMT strain pool	$FDS = \frac{N_{shared\ strains\ (post-FMT\ vs.\ donor)}}{N_{strains\ profiled\ in\ post-FMT}}$	<i>FDS</i> represents the proportion of the post-FMT recipient's strain pool that originated from the donor	Quantify donor contribution to recipient microbiome
Fraction of Retained Strains (FRS)	Baseline-derived proportion of post-FMT strain pool	$FRS = \frac{N_{shared\ strains\ (post-FMT\ vs.\ pre-FMT)}}{N_{strains\ profiled\ in\ post-FMT}}$	<i>FRS</i> represents the proportion of the post-FMT strain pool that persisted from the recipient's baseline	Quantify persistence of recipient baseline strains

Bray-Curtis dissimilarity quantifies compositional differences based on relative abundances and will be applied to donor-post-FMT, pre-FMT-post-FMT, and donor-pre-FMT pairs. Donor Similarity Index normalizes post-transplant similarity to the donor relative to baseline, enabling longitudinal tracking independent of inter-individual variation. Jaccard index measures presence/absence overlap, Aitchison distance assesses functional convergence via centered log-ratio (CLR) transformation, and Microbiota Compatibility Index (scaled 0 – 100) integrates phylogenetic, functional, and metabolic compatibility between donor and recipient, with scores ≥ 80 indicating high compatibility, 60 – 79 moderate compatibility, and <60 low compatibility(19, 21-24). SER is defined as the number of donor-derived strains detected post-FMT divided by the number of donor strains profiled(19). SSR quantifies shared strains between sample pairs, and FDS/FRS represent the donor-derived and baseline-retained proportions of the post-FMT strain pool. For comparison with published literature, we will additionally calculate the species-level engraftment rate without subtracting recipient-baseline strains.

We will explore donor-related (e.g., α -diversity, specific taxa abundance), recipient-related (e.g., baseline diversity, immune status), and procedural (e.g., bowel preparation) factors influencing engraftment using generalized linear models, adjusted for potential confounders. Candidate predictors for the engraftment determinants model will include: (1) Donor-related: donor alpha diversity, Bifidobacterium abundance, proportion of high-engraftment phyla (Bacteroidetes, Actinobacteria); (2) Recipient-related: pre-FMT alpha diversity (Shannon), pre-FMT beta distance to

donor, antibiotic-free interval, antipsychotic duration; (3) Capsule fidelity: capsule – stool Aitchison distance; (4) Procedural: cumulative FMT dose, administration frequency. The modeling strategy will proceed in three stages: (1) linear regression with all candidate predictors; (2) Random Forest for feature importance ranking; (3) LASSO regression for a parsimonious final model.

4.3.2 Engraftment–clinical association

We will examine the relationship between engraftment metrics and clinical outcomes using multiple approaches: (1) Spearman correlation between SER and Δ PANSS, adjusted for baseline PANSS and antipsychotic class via linear regression; (2) comparison of Δ PANSS between high-SER and low-SER groups (upper vs lower tertile) using Welch’s t-test or Mann–Whitney test, and response rate via χ^2 test; (3) Jonckheere–Terpstra test for trend across SER tertiles; and (4) LASSO regression selecting engrafted species as predictors of Δ PANSS to identify species-level contributions to clinical improvement.

4.3.3 Predictive Modeling of Post-FMT Microbiome Composition

Input features for species presence/absence prediction will include: donor species abundance, recipient baseline species abundance, species prevalence in healthy reference populations, and taxonomic affiliation. Model performance will be evaluated by 5-fold cross-validation and leave-one-donor-out validation, with AUROC as the primary metric(19, 20). For strain-level colonization prediction, LASSO regression will incorporate: recipient baseline species richness, donor-to-recipient species abundance ratio (propagule pressure), donor–recipient

strain population diversity, and phylogenetic distance(20).

Magnitude of Donor Colonization (MDC) and Loss of Baseline SGBs (Species-Level Genome Bins): The magnitude of donor colonization will be calculated as the sum of relative abundances of all donor-derived strains in the post-FMT recipient. SGBs detected in the recipient at baseline but no longer detectable post-FMT will be classified as “lost”(25). The proportion of lost SGBs will be calculated as the number of lost SGBs divided by the total number of SGBs present at baseline.

$$MDC = \sum_{i \in S_{shared}} RA_{i, post-FMT}$$

S_{shared} is the set of donor-derived strains detected in the post-FMT recipient, and $RA_{i, post-FMT}$ is the relative abundance of strain i in the post-FMT recipient sample.

$$Proportion\ of\ lost\ SGBs = \frac{N_{SGBs\ lost}}{N_{SGBs\ present\ at\ baseline}}$$

$$N_{SGBs\ lost} = |SGB_{baseline} \setminus SGB_{post-FMT}|$$

$SGB_{baseline}$ is the set of SGBs detected in the recipient at baseline, and $SGB_{post-FMT}$ is the set of SGBs detected post-FMT. The operator $\setminus$ denotes set difference (elements present in the first set but not in the second).

4.4 Multi-Omics Integrative Analysis

To systematically elucidate the mechanisms linking FMT-induced microbial changes to clinical outcomes, an integrated multi-omics analysis framework will be implemented across five complementary layers: (1) microbiome layer (fecal DNA): metagenomic species/strain composition and functional gene repertoire; (2) functional activity layer (fecal RNA): metatranscriptomic active pathway expression; (3)

metabolite layer (fecal raw and plasma): targeted and untargeted metabolomics of SCFAs, bile acids, and tryptophan metabolites; (4) immune response layer (plasma + PBMC): inflammatory proteomics, high-dimensional immunophenotyping, and PBMC functional assays; and (5) host transcriptome layer (PBMC bulk RNA): whole-blood gene expression profiling.

Integration across these layers will be performed using: (i) Spearman correlation to assess pairwise associations among microbiota, metabolites, immune factors, and clinical parameters; (ii) mediation analysis (mediation R package, 1,000 bootstrap iterations) to construct “microbiota-metabolite-immune-clinical” mediation models, reporting the average causal mediation effect (ACME); (iii) DIABLO (Data Integration Analysis for Biomarker discovery using Latent components) from the mixOmics package for multi-block integration; and (iv) canonical correspondence analysis (CCA) to assess associations between microbial communities and metabolite profiles. This five-layer integrative strategy enables systematic mapping of the “microbiota-metabolite-immune-clinical” landscape, consistent with the multi-omics methodological frameworks established in recent FMT trials(19, 20, 26, 27). The five-layer integrative framework, including data sources, analytical platforms, and targets for each layer, is summarized in **Table 8**.

Table 8 Multi-Omics Integrative Analysis Framework

Layer	Data Source	Technology	Analytical Target
-------	----------------	------------	-------------------

Layer 1:	Fecal	Shotgun	
Microbiome	DNA	metagenomics (Illumina NovaSeq)	Species/strain composition, functional gene repertoire
Layer 2:	Fecal	Metatranscriptomics	
Functional	RNA	(Illumina NovaSeq)	Active pathway expression
Activity			
		Targeted /	
Layer 3:	Fecal	untargeted	
Metabolome	raw /	metabolomics	SCFAs, bile acids (>20 species),
	plasma	(UHPLC-HRMS, LC-MS/MS)	tryptophan metabolites
		Inflammatory	
		proteomics	
Layer 4:	Plasma	(Olink/Luminex),	
Immune	/	spectral flow	Cytokine profiles, immune cell subsets,
Response	PBMC	cytometry (≥ 40 markers), functional assays	T-cell function
Layer 5: Host			
Transcriptome	PBMC	Bulk RNA-seq	Global gene expression (immune/mitochondrial pathways)
		Spearman	
Integration		correlation,	Map “microbiota–metabolite–immune–
Methods		mediation analysis, DIABLO, CCA	clinical” landscape

4.5 Sensitivity & Supplementary Analyses

To assess the robustness of the primary and secondary findings, a series of sensitivity and supplementary analyses will be performed as summarized in **Table 9**.

Table 9 Sensitivity and Supplementary Analyses

Analysis	Method	Purpose
Per-protocol analysis	Exclude major protocol violators (e.g., <80 capsules consumed, prohibited medications)	Assess robustness of ITT findings
Subgroup analyses	Stratify by sex, age (<40 vs ≥40 years), disease duration (<5 vs 5–10 years), antipsychotic class (olanzapine-based vs others), and baseline PANSS severity (median split); test interaction terms in LMM	Identify potential effect modifiers
Dietary adjustment	Include FFQ data as covariate in LMM (if collected)	Control for dietary confounding
Engraftment threshold sensitivity	Test alternative SER cut-offs (median, quartiles)	Validate robustness of high-vs-low engraftment comparisons
Missing data	MICE vs completers-only analysis	Evaluate impact of missing data on outcomes

Notes: ITT, intention-to-treat; LMM, linear mixed model; FFQ, food frequency questionnaire;

SER, strain engraftment rate; MICE, multiple imputation by chained equations.

5. Ethics and Compliance

5.1 Ethical Approval

This study protocol has been reviewed and approved by the Ethics Committee of the First Affiliated Hospital of Xi'an Jiaotong University (No: XJTU1AF2022LSK-435-01) and the ethics committees of all participating centers. The trial will be conducted in accordance with the principles of the Declaration of Helsinki (Fortaleza 2013 revision) and the International Conference on Harmonisation Good Clinical Practice (ICH-GCP E6 R2) guidelines.

5.2 Protocol Amendments

No changes have been made to the trial protocol since its commencement. The protocol described herein represents the final approved version. Any modifications to the study protocol that may affect the scientific integrity, participant safety, or conduct of the trial will be formally documented and submitted for review and approval by the lead Ethics Committee and all participating site ethics committees prior to implementation. Substantial amendments will also be updated in the *ClinicalTrials.gov* registry (NCT05850585) in accordance with regulatory requirements, and all amendments will be reported in the final trial publication, including the nature of the change, the date of amendment, and the rationale.

5.3 Informed Consent

Written informed consent will be obtained from all participants or their legally

authorized representatives (LARs) prior to enrollment. For participants with impaired decision-making capacity at screening, consent will be obtained from their LARs, and re-consent will be sought from the participant personally as soon as their decisional capacity is restored to the extent permitted by clinical evaluation. All participants (or their LARs) will be informed of the study's purpose, procedures, potential risks, benefits, and their right to withdraw from the study at any time without penalty or loss of standard medical care.

5.4 Data Protection and Confidentiality

All participant-identifiable information will be anonymized and replaced with unique coded identifiers. The electronic database will be password-protected, encrypted, and accessible only to authorized research personnel directly involved in the trial. Data will be retained for a period of 10 years after trial completion at the lead center's secure server, in compliance with national data protection regulations, and will be destroyed thereafter in accordance with the approved data management plan. Secondary use of anonymized data for future research purposes will require separate ethics approval and, where applicable, re-consent from participants.

5.5 Data Monitoring

An independent Data Monitoring Committee (DMC), comprising clinical experts, biostatisticians, and an ethicist, will review cumulative safety data at predefined intervals (every 3 months) throughout the trial. The DMC may recommend early termination or protocol modification if unacceptable safety signals emerge.

5.6 Trial Registration

This trial has been prospectively registered with the U.S. National Institutes of Health *ClinicalTrials.gov* database (registration number: NCT05850585). The results will be disseminated through peer-reviewed publications regardless of the outcome direction, and will be made publicly available in accordance with the International Committee of Medical Journal Editors (ICMJE) data-sharing requirements.

5.7 Patient and Public Involvement

Patients and the public were not involved in the design, conduct, or reporting of this trial protocol. The trial was designed by academic investigators based on preclinical evidence and prior clinical observations. However, the findings of this trial will be disseminated through peer-reviewed publications and presented at patient advocacy group meetings to ensure that results are accessible to the schizophrenia community. Patient representatives will be consulted for the development of patient-friendly summaries of the trial results upon study completion.

6. Discussion

Schizophrenia treatment faces the dual challenges of limited efficacy and significant adverse effects. The gut microbiota, through the microbiota-gut-brain axis, influences brain function and behavior, providing a novel therapeutic target for schizophrenia(28, 29). FMT has demonstrated efficacy in recurrent *Clostridium difficile* infection and certain gastrointestinal disorders, and has shown preliminary positive effects in autism spectrum disorder, yet its role in schizophrenia remains unclear(14). This trial addresses the gap by providing a rigorous evaluation of FMT as

an adjunctive therapy in non-treatment-resistant schizophrenia.

A critical determinant of FMT efficacy that warrants prospective evaluation is the degree of donor microbiota engraftment, the extent to which transplanted strains establish and persist in the recipient. Engraftment is shaped by donor-related factors (microbiota diversity, diet, and phylogenetic compatibility)(19, 24), recipient-related factors (baseline microbial diversity, immune status, and “gatekeeper” species such as *Bacteroides uniformis* and *Alistipes shahii*)(30), and procedural variables (bowel preparation, route, and dosing frequency)(19). Recipient factors, particularly pre-treatment dysbiosis characterized by reduced species richness, consistently outweigh donor factors in driving strain-level outcomes, suggesting that patients with more disrupted microbiomes may derive greater benefit(19, 24). Meta-analytic evidence supports a positive association between higher donor strain engraftment and clinical success ($P = 0.017$ across 24 studies)(19), although the heterogeneity of FMT responses across diseases underscores the need for disease-specific evaluation(23, 30). Strain-level engraftment patterns, including donor colonization, recipient persistence, and coexistence, may further refine outcome prediction(20), and machine learning classifiers using baseline donor-recipient features have achieved an average AUROC of 0.77 in predicting post-FMT species presence(19).

Conducting this trial across 11 centers presents several practical challenges. First, recruitment of non-treatment-resistant schizophrenia patients requires coordinated efforts across sites to ensure steady enrollment within the 18-month window. Early engagement with community mental health services and clear communication of the

trial' s adjunctive nature (standard antipsychotic treatment continues unchanged) will be essential to address potential participant concerns. Second, maintaining adherence to the 120-capsule regimen (10 capsules/day for 6 days, then 5 capsules/day for 12 days) requires robust monitoring. Third, the integrity of blinding depends on the visual and olfactory indistinguishability of FMT and placebo capsules, which is ensured by a single third-party manufacturer. Fourth, biospecimen handling, particularly fecal sample collection and storage at -80° C across multiple sites, requires standardized protocols for collection, aliquoting, and transport to maintain sample integrity for metagenomic, metatranscriptomic, and metabolomic analyses. Site staff will receive centralized training on these procedures.

The present trial is uniquely positioned to address these engraftment-related questions through its integrated multi-omics framework and strain-level tracking using StrainPhlAn 4. By systematically linking baseline microbial features, engraftment metrics (Bray-Curtis, DSI, Jaccard, Aitchison, MCI, SER, SSR, FDS/FRS), and clinical outcomes, this study will: (i) identify predictive biomarkers of FMT response in schizophrenia; (ii) elucidate whether engraftment magnitude correlates with PANSS improvement, MCCB cognitive gains, or metabolic changes; and (iii) generate hypotheses for personalized donor-recipient matching strategies. This mechanistic depth distinguishes the present trial from existing FMT studies in psychiatry, which have largely employed descriptive microbiome profiling without strain-level resolution.

The novelty of this trial lies in three aspects. First, it is the first multicenter RCT to

678 evaluate FMT specifically in non-treatment-resistant schizophrenia, complementing
679 the ongoing TRS-focused trial (ChiCTR2400092799) and addressing a substantially
680 larger patient population. Second, the integration of multidimensional data, clinical
681 symptoms, cognitive function (MCCB), gut metagenomics/metabolomics,
682 inflammatory proteomics, and high-dimensional immunophenotyping, enables
683 systematic exploration of the microbiota-immune-cognition pathway. Third, the use of
684 uniformly prepared FMT capsules by a third-party manufacturer ensures intervention
685 standardization and reproducibility(31).

686 Compared with the published TRS-FMT protocol (ChiCTR2400092799), the
687 present trial offers complementary advantages: a two-arm design providing greater
688 statistical power for the primary PANSS endpoint, exclusive use of oral capsules for
689 superior scalability and patient acceptability, systematic cognitive assessment via
690 MCCB, comprehensive metabolic monitoring, and a 24-week follow-up to assess
691 durability. While the TRS protocol addresses a high-unmet-need population, the
692 present study targets a broader group where earlier intervention may prevent treatment
693 resistance from consolidating.

694 Several limitations should be acknowledged. First, patients with schizophrenia
695 often present with multiple comorbidities and complex medication regimens, which
696 may confound the efficacy evaluation of FMT. Second, long-term efficacy and safety
697 beyond 24 weeks require further validation through extended follow-up studies(17).
698 Third, donor-recipient matching strategies have not been systematically implemented
699 in this trial, and the multi-omics analyses are inherently exploratory. Findings will

require independent replication before clinical translation(13).

Looking forward, the results of this trial will inform several key areas for future investigation. If engraftment metrics correlate with clinical response, subsequent studies could prospectively test donor selection algorithms to maximize colonization efficiency. The multi-omics data may reveal specific microbial taxa, metabolites, or immune signatures that mediate therapeutic effects, potentially guiding the development of next-generation microbiome therapeutics (e.g., defined bacterial consortia or postbiotics). Finally, the durability of FMT effects will determine whether maintenance dosing schedules are needed, an question that the 24-week follow-up is designed to preliminarily address.

In conclusion, this trial will provide high-level evidence on the efficacy and safety of FMT as an adjunctive therapy for non-treatment-resistant schizophrenia, contributing to the evidence base for microbiome-based interventions in psychiatric disorders and informing the design of future implementation studies.

Abbreviations

ACME	Average Causal Mediation Effect
ANCOVA	Analysis of Covariance
BSFS	Bristol Stool Form Scale
CCA	Canonical Correspondence Analysis
CFU	Colony-Forming Units

CGI	Clinical Global Impression
CTCAE	Common Terminology Criteria for Adverse Events
DIABLO	Data Integration Analysis for Biomarker discovery using Latent components
FDS	Fraction of Donor Strains
FMT	Fecal Microbiota Transplantation
FRS	Fraction of Retained Strains
GSRS	Gastrointestinal Symptom Rating Scale
HOMA-IR	Homeostatic Model Assessment of Insulin Resistance
ITT	Intention-to-Treat
LASSO	Least Absolute Shrinkage and Selection Operator
LMM	Linear Mixed Model
MCCB	MATRICES Consensus Cognitive Battery
MICE	Multiple Imputation by Chained Equations
PANSS	Positive and Negative Syndrome Scale
PP	Per-Protocol
RF	Random Forest
SER	Strain Engraftment Rate

SGBs	Species-Level Genome Bins
SSR	Strain-Sharing Rate
TESS	Treatment Emergent Symptom Scale

716

717 **Declarations**

718 **Ethics approval and consent to participate**

719 This study protocol has been reviewed and approved by the Ethics Committee of
720 the First Affiliated Hospital of Xi'an Jiaotong University (Approval No.
721 XJTU1AF2022LSK-435-01) and the ethics committees of all participating centers.
722 The additional movie files show this in more detail (see **Additional file 1 and 2**). The
723 trial will be conducted in accordance with the principles of the Declaration of
724 Helsinki (Fortaleza 2013 revision) and the International Conference on Harmonisation
725 Good Clinical Practice (ICH-GCP E6 R2) guidelines. Written informed consent will
726 be obtained from all participants or their legally authorized representatives prior to
727 enrollment. For participants with impaired decision-making capacity at screening,
728 consent will be obtained from their legally authorized representatives, and re-consent
729 will be sought from the participant personally as soon as their decisional capacity is
730 restored. All participants (or their legally authorized representatives) will be informed
731 of the study's purpose, procedures, potential risks, benefits, and their right to
732 withdraw from the study at any time without penalty or loss of standard medical care.

733 **Consent for publication**

Written informed consent will be obtained from all participants or their legally authorized representatives (LARs) prior to enrollment (see **Additional file 3**).

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was funded by the National Key Research and Development Program of China (No. 2025YFC2511200) and the National Natural Science Foundation of China (No. 82271572) (see **Additional file 4 and 5**).

Author's contributions

CW is the principal investigator of the grant; **CW, YL, MJ and JL** conceptualized the study. **CW, XM, BH, MJ, YJ, and YZ** adapted the intervention and developed the implementation strategies. **RG, XS, XM, and FZ** contributed to the refinement of the implementation strategies. **CW** drafted the manuscript, **YJ, MJ, XM, BH, MJ, YJ, YZ, RG, XS, XM, and FZ** revised it. All authors (**CW, YJ, MJ, XM, BH, MJ, YJ, YZ, RG, XS, XM, and FZ**) critically reviewed the manuscript for intellectual content and approved the final version.

Acknowledgements

We thank Dr. Dunbao Sun and Dr. Jiarong Tian (Ankang Central Hospital) for coordinating the participation of local hospitals; Dr. Dunhua Sun (Xunyang Disabled

Persons' Federation Bo'ai Hospital), Dr. Kang Liu (Ankang Jinzhou Mental Health Rehabilitation Hospital), Dr. Juntao Zhang (Ankang Antai Hospital), and Dr. Huitang Du (Ankang Hanbin District Third Hospital) for their substantial support in clinical implementation; and Qingyan Ma, Zai Yang, Yiyang Wang, Lu Yu, Yufeng Xun and Jing Liu (The First Affiliated Hospital of Xi'an Jiaotong University) for administrative and technical assistance. We also thank all participating centers, investigators, and study coordinators for their contributions to the trial.

CONSORT Checklist

This protocol is reported in accordance with the CONSORT 2025 Statement for randomized controlled trials. A completed CONSORT 2025 checklist is provided in **Additional file 6** (items pertaining to results will be completed at trial conclusion). The trial was prospectively registered at *ClinicalTrials.gov* (NCT05850585) prior to participant enrollment.

References

1. Luo W, Gao J, Guo Z, Han X, Song J, Su X, et al. Trends and cross-country inequalities in schizophrenia from 1990 to 2021, with prediction to 2035: a systematic analysis of the global burden of disease study 2021. *BMC Psychiatry*. 2025;25(1):928.
2. McCutcheon RA, Pillinger T, Varvari I, Halstead S, Ayinde OO, Crossley NA, et al. INTEGRATE: international guidelines for the algorithmic treatment of schizophrenia. *Lancet Psychiatry*. 2025;12(5):384–94.

- 778 3. Molstrom IM, Nordgaard J, Urfer-Parnas A, Handest R, Berge J, Henriksen MG. The prognosis of
779 schizophrenia: A systematic review and meta-analysis with meta-regression of 20-year follow-up
780 studies. *Schizophr Res.* 2022;250:152–63.
- 781 4. Romeo B, Willaime L, Rari E, Benyamina A, Martelli C. Efficacy of 5-HT_{2A} antagonists on negative
782 symptoms in patients with schizophrenia: A meta-analysis. *Psychiatry Res.* 2023;321:115104.
- 783 5. Leucht S, Priller J, Davis JM. Antipsychotic Drugs: A Concise Review of History, Classification,
784 Indications, Mechanism, Efficacy, Side Effects, Dosing, and Clinical Application. *Am J Psychiatry.*
785 2024;181(10):865–78.
- 786 6. Pillinger T, Howes OD, Correll CU, Leucht S, Huhn M, Schneider-Thoma J, et al. Antidepressant
787 and antipsychotic side-effects and personalised prescribing: a systematic review and digital tool
788 development. *Lancet Psychiatry.* 2023;10(11):860–76.
- 789 7. Dias MF, Nogueira YJA, Romano-Silva MA, Marques de Miranda D. Effects of antipsychotics on the
790 gastrointestinal microbiota: A systematic review. *Psychiatry Res.* 2024;336:115914.
- 791 8. Gao Y, Fan Y, Yang Z, Ma Q, Zhao B, He X, et al. Systems biological assessment of altered cytokine
792 responses to bacteria and fungi reveals impaired immune functionality in schizophrenia. *Mol*
793 *Psychiatry.* 2022;27(2):1205–16.
- 794 9. Pennisi E. Meet the psychobiome. *Science.* 2020;368(6491):570–3.
- 795 10. Settanni CR, Ianaro G, Bibbò S, Cammarota G, Gasbarrini A. Gut microbiota alteration and
796 modulation in psychiatric disorders: Current evidence on fecal microbiota transplantation. *Prog*
797 *Neuropsychopharmacol Biol Psychiatry.* 2021;109:110258.

798 11. Wei N, Ju M, Su X, Zhang Y, Huang Y, Rao X, et al. Transplantation of gut microbiota derived from
799 patients with schizophrenia induces schizophrenia-like behaviors and dysregulated brain transcript
800 response in mice. *Schizophrenia (Heidelb)*. 2024;10(1):44.

801 12. Zhu F, Ju Y, Wang W, Wang Q, Guo R, Ma Q, et al. Metagenome-wide association of gut
802 microbiome features for schizophrenia. *Nat Commun*. 2020;11(1):1612.

803 13. Chen Q, Wu C, Xu J, Ye C, Chen X, Tian H, et al. Donor-recipient intermicrobial interactions impact
804 transfer of subspecies and fecal microbiota transplantation outcome. *Cell Host Microbe*.
805 2024;32(3):349–65.e4.

806 14. Wan Y, Su Q, Ng SC. New insights on gut microbiome and autism. *Trends Mol Med*.
807 2024;30(12):1100–2.

808 15. Ding W, Li R, Gong L, Yi J, Wang Y, Xiao D, et al. Efficacy and Safety of Fecal Microbiota
809 Transplantation for Patients with Treatment-resistant Schizophrenia: A Double-blind, Randomized
810 Clinical Trial Protocol. *Curr Neuropsychopharmacol*. 2026.

811 16. Xiong Y, Krebs K, Jermy B, Karlsson R, Pasman JA, Nguyen TD, et al. Genome-wide association
812 meta-analysis and rare copy number variant analysis of treatment-resistant depression. *Mol Psychiatry*.
813 2025;30(11):5024–33.

814 17. Meng Y, Sun J, Zhang G. Pick fecal microbiota transplantation to enhance therapy for major
815 depressive disorder. *Prog Neuropsychopharmacol Biol Psychiatry*. 2024;128:110860.

816 18. Fernandes R, Jabbarizadeh B, Rajeh A, Hong MMY, Baines KJ, Ernst S, et al. Fecal microbiota
817 transplantation plus immunotherapy in metastatic renal cell carcinoma: the phase 1 PERFORM trial.
818 *Nat Med*. 2026;32(4):1325–36.

819 19. Ianiro G, Punčochář M, Karcher N, Porcari S, Armanini F, Asnicar F, et al. Variability of strain
820 engraftment and predictability of microbiome composition after fecal microbiota transplantation
821 across different diseases. *Nat Med.* 2022;28(9):1913–23.

822 20. Schmidt TSB, Li SS, Maistrenko OM, Akanni W, Coelho LP, Dolai S, et al. Drivers and determinants
823 of strain dynamics following fecal microbiota transplantation. *Nat Med.* 2022;28(9):1902–12.

824 21. Liu Y, Li X, Chen Y, Yao Q, Zhou J, Wang X, et al. Fecal microbiota transplantation: application
825 scenarios, efficacy prediction, and factors impacting donor-recipient interplay. *Front Microbiol.*
826 2025;16:1556827.

827 22. Kazemian N, Ramezankhani M, Sehgal A, Khalid FM, Kalkhoran AHZ, Narayan A, et al. The
828 trans-kingdom battle between donor and recipient gut microbiome influences fecal microbiota
829 transplantation outcome. *Sci Rep.* 2020;10(1):18349.

830 23. Reddi S, Senyshyn L, Ebadi M, Podlesny D, Minot SS, Gooley T, et al. Fecal microbiota
831 transplantation to prevent acute graft-versus-host disease: pre-planned interim analysis of donor
832 effect. *Nat Commun.* 2025;16(1):1034.

833 24. Behling AH, Portlock T, Ho D, Wilson BC, Paramsothy S, Kamm MA, et al. Cohort-specific
834 determinants of donor strain engraftment following multi-donor faecal microbiota transplantation in
835 two randomised clinical trials. *Gut Microbes.* 2025;17(1):2597628.

836 25. Duttagupta S, Messaoudene M, Hunter S, Desilets A, Jamal R, Mihalcioiu C, et al. Fecal
837 microbiota transplantation plus immunotherapy in non-small cell lung cancer and melanoma: the
838 phase 2 FMT-LUMINate trial. *Nat Med.* 2026;32(4):1337–50.

26. Li J, Wang Y, Li R, Huang H, Zhu X, Li X, et al. Adjunctive fecal microbiota transplantation for major depressive disorder: A randomized, double-blind, placebo-controlled trial. *Cell Host Microbe*. 2026;34(7):1367–82.e7.
27. Woodworth MH, Conrad RE, Haldopoulos M, Pouch SM, Babiker A, Mehta AK, et al. Fecal microbiota transplantation promotes reduction of antimicrobial resistance by strain replacement. *Sci Transl Med*. 2023;15(720):eabo2750.
28. Yassin LK, Skrabulyte-Barbulescu J, Alshamsi SH, Saeed S, Alkuwaiti SH, Almazrouei S, et al. The microbiota-gut-brain axis in mental and neurodegenerative disorders: opportunities for prevention and intervention. *Front Aging Neurosci*. 2025;17:1667448.
29. Nuzum ND, Deady C, Kittel-Schneider S, Cryan JF, O'Mahony SM, Clarke G. More than just a number: the gut microbiota and brain function across the extremes of life. *Gut Microbes*. 2024;16(1):2418988.
30. Porcari S, Benech N, Valles-Colomer M, Segata N, Gasbarrini A, Cammarota G, et al. Key determinants of success in fecal microbiota transplantation: From microbiome to clinic. *Cell Host Microbe*. 2023;31(5):712–33.
31. Hou S, Yu J, Li Y, Zhao D, Zhang Z. Advances in Fecal Microbiota Transplantation for Gut Dysbiosis-Related Diseases. *Adv Sci (Weinh)*. 2025;12(13):e2413197.

Figure Legend

Figure 1. CONSORT flow diagram of the trial design.

Eligible patients with non-treatment-resistant schizophrenia (PANSS total score ≥ 60 , disease duration ≤ 10 years) will be randomized 1:1 to receive either FMT capsules or placebo

capsules (identical in appearance, weight, and odor, containing lyophilized protectant only) adjunctive to standard antipsychotic treatment, with both groups following the same administration schedule. Follow-up assessments are scheduled at baseline (V1), day 7 (V2), day 18 (V3, end of intervention), week 4 (V4), week 8 (V5), week 12 (V6), and week 24 (V7, endpoint). Assessments include clinical scales (PANSS, MCCB, TESS, CGI, GSRS, BSFS), laboratory examinations (blood counts, liver/renal function, lipid profile, fasting insulin), and biospecimen collection (blood and feces) for multi-omics profiling. The analytical framework comprises four layers: Layer 1 evaluates the clinical efficacy of FMT versus placebo (primary and secondary outcomes). Layer 2 assesses capsule fidelity by comparing the finished capsule microbiota to the original donor stool. Layer 3 quantifies donor strain engraftment, identifies its determinants, and examines its association with clinical outcomes. Layer 4 integrates multi-omics data (microbiome, metabolome, immune, and host transcriptome) to explore mechanistic pathways. Clinical scale abbreviations: FMT, fecal microbiota transplantation; PANSS, Positive and Negative Syndrome Scale; MCCB, MATRICS Consensus Cognitive Battery; TESS, Treatment Emergent Symptom Scale; CGI, Clinical Global Impression; GSRS, Gastrointestinal Symptom Rating Scale; BSFS, Bristol Stool Form Scale. Statistical and analytical abbreviations are defined in the main text abbreviations table.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryMaterials12.docx](#)