



Outcome: Urinary continence at 3 months

Primary analysis: all study designs | random effects (REML) | OR > 1 favours treatment versus Conventional




Outcome: Urinary continence at 12 months

Primary analysis: all study designs | random effects (REML) | OR > 1 favours treatment versus Conventional




Outcome: Overall positive surgical margins

Primary analysis: all study designs | random effects (REML) | OR < 1 favours treatment versus Conventional




Outcome: Overall complications

Primary analysis: all study designs | random effects (REML) | OR < 1 favours treatment versus Conventional




Outcome: Clavien–Dindo grade I–II complications

Primary analysis: all study designs | random effects (REML) | OR < 1 favours treatment versus Conventional




Outcome: Clavien-Dindo grade $\geq$ III complications

Primary analysis: all study designs | random effects (REML) | OR < 1 favours treatment versus Conventional




Outcome: Operative time

Primary analysis: all study designs | random effects (REML) | MD < 0 favours treatment versus Conventional




Outcome: Estimated blood loss

Primary analysis: all study designs | random effects (REML) | MD < 0 favours treatment versus Conventional




Outcome: Immediate urinary continence

Primary analysis: all study designs | random effects (REML) | OR > 1 favours treatment versus Conventional




Outcome: Urinary continence at 2 weeks

Primary analysis: all study designs | random effects (REML) | OR > 1 favours treatment versus Conventional




Outcome: Urinary continence at 6 months

Primary analysis: all study designs | random effects (REML) | OR > 1 favours treatment versus Conventional




Outcome: Blood transfusion

Primary analysis: all study designs | random effects (REML) | OR < 1 favours treatment versus Conventional



Outcome: Console time

Primary analysis: all study designs | random effects (REML) | MD < 0 favours treatment versus Conventional



Outcome: Length of hospital stay

Primary analysis: all study designs | random effects (REML) | MD < 0 favours treatment versus Conventional