


Alpha Oscillatory Activity Associated with Impaired Visual Search in Cerebral Visual Impairment

Christopher R. Bennett, Corinna M. Bauer, Peter J. Bex, Davide Bottari, Lotfi B. Merabet

Supplemental Figure 1



Supplementary Figure 1. Thalamic Segmentation Map. Location of the 26 segmented thalamic nuclei used in the correlation analysis. Nuclei are identified based on a probabilistic atlas (Iglesias et al., 2018) and overlaid on axial T_1 -weighted images from a representative control participant. Note that the data from each hemisphere were evaluated separately. Slice projections move from ventral (left) to more dorsal (right) and are presented in radiologic convention. *Abbreviations:* AV: anteroventral, CeM: central medial, CL: central lateral, CM: centromedian, LD: laterodorsal, LGN: lateral geniculate, LP: lateral posterior, L-Sg: limitans (suprageniculata), MDI: mediodorsal lateral parvocellular, MDm: mediodorsal medial magnocellular, MGN: medial geniculate, MV(Re): reuniens (medial ventral), Pc: paracentral, Pf: parafascicular, Pt: paratenial, PuA: pulvinar anterior, Pul: pulvinar inferior, PuL: pulvinar lateral, PuM: pulvinar medial, R: reticular, VA: ventral anterior, VAmc: ventral anterior magnocellular, VLa: ventral lateral anterior, VLp: ventral lateral posterior, VM: ventromedial, VPL: ventral posterolateral.