Full wwPDB X-ray Structure Validation Report (i)

May 6, 2026 – 04:46 PM EDT

PDB ID	:	9ZY8 / pdb_00009zy8
Title	:	Crystal structure of human P450 CYP8B1 enzyme
Deposited on	:	2026-01-05
Resolution	:	2.79 Å(reported)

This wwPDB validation report is for manuscript review

This is a Full wwPDB X-ray Structure Validation Report.

This report is produced by the wwPDB biocuration pipeline after annotation of the structure.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)


For Manuscript Review

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	489	2% 80% 9% 10%

For Manuscript Review

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 3688 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 7-alpha-hydroxycholest-4-en-3-one 12-alpha-hydroxylase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	439	3629	2369	619	620	21	0	0	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	19	MET	-	expression tag	UNP Q9UNU6
A	20	ALA	-	expression tag	UNP Q9UNU6
A	21	LYS	-	expression tag	UNP Q9UNU6
A	22	LYS	-	expression tag	UNP Q9UNU6
A	23	THR	-	expression tag	UNP Q9UNU6
A	24	SER	-	expression tag	UNP Q9UNU6
A	25	SER	-	expression tag	UNP Q9UNU6
A	502	HIS	-	expression tag	UNP Q9UNU6
A	503	HIS	-	expression tag	UNP Q9UNU6
A	504	HIS	-	expression tag	UNP Q9UNU6
A	505	HIS	-	expression tag	UNP Q9UNU6
A	506	HIS	-	expression tag	UNP Q9UNU6
A	507	HIS	-	expression tag	UNP Q9UNU6


- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).

For Manuscript Review



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C Fe N O 43 34 1 4 4	0	0

For Manuscript Review

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	4	Total O 4 4	0	0


For Manuscript Review

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 7-alpha-hydroxycholest-4-en-3-one 12-alpha-hydroxylase

Chain A:



For Manuscript Review

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	57.82Å 89.18Å 107.29Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.31 – 2.79 35.31 – 2.79	Depositor EDS
% Data completeness (in resolution range)	99.8 (35.31-2.79) 99.8 (35.31-2.79)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.26 (at 2.81Å)	Xtriage
Refinement program	PHENIX (1.21.2_5419: ???)	Depositor
R, R_{free}	0.224 , 0.269 0.225 , 0.268	Depositor DCC
R_{free} test set	721 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å^2)	46.7	Xtriage
Anisotropy	0.265	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.30 , 26.7	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.92	EDS
Total number of atoms	3688	wwPDB-VP
Average B, all atoms (Å^2)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.71% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

For Manuscript Review

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	0.09	0/3736	0.22	0/5042

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3629	0	3580	25	0
2	A	12	0	16	0	0
3	A	43	0	30	0	0
4	A	4	0	0	0	0
All	All	3688	0	3626	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

For Manuscript Review

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:300:LYS:HG3	1:A:472:LEU:HD21	1.63	0.79
1:A:343:VAL:HG12	1:A:405:TYR:HB2	1.74	0.69
1:A:433:TRP:HB3	1:A:440:CYS:HB3	1.75	0.68
1:A:128:THR:HA	1:A:132:ARG:HD3	1.76	0.68
1:A:104:LYS:HG3	1:A:105:LYS:HE2	1.77	0.66
1:A:240:LEU:HD11	1:A:280:LEU:HD22	1.89	0.55
1:A:113:TYR:CD2	1:A:278:MET:HE1	2.48	0.49
1:A:33:PRO:HD3	1:A:374:TYR:HB3	1.95	0.48
1:A:110:VAL:HG11	1:A:213:PHE:HE1	1.80	0.47
1:A:445:PHE:O	1:A:449:GLU:HG3	2.14	0.47
1:A:50:ARG:HD2	1:A:218:TYR:HB3	1.96	0.46
1:A:98:ASP:OD2	1:A:101:GLN:HG2	2.17	0.45
1:A:90:LEU:HD11	1:A:431:MET:HE1	1.99	0.44
1:A:377:ARG:HB2	1:A:380:ASP:OD2	2.17	0.44
1:A:473:PRO:HB3	1:A:489:HIS:CE1	2.53	0.44
1:A:110:VAL:HA	1:A:216:PHE:CE2	2.53	0.43
1:A:110:VAL:HG11	1:A:213:PHE:CE1	2.53	0.43
1:A:124:HIS:CE1	1:A:441:PRO:HG3	2.54	0.43
1:A:202:LEU:HD21	1:A:239:MET:HG2	2.00	0.43
1:A:146:SER:O	1:A:150:VAL:HG12	2.19	0.42
1:A:356:LEU:HB2	1:A:384:LEU:HB2	2.00	0.42
1:A:185:LEU:HA	1:A:252:SER:HB3	2.03	0.41
1:A:440:CYS:HA	1:A:441:PRO:HD3	1.96	0.41
1:A:293:TRP:CG	1:A:351:ARG:HG3	2.55	0.40
1:A:122:MET:HG3	1:A:261:PHE:CG	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	423/489 (86%)	411 (97%)	12 (3%)	0	100 100

For Manuscript Review

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	392/436 (90%)	391 (100%)	1 (0%)	86 95

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	242	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	124	HIS
1	A	257	ASN
1	A	272	GLN
1	A	410	ASN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

For Manuscript Review

5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	HEM	A	603	4	50,50,50	1.97	16 (32%)	67,82,82	1.51	10 (14%)
2	GOL	A	601	-	5,5,5	1.08	0	5,5,5	0.02	0
2	GOL	A	602	-	5,5,5	1.07	0	5,5,5	0.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	603	4	-	3/14/54/54	-
2	GOL	A	601	-	-	4/4/4/4	-
2	GOL	A	602	-	-	2/4/4/4	-

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	603	HEM	FE-NC	7.22	2.19	1.95
3	A	603	HEM	FE-NA	3.72	2.07	1.95
3	A	603	HEM	CHD-C4C	3.67	1.45	1.38
3	A	603	HEM	CHC-C1C	3.26	1.44	1.38
3	A	603	HEM	CHA-C1A	3.23	1.46	1.39
3	A	603	HEM	CBA-CAA	2.80	1.61	1.51
3	A	603	HEM	C4C-NC	2.61	1.44	1.39
3	A	603	HEM	CHB-C4A	2.49	1.45	1.39
3	A	603	HEM	C3C-C4C	-2.44	1.41	1.46
3	A	603	HEM	FE-NB	2.44	2.02	1.94
3	A	603	HEM	C1B-C2B	2.42	1.49	1.44
3	A	603	HEM	C4A-NA	-2.42	1.35	1.39
3	A	603	HEM	C4B-NB	2.32	1.43	1.38

Continued on next page...

For Manuscript Review

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	603	HEM	C3C-C2C	2.19	1.41	1.37
3	A	603	HEM	CBC-CAC	2.19	1.40	1.30
3	A	603	HEM	C4A-C3A	2.02	1.47	1.43

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	A	603	HEM	C3B-C4B-NB	4.41	112.64	109.47
3	A	603	HEM	CHB-C1B-NB	3.87	129.16	124.37
3	A	603	HEM	CHC-C4B-NB	3.71	128.41	124.42
3	A	603	HEM	CHC-C4B-C3B	-3.07	118.95	125.07
3	A	603	HEM	C4A-NA-C1A	2.96	110.64	105.82
3	A	603	HEM	CHA-C1A-NA	2.57	128.52	123.86
3	A	603	HEM	CHB-C1B-C2B	-2.42	120.08	126.95
3	A	603	HEM	C1B-NB-C4B	-2.27	102.52	105.21
3	A	603	HEM	O2D-CGD-CBD	2.08	120.58	114.00
3	A	603	HEM	C4A-CHB-C1B	-2.01	121.51	126.25

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	GOL	O1-C1-C2-C3
2	A	601	GOL	C1-C2-C3-O3
2	A	602	GOL	C1-C2-C3-O3
2	A	601	GOL	O2-C2-C3-O3
3	A	603	HEM	C2B-C3B-CAB-CBB
3	A	603	HEM	C2C-C3C-CAC-CBC
3	A	603	HEM	C4B-C3B-CAB-CBB
2	A	602	GOL	O2-C2-C3-O3
2	A	601	GOL	O1-C1-C2-O2

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

For Manuscript Review

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand HEM A 603



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

For Manuscript Review

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	439/489 (89%)	0.29	12 (2%) 56 46	28, 45, 70, 99	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	188	TYR	3.3
1	A	239	MET	2.5
1	A	243	SER	2.4
1	A	314	GLN	2.3
1	A	91	LYS	2.3
1	A	468	PRO	2.3
1	A	462	ASP	2.3
1	A	437	VAL	2.2
1	A	418	ASP	2.2
1	A	434	GLY	2.1
1	A	332	GLY	2.1
1	A	251	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

For Manuscript Review


median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	GOL	A	601	6/6	0.85	0.15	42,49,57,61	0
2	GOL	A	602	6/6	0.93	0.12	42,48,53,54	0
3	HEM	A	603	43/43	0.95	0.11	31,40,53,67	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around HEM A 603:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.

For Manuscript Review