Tittle	**Differentiation of *Mitragyna speciosa*, a narcotic plant, from allied *Mitragyna* species using DNA barcoding-high-resolution melting (Bar-HRM) analysis**
Authors	Chayapol Tungphatthong[1,3], Santhosh Kumar J. Urumarudappa1[,3], Supita Awachai[1], Thongchai Sooksawate[2] and Suchada Sukrong[1]*
Affiliation	[1]Research Unit of DNA Barcoding of Thai Medicinal Plants, Department of Pharmacognosy and Pharmaceutical Botany, Faculty of Pharmaceutical Sciences, Chulalongkorn University, Bangkok 10330, Thailand
	[2]Department of Pharmacology and Physiology, Faculty of Pharmaceutical Sciences, Chulalongkorn University, Bangkok 10330, Thailand
	[3]These authors contributed equally: Chayapol Tungphatthong and Santhosh Kumar J. Urumarudappa
*Corresponding author	Professor Suchada Sukrong, Ph.D.
	Research Unit of DNA Barcoding of Thai Medicinal Plants,
	Department of Pharmacognosy and Pharmaceutical Botany,
	Faculty of Pharmaceutical Sciences, Chulalongkorn University,
	Bangkok 10330, Thailand
	Phone: +6681-819-6742, Fax: +6622-558-227
	Email: suchada.su@chula.ac.th



Fig. S1: Multiple sequence alignment of ITS2, *mat*K, *rbc*L and *trn*H-*psb*A intergenic spacer from four *Mitragyna* species showing the positions of primers used in HRM analysis. *Mitragyna* species indicates; MS- *M. speciosa*, MD- *M. diversifolia*, MH- *M. hirsuta* and MR- *M. rotundifolia*.