

Persistent phoQ upregulation and sigma-factor modulation link polymyxin resistance to broader stress responses in *Klebsiella pneumoniae* ST11

Michelly Maria Pereira e Oliveira

University of Pernambuco

Paula Mariana Salgueiro de Souza

`paulasalgueiro.db@ufrpe.br`

Federal Rural University of Pernambuco

Francisca Gama Silva

Federal Rural University of Pernambuco

Lara Valesca Mendonça da Costa Santos

Federal Rural University of Pernambuco

Rodrigo Tenório Gomes Pereira

University of Pernambuco

Márcia Maria Camargo Moraes

University of Pernambuco

Anna Carolina Soares Almeida

Federal Rural University of Pernambuco

Research Article

Keywords: antimicrobial resistance, lipid A remodeling, two-component regulatory systems, mgrB disruption, transcriptional regulation, reverse transcription quantitative PCR

Posted Date: October 19th, 2026

DOI: <https://doi.org/10.21203/rs.3.rs-10999995/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Additional Declarations: No competing interests reported.

Abstract

Background

Polymyxin resistance in *Klebsiella pneumoniae* frequently involves lipid A remodeling controlled by the PhoPQ and PmrAB two-component systems. However, interactions between these pathways and sigma factor-mediated stress responses remain incompletely understood in high-risk lineages.

Objective

To determine how polymyxin B exposure affects targeted transcriptional responses in the polymyxin-resistant *K. pneumoniae* sequence type 11 (ST11) clinical isolate Pol16 and to relate these responses to resistance- and stress-associated genomic alterations.

Methods

Pol16 was characterized by broth microdilution, whole-genome sequencing, and targeted sequence analysis. Expression of *phoQ*, *pmrD*, *pmrB*, *rpoD*, *rpoS*, and *rpoE* was quantified by reverse transcription quantitative PCR in exponential-phase cultures grown with or without 4 µg/mL polymyxin B. Expression was normalized to *rpoC* and calculated relative to *K. pneumoniae* ATCC 700603.

Results

Pol16 belonged to ST11 and carried a large deletion affecting *mgrB*, together with alterations in genes associated with two-component systems, lipid A modification, and σE regulation. *phoQ* was upregulated under both growth conditions and increased further following polymyxin exposure. Exposure also induced marked increases in *rpoD* (5.291-fold) and *rpoS* (2.453-fold). *pmrD* remained upregulated but did not differ significantly between the conditions tested, whereas *pmrB* remained below the calibration baseline. Although *rpoE* remained below baseline, its expression increased significantly after exposure.

Conclusion

The integrated genomic and transcriptional profile supports persistent PhoPQ signaling, potentially favored by *mgrB* disruption, accompanied by RpoD/RpoS-associated stress responses. The limited *pmrB* response and distinct *rpoE* behavior highlight a regulatory configuration that broadens the understanding of polymyxin resistance in a clinically important ST11 lineage.

Introduction

The prevalence of polymyxin resistance has increased among clinically important Gram-negative pathogens, including *K. pneumoniae*, a frequent cause of community- and hospital-acquired infections. Multidrug-resistant *K. pneumoniae* is recognized as a critical public-health threat, underscoring the need to clarify resistance mechanisms and improve control strategies [1–5].

Polymyxins are used as last-line agents against infections caused by multidrug-resistant Gram-negative bacteria. Their cationic residues bind to negatively charged phosphate groups in lipid A, disrupting the bacterial envelope. Resistance commonly arises when the net negative charge of lipid A is reduced through the addition of 4-amino-4-deoxy-L-arabinose (L-Ara4N) or phosphoethanolamine (pEtN) [6–11].

In *K. pneumoniae*, lipid A remodeling is regulated mainly by the PhoPQ, PmrAB, and CrrAB two-component systems. PhoPQ can activate the *arnBCADTEF-ugd* locus directly and can also communicate with PmrAB through the connector PmrD. Loss of the negative regulator MgrB is a frequent route to enhanced PhoPQ signaling and polymyxin resistance [5, 12]. In parallel, sigma factors coordinate basal transcription and stress-responsive programs, but changes in sigma-factor transcripts cannot by themselves establish sigma-factor activity or a direct contribution to resistance [13, 14].

This study characterized the polymyxin-resistant *K. pneumoniae* ST11 clinical isolate Pol16 and examined the expression of *phoQ*, *pmrD*, and *pmrB* genes, together with the sigma-factor genes *rpoD* ($\sigma 70$), *rpoS* (σS), and *rpoE* (σE), during growth with and without polymyxin B. Sequence alterations in polymyxin resistance-associated genes and in regulators of the σE pathway were also assessed.

Materials and Methods

Bacterial isolate and antimicrobial susceptibility testing

The *K. pneumoniae* isolate Pol16 was recovered from a urinary tract infection in a hospitalized patient in northeastern Brazil. Species identification was performed using the Vitek 2 system (bioMérieux, Marcy-l'Étoile, France). The polymyxin B minimum inhibitory concentration was determined by broth microdilution using the Policimbac kit (Probac, Brazil). The antimicrobial susceptibility profile is provided in Supplementary Table S1 (Online Resource 1).

Whole-genome sequencing and in silico analysis

Genomic DNA was extracted with the PureLink Genomic DNA Mini Kit (Invitrogen) and sequenced on the Illumina NextSeq 550 platform (Illumina, San Diego, CA, USA) using 2 × 75-bp paired-end reads. Reads were assembled with Velvet and the VelvetOptimiser script, using N50 optimization for k-mer selection [15]. Coding sequences were predicted and annotated with the NCBI Prokaryotic Genome Annotation Pipeline v6.1 [16]. The whole-genome shotgun sequence is available under GenBank accession JAJQQO000000000.1.

Resistance determinants were investigated with BIGSdb v1.34.0 [17], ResFinder v4.1 [18], and the Comprehensive Antibiotic Resistance Database [19]. Virulence-associated *loci* were assessed with BIGSdb, VirulenceFinder v2.0 [20–22], and VFDB Analyzer [23,24]. Kleborate v2.2.0 was used for species confirmation, multilocus sequence typing, and characterization of ICEKp-associated *loci*, capsule, K locus, and O antigen. Plasmid replicons were identified with PlasmidFinder v2.1 [21,22], and the sequence type was verified in PubMLST [25]. Tools were run with default settings unless otherwise stated; software details are provided in Supplementary Table S3 (Online Resource 1).

Coding and predicted protein sequences of *pmrA*, *pmrB*, *phoP*, *phoQ*, *crrA*, *crrB*, *arnA*, *arnB*, *arnC*, *arnD*, *arnT*, *mgrB*, *pmrD*, *ugd*, *eptA*, *eptB*, *rseA*, *rseB*, *rseP*, *surA*, *degS*, *rsd*, *rpoD*, *rpoS*, and *rpoE* were compared in Geneious Prime

v2023.0.4 and SnapGene 2023. *K. pneumoniae* ATCC 700603 (GenBank accession NZ_AOGO000000000.1) was used as the sequence reference. Alterations are summarized in Supplementary Table S2 and Figure S2 (Online Resource 1).

Primer design, RNA extraction, and RT-qPCR

Primers were designed from conserved target-gene regions using PrimerQuest. Specificity, GC content, melting temperature, and potential dimer formation were evaluated within the silico PCR and OligoAnalyzer. Amplification efficiency was determined from standard-curve slopes, and melt curves were inspected for single products and absence of detectable primer-dimer or nonspecific amplification (Table 1; Supplementary Figure S1, Online Resource 1). The *rpoC* primer pair was obtained from Gomes et al. [26].

Pol16 and *K. pneumoniae* ATCC 700603 cultures were grown in cation-adjusted Mueller-Hinton broth at 37 °C with shaking. Pol16 was cultivated with or without 4 µg/mL polymyxin B and harvested during exponential growth at OD600 = 0.5. Total RNA was extracted with TRIzol reagent (Thermo Fisher Scientific), treated with RQ1 RNase-Free DNase (Promega), and quantified with a NanoDrop Lite spectrophotometer (Thermo Scientific).

Complementary DNA was synthesized with the EasyScript Reverse Transcriptase/EasyScript cDNA Synthesis Kit (Applied Biological Materials). Expression of *phoQ*, *pmrD*, *pmrB*, *rpoD*, *rpoS*, and *rpoE* genes was quantified on an Amplio96 instrument (Locus) with SYBR Green I detection and 25 ng of total RNA equivalent per reaction. *rpoC* was used as the endogenous control, and polymyxin-susceptible *K. pneumoniae* ATCC 700603 as the calibration strain. Three independent biological replicates were analyzed, each in technical triplicate. Relative expression was calculated using the $2^{-\Delta\Delta C_q}$ method.

Statistical analysis

Between-condition comparisons were performed using Student's *t* test with Bonferroni correction for normally distributed data and the Wilcoxon test for non-normally distributed data. A two-sided *P* value < 0.05 was considered statistically significant. Analyses were performed in R using the stats package [27].

Ethics approval

The study was approved by the Research Ethics Committee of the University of Pernambuco, Brazil (reference 265.604).

Table 1 Primers used for RT-qPCR and amplification-performance parameters

Gene	Product/function	Primer sequence, 5'-3'	Amplicon bp	T _m , °C	GC, %	Efficiency, %	Slope
<i>phoQ</i>	PhoPQ sensor histidine kinase	F ACCCCTTTGCGCTGATAGAC R TGAGCCGGGAATTACACCC	83	64.2 64.2	55.0 55.0	98.537	-3.358
<i>pmrD</i>	PhoPQ-PmrAB connector protein	F GGCGACAGGCTTCAATCTCT R TACAGGACAACGCTTCGGC	87	64.1 64.3	55.0 57.9	97.980	-3.371
<i>pmrB</i>	PmrAB sensor histidine kinase	F TCACTCCTATCGTACTCAGC R TCGAGGGTCAGATTGAGG	94	60.1 60.0	50.0 55.6	102.936	-3.254
<i>rpoD</i>	RNA polymerase sigma factor RpoD (σ 70)	F CCGAGCTGTCTCAGGAAGAG R GTTGTCGTCATCGCTGTTGT	83	63.3 63.0	60.0 50.0	92.097	-3.527
<i>rpoS</i>	RNA polymerase sigma factor RpoS (σ S)	F CGCGAGTTGTCCCATAAG R CGTTCAGACGCAGCATAC	103	59.8 60.4	55.6 55.6	95.941	-3.423
<i>rpoE</i>	RNA polymerase sigma factor RpoE (σ E)	F CGGATCAGGTCCTTGTTG R CGCGACCTTATGCTGATAG	86	60.0 60.0	55.6 52.6	96.980	-3.396
<i>rpoC</i>	RNA polymerase β' subunit	F TATTCTGGTTCCACGCAACA R GGATACAACGGAACGCACTT	97	62.0 62.6	45.0 50.0	96.048	-3.420

T_m, melting temperature; GC, guanine-cytosine content. Target-gene primers were designed in this study; the *rpoC* primer pair was reported by Gomes et al. [26].

Results

Genomic findings associated with polymyxin resistance

Pol16 was classified as polymyxin resistant and assigned to sequence type ST11. Targeted sequence comparisons identified missense substitutions and deletions in genes encoding two-component systems and their regulators, including *pmrB*, *rrrB*, *pmrD*, and *mgrB* genes. A large deletion affected *mgrB* gene, the negative regulator of PhoQ. Alterations were also detected in the σ E regulatory genes *rseB*, *rseP*, and *degS* and in genes involved in L-Ara4N synthesis or transfer (*arnA*, *arnC*, *arnD*, and *arnT*). The remaining genes examined were intact or contained only synonymous substitutions relative to *K. pneumoniae* ATCC 700603. Detailed alterations and alignments are provided in Supplementary Table S2 and Figure S2 (Online Resource 1).

Target-gene expression during polymyxin B exposure

In Pol16 grown without polymyxin B, *phoQ* and *pmrD* expression was 2.542-fold and 3.648-fold, respectively, relative to the ATCC 700603 calibration strain. Expression of the *pmrB*, *rpoD*, *rpoS*, and *rpoE* genes remained below

the calibration baseline under this condition (Fig. 1).

Exposure to polymyxin B was associated with a significant increase in *phoQ* expression compared with untreated Pol16, while *pmrD* remained at a similar level at the different conditions. *pmrB* expression remained below the calibration baseline. The most pronounced change at the different conditions tested was observed for *rpoD*, which reached 5.291-fold expression relative to the calibration strain during polymyxin exposure. *rpoS* increased to 2.453-fold. Although *rpoE* increased significantly between Pol16 distinct culture conditions, its expression remained below the ATCC 700603 calibration baseline (Fig. 1).

Discussion

The findings of this study indicate that polymyxin resistance in *K. pneumoniae* Pol16 is associated with a regulatory profile dominated by persistent *phoQ* upregulation and accompanied by changes in transcriptional responses related to bacterial stress adaptation. The large deletion affecting *mgrB*, together with increased *phoQ* expression in both untreated and polymyxin-exposed conditions, provides a coherent genomic and transcriptional context for enhanced PhoPQ-associated signaling. Moreover, polymyxin exposure further increased *phoQ* expression and was accompanied by increased transcription of *rpoD* and *rpoS*, suggesting that exposure to the antibiotic intensifies not only the response linked to polymyxin resistance but also broader mechanisms involved in cellular adaptation to stress.

The PhoPQ system is one of the major regulatory pathways involved in polymyxin resistance in *K. pneumoniae*, particularly through modifications of lipid A that reduce the interaction of polymyxins with the bacterial outer membrane [12, 36, 37]. Inactivation of *mgrB*, a negative regulator of PhoPQ, is a well-established mechanism associated with constitutive activation of this system and polymyxin resistance [38, 39, 43, 44]. In Pol16, the presence of a large *mgrB* deletion together with constitutively increased *phoQ* transcription strongly supports the involvement of this regulatory axis in the resistant phenotype. The additional increase in *phoQ* expression during polymyxin exposure suggests that, despite an already elevated basal transcriptional state, the system remains responsive to antibiotic-induced envelope stress. This observation is relevant because PhoPQ activity is influenced by multiple environmental and regulatory signals and may be further modulated under conditions that perturb membrane homeostasis [40–42].

The transcriptional behavior of *pmrD* and *pmrB* provides additional insights into the organization of this response. The *pmrD* gene was upregulated in Pol16 relative to the calibration strain under both conditions, although its expression did not increase significantly following polymyxin exposure. PmrD functions as a connector between PhoPQ and PmrAB by stabilizing phosphorylated PmrA, thereby contributing to the regulatory network controlling lipid A remodeling [38, 40, 45, 46]. Its sustained expression in Pol16 may therefore reflect a regulatory background already adapted to polymyxin resistance rather than an acute response specifically induced by antibiotic exposure. In contrast, *pmrB* transcript levels remained below those observed in ATCC 700603 in both conditions. Taken together, these findings suggest that the transcriptional response detected in Pol16 is more strongly associated with the PhoPQ axis than with increased *pmrB* expression. This pattern is also compatible with evidence that PhoPQ can promote expression of lipid A modification pathways in *K. pneumoniae* without requiring strong transcriptional activation of the PmrAB system [38, 39].

Beyond these classical polymyxin resistance pathways, exposure of Pol16 to polymyxin B was associated with marked changes in sigma-factor transcription. *rpoD* showed a 5.291-fold increase during polymyxin exposure. RpoD (σ 70) is the primary housekeeping sigma factor and controls a broad range of genes required for bacterial

growth and cellular maintenance [14, 28]. Its increased expression may therefore reflect a substantial transcriptional adjustment to the physiological disturbance caused by polymyxin B rather than a resistance-specific response. At the same time, *rpoS*, encoding the general stress sigma factor σ S, was also induced during exposure. RpoS coordinates bacterial adaptation to multiple adverse conditions and contributes to survival during environmental and physiological stress [31, 32]. The simultaneous increase in *rpoD* and *rpoS* suggests that polymyxin exposure triggers a broader transcriptional reorganization in Pol16, combining maintenance of essential cellular functions with activation of stress-adaptation pathways.

The σ E pathway adds another layer to this response because it is closely associated with envelope homeostasis in Gram-negative bacteria. Pol16 carried sequence alterations in *degS*, *rseB*, and *rseP*, including a 171-bp deletion in *degS* and a 21-bp deletion together with an amino-acid substitution in *rseB*. These genes participate in the regulatory cascade controlling σ E release through the RseA anti-sigma factor [29, 30]. Although *rpoE* transcription remained below the calibration strain baseline, its expression increased significantly after polymyxin exposure. Thus, the σ E-associated response appears to remain responsive to antibiotic-induced envelope stress despite the genomic alterations identified in components of its regulatory pathway. The combination of genomic variation and inducible *rpoE* expression suggests a potentially modified regulatory architecture that may influence how Pol16 responds to membrane damage [29, 30, 33–35].

Collectively, these findings support a model in which polymyxin resistance in Pol16 is associated with a constitutively altered regulatory background centered on the *mgrB*–PhoPQ axis, while polymyxin exposure further amplifies *phoQ* transcription and recruits broader stress-related responses involving RpoD, RpoS, and, to a lesser extent, RpoE. Rather than representing isolated transcriptional events, these changes indicate coordinated adaptation to the membrane and physiological stress imposed by polymyxin B. The comparatively stable expression of *pmrD* and low *pmrB* transcription further indicate that the different regulatory components involved in polymyxin resistance do not respond uniformly to antibiotic exposure.

Because the study focused on a single clinical isolate and a targeted set of transcripts, these findings primarily define the regulatory profile of Pol16 and provide a basis for further mechanistic investigation. Functional analyses of pathway activity and lipid A composition, as well as evaluation of additional polymyxin-resistant isolates, will be important to determine how broadly this transcriptional pattern is present. Nevertheless, the combined genomic and transcriptional data provide evidence that the polymyxin response in Pol16 extends beyond classical resistance determinants and involves a broader regulatory adaptation to antibiotic-induced stress.

Conclusion

Together, these findings support a resistance-associated regulatory profile predominantly linked to the *mgrB*–PhoPQ axis and accompanied by a broader transcriptional response to polymyxin-induced stress. This integrated genomic and transcriptional perspective expands the understanding of regulatory responses associated with polymyxin resistance in high-risk ST11 *K. pneumoniae*.

Declarations

Funding

This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brazil (CAPES), Finance Code 001. Financial support was also provided by the Fundação de Amparo à Ciência e

Competing interests

The authors declare no financial or non-financial interests that are directly or indirectly related to this work.

Author contributions

Michelly Maria Pereira e Oliveira: Methodology, Investigation, Writing - original draft. Paula Mariana Salgueiro de Souza: Writing - review and editing. Patrícia Francisca Gama da Silva: Investigation. Lara Valesca Mendonça da Costa Santos: Formal analysis. Rodrigo Tenório Gomes Pereira: Investigation. Márcia Maria Camargo de Moraes: Conceptualization, Writing - review and editing. Anna Carolina Soares Almeida: Conceptualization, Supervision, Data curation, Writing - review and editing.

Data availability

The whole-genome shotgun sequence of *K. pneumoniae* Pol16 is available in GenBank under accession JAJQQ0000000000.1. Additional data supporting the study are included in the article and its supplementary information.

Ethics approval

This study was approved by the Research Ethics Committee of the University of Pernambuco, Brazil (reference 265.604).

References

1. European Centre for Disease Prevention and Control (2022) Antimicrobial resistance surveillance in Europe 2022 – 2020 data. <https://www.ecdc.europa.eu/en/publications-data/antimicrobial-resistance-surveillance-europe-2022-2020-data>. Accessed 20 March 2023
2. World Health Organization (2024) WHO bacterial priority pathogens list 2024: bacterial pathogens of public health importance to guide research, development, and strategies to prevent and control antimicrobial resistance. World Health Organization, Geneva
3. Centers for Disease Control and Prevention (2022) The biggest antibiotic-resistant threats in the United States. <https://www.cdc.gov/drugresistance/biggest-threats.html>. Accessed 28 March 2023
4. Martin RM, Bachman MA (2018) Colonization, infection, and the accessory genome of *Klebsiella pneumoniae*. *Front Cell Infect Microbiol* 8:4
5. Stojowska-Swędryńska K, Łupkowska A, Kuczyńska-Wiśnik D, Laskowska E (2021) Antibiotic heteroresistance in *Klebsiella pneumoniae*. *Int J Mol Sci* 23:449
6. Andrade FF, Silva D, Rodrigues A, Pina-Vaz C (2020) Colistin: update on its mechanism of action and resistance, present and future challenges. *Microorganisms* 8:1716

7. MacNair CR, Stokes JM, Carfrae LA, Fiebig-Comyn AA, Coombes BK, Mulvey MR et al (2018) Overcoming *mcr-1*-mediated colistin resistance with colistin in combination with other antibiotics. *Nat Commun* 9:458
8. Boszczowski I, Salomão MC, Moura ML, Freire MP, Guimarães T, Cury AP et al (2019) Multidrug-resistant *Klebsiella pneumoniae*: genetic diversity, mechanisms of resistance to polymyxins and clinical outcomes in a tertiary teaching hospital in Brazil. *Rev Inst Med Trop Sao Paulo* 61:e29
9. Hancock RE (1997) Peptide antibiotics. *Lancet* 349:418–422
10. Olaitan AO, Morand S, Rolain JM (2014) Mechanisms of polymyxin resistance: acquired and intrinsic resistance in bacteria. *Front Microbiol* 5:643. <https://doi.org/10.3389/fmicb.2014.00643>
11. Sabnis A, Hagart KL, Klöckner A, Becce M, Evans LE, Furniss RCD et al (2021) Colistin kills bacteria by targeting lipopolysaccharide in the cytoplasmic membrane. *eLife* 10:e65836
12. Huang J, Li C, Song J, Velkov T, Wang L, Zhu Y et al (2020) Regulating polymyxin resistance in Gram-negative bacteria: roles of two-component systems PhoPQ and PmrAB. *Future Microbiol* 15:445–459
13. Helmann JD, Zaher H, MacLellan SR (2021) Transcription: sigma factors. In: Jez J (ed) *Encyclopedia of Biological Chemistry III*. Elsevier, Oxford, pp 379–382
14. Paget MS, Helmann JD (2003) The $\sigma 70$ family of sigma factors. *Genome Biol* 4:203
15. Zerbino DR, Birney E (2008) Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res* 18:821–829
16. Wyres KL, Wick RR, Gorrie C, Jenney A, Follador R, Thomson NR et al (2016) Identification of *Klebsiella* capsule synthesis loci from whole-genome data. *Microb Genom* 2:e000102. <https://doi.org/10.1099/mgen.0.000102>
17. Jolley KA, Maiden MC (2010) BIGSdb: scalable analysis of bacterial genome variation at the population level. *BMC Bioinformatics* 11:595
18. Florensa AF, Kaas RS, Clausen PTLC, Aytan-Aktug D, Aarestrup FM (2022) ResFinder: an open online resource for identification of antimicrobial resistance genes in next-generation sequencing data and prediction of phenotypes from genotypes. *Microb Genom* 8:e000748. <https://doi.org/10.1099/mgen.0.000748>
19. Alcock BP, Huynh W, Chalil R, Smith KW, Raphenya AR, Wlodarski MA et al (2023) CARD 2023: expanded curation, support for machine learning, and resistome prediction at the Comprehensive Antibiotic Resistance Database. *Nucleic Acids Res* 51:D690–D699
20. Joensen KG, Scheutz F, Lund O, Hasman H, Kaas RS, Nielsen EM et al (2014) Real-time whole-genome sequencing for routine typing, surveillance, and outbreak detection of verotoxigenic *Escherichia coli*. *J Clin Microbiol* 52:1501–1510
21. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K et al (2009) BLAST+: architecture and applications. *BMC Bioinformatics* 10:421
22. Clausen PTLC, Aarestrup FM, Lund O (2018) Rapid and precise alignment of raw reads against redundant databases with KMA. *BMC Bioinformatics* 19:307
23. Liu B, Zheng D, Zhou S, Chen L, Yang J (2022) VFDB 2022: a general classification scheme for bacterial virulence factors. *Nucleic Acids Res* 50:D912–D917
24. Zhou S, Liu B, Zheng D, Chen L, Yang J (2025) VFDB 2025: an integrated resource for exploring anti-virulence compounds. *Nucleic Acids Res* 53:D871–D877
25. Jolley KA, Bray JE, Maiden MCJ (2018) Open-access bacterial population genomics: BIGSdb software, the PubMLST.org website and their applications. *Wellcome Open Res* 3:124

26. Gomes AEI, Stuchi LP, Siqueira NMG, Henrique JB, Vicentini R, Ribeiro ML et al (2018) Selection and validation of reference genes for gene expression studies in *Klebsiella pneumoniae* using reverse-transcription quantitative real-time PCR. *Sci Rep* 8:9001. <https://doi.org/10.1038/s41598-018-27420-2>
27. Hornik K, R Core Team (2024) The R FAQ. <https://CRAN.R-project.org/doc/manuals/R-FAQ.html>. Accessed 9 September 2025
28. Treviño-Quintanilla LG, Freyre-González JA, Martínez-Flores I (2013) Anti-sigma factors in *Escherichia coli*: common regulatory mechanisms controlling sigma factor availability. *Curr Genomics* 14:378–387
29. Zeng X, Hinenoya A, Guan Z, Xu F, Lin J (2023) Critical role of the RpoE stress response pathway in polymyxin resistance of *Escherichia coli*. *J Antimicrob Chemother* 78:732–746
30. Paget MS (2015) Bacterial sigma factors and anti-sigma factors: structure, function and distribution. *Biomolecules* 5:1245–1265
31. Lonetto M, Gribskov M, Gross CA (1992) The sigma 70 family: sequence conservation and evolutionary relationships. *J Bacteriol* 174:3843–3849
32. Weber H, Polen T, Heuveling J, Wendisch VF, Hengge R (2005) Genome-wide analysis of the general stress response network in *Escherichia coli*: σ S-dependent genes, promoters, and sigma factor selectivity. *J Bacteriol* 187:1591–1603
33. Helmann JD (1999) Anti-sigma factors. *Curr Opin Microbiol* 2:135–141
34. Pinto D, Liu Q, Mascher T (2019) ECF σ factors with regulatory extensions: the one-component systems of the σ universe. *Mol Microbiol* 112:399–409
35. Wösten MM (1998) Eubacterial sigma factors. *FEMS Microbiol Rev* 22:127–150
36. Chen HD, Groisman EA (2013) The biology of the PmrA/PmrB two-component system: the major regulator of lipopolysaccharide modifications. *Annu Rev Microbiol* 67:83–112
37. Jayol A, Nordmann P, Brink A, Poirel L (2015) Heteroresistance to colistin in *Klebsiella pneumoniae* associated with alterations in the PhoPQ regulatory system. *Antimicrob Agents Chemother* 59:2780–2784
38. Chen AI, Albicoro FJ, Zhu J, Goulian M (2021) Effects of regulatory network organization and environment on PmrD connector activity and polymyxin resistance in *Klebsiella pneumoniae* and *Escherichia coli*. *Antimicrob Agents Chemother* 65:e00889–e00820
39. Cheung CHP, Dulyayangkul P, Heesom KJ, Avison MB (2020) Proteomic investigation of the signal transduction pathways controlling colistin resistance in *Klebsiella pneumoniae*. *Antimicrob Agents Chemother* 64:e00790–e00720
40. Groisman EA, Duprey A, Choi J (2021) How the PhoP/PhoQ system controls virulence and Mg²⁺ + homeostasis. *Microbiol Mol Biol Rev* 85:e00176–e00120
41. Han J, Gao X, Luo X, Zhu L, Zhang Y, Dong P (2023) Role of the PhoP/PhoQ system in regulating stress adaptation responses in *Escherichia coli* O157:H7. *Food Microbiol* 112:104244
42. Soncini FC, Vescovi EG, Groisman EA (1995) Transcriptional autoregulation of the *Salmonella typhimurium* *phoPQ* operon. *J Bacteriol* 177:4364–4371
43. Aires CAM, Pereira PS, Asensi MD, Carvalho-Assef APD (2016) *mgrB* mutations mediating polymyxin B resistance in *Klebsiella pneumoniae* isolates from rectal surveillance swabs in Brazil. *Antimicrob Agents Chemother* 60:6969–6972
44. Bray AS, Smith RD, Hudson AW, Hernandez GE, Young TM, George HE et al (2022) MgrB-dependent colistin resistance in *Klebsiella pneumoniae* is associated with increased host-to-host transmission. *mBio* 13:e03595–

45. Elias R, Duarte A, Perdigão J (2021) A molecular perspective on colistin and *Klebsiella pneumoniae*: mode of action, resistance genetics, and phenotypic susceptibility. *Diagnostics* 11:1165
46. Li L, Ma J, Cheng P, Li M, Yu Z, Song X et al (2023) Roles of two-component regulatory systems in *Klebsiella pneumoniae*: regulation of virulence, antibiotic resistance, and stress responses. *Microbiol Res* 272:127374

Figures




Figure 1

Relative expression of *phoQ*, *pmrB*, *pmrD*, *rpoD*, *rpoE*, and *rpoS* in *K. pneumoniae* Pol16 grown with 4 µg/mL polymyxin B (black bars) or without antibiotic (gray bars). Expression was normalized to *rpoC* and calculated relative to *K. pneumoniae* ATCC 700603 using the $2^{-\Delta\Delta C_q}$ method. The red line denotes the calibration baseline (fold change = 1). Bars show mean $\pm$ standard deviation for three biological replicates. Brackets compare the two Pol16 growth conditions by Student's *t* test. **P* < 0.05; ***P* $\leq$ 0.01; ****P* $\leq$ 0.001; n.s., not significant

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryMaterial1.docx](#)