

# Research Integrated clinical, environmental, and healthcare-worker hand surveillance for unit-level infection prevention prioritisation in a Mongolian tertiary hospital: a two-component observational study

**Odgerel Tsogbadrakh**

**zogbadrakhodgerel@gmail.com**

National University of Mongolia

**Tumenjargal Davaasuren**

National University of Mongolia

**Gantumur Tsagaan**

Medclean LLC

**Uyanga Tsogtbaatar**

Second State Central Hospital

**Orkhon Munkhtsetseg**

Second State Central Hospital

**Khulan Battulga**

Second State Central Hospital

**Tsendsuren Tumur**

Second State Central Hospital

**Shuree Enkh-Amgalan**

National University of Mongolia

**Aminaa Bayaraa**

National University of Mongolia

**Anudari Badral**

National University of Mongolia

---

## Research Article

**Keywords:** healthcare-associated infection, multidrug-resistant organisms, environmental contamination, healthcare-worker hands, methicillin-resistant *Staphylococcus aureus*, infection prevention and control, antimicrobial resistance, Mongolia

**Posted Date:** October 7th, 2026

**DOI:** <https://doi.org/10.21203/rs.3.rs-11017403/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.  
[Read Full License](#)

**Additional Declarations:** No competing interests reported.

---

# Abstract

## Background

Integrating clinical surveillance with environmental cultures and healthcare-worker (HCW) hand sampling may help target infection prevention and control (IPC) action, but environmental or hand recovery alone does not establish transmission. We assessed unit-level IPC priorities in a Mongolian tertiary hospital and contextualised the institutional profile using national surveillance.

## Methods

We combined a retrospective analysis of anonymised, IPC-reviewed, laboratory-positive healthcare-associated infection (HAI) records from 28 December 2022 to 31 December 2025 with a cross-sectional survey conducted in six units from 16 June to 3 July 2025. The survey sampled 131 high-touch surfaces, nine field-negative controls, and 68 HCW hand imprints. Cross-sectional isolates underwent MALDI-TOF mass spectrometry identification and VITEK 2 susceptibility testing. Monthly phenotype-confirmed MDRO-positive HAI records were analysed by negative-binomial regression with patient-days as an offset.

## Results

Across 43,922 admissions and 275,922 patient-days, 2,458 HAIs were recorded (8.91 per 1,000 patient-days). Of 1,819 IPC-classified MDRO-positive HAI cases, 91.6% occurred in four service categories. Among 1,999 organisms recovered from these records, MRSA (32.3%) and ESBL-producing *Escherichia coli* (23.4%) predominated; *Acinetobacter baumannii* was most frequent in intensive care. After exclusion of six data-incomplete 2023 months, the temporal model did not provide evidence of an increase (incidence-rate ratio per year 1.08, 95% confidence interval 0.98–1.18;  $p = 0.11$ ). Twenty-two of 131 surfaces (16.8%) were MDRO-positive, and 23 of 68 HCW hand imprints (33.8%) were MRSA-positive. Surgery Unit I and the intensive care unit were ranked top priority, and the Stroke Unit high priority.

## Conclusions

Integrated surveillance identified a small number of units where clinical burden overlapped with environmental-surface and HCW-hand contamination. These findings support risk-stratified IPC and antimicrobial stewardship while remaining insufficient to infer directional transmission or clonal linkage.

## Background

Antimicrobial resistance (AMR) and healthcare-associated infections (HAIs) remain major threats to patient safety, particularly in tertiary hospitals where invasive care, broad-spectrum antimicrobial

exposure, and high patient acuity concentrate risk [1–5]. Within these settings, multidrug-resistant organisms (MDROs) may be distributed unevenly across clinical units, making local surveillance essential for targeted infection prevention and control (IPC).

High-touch surfaces, shared equipment, and healthcare-worker (HCW) hands can form an ecological interface through which clinically important organisms persist or are transiently redistributed [6–11]. Environmental or hand cultures can therefore identify priorities for IPC verification, but they cannot establish source-reservoir relationships or direction of transmission without molecular epidemiology [12]. High-touch frequency is also an imperfect proxy for transmission relevance because mobile shared items and network-central surfaces may be more important than touch counts alone [13].

Mongolian national surveillance indicates substantial antimicrobial resistance pressure. The ARMOR-MN 2019–2023 report documented high resistance among major healthcare-associated pathogens and identified *Acinetobacter baumannii* as a leading organism in critical care [14]. Earlier Mongolian studies have reported under-recognised HAI burden, elevated device-associated infection rates, resistant ICU pathogens, and substantial Gram-negative MDRO carriage among hospitalised patients [15–19].

However, adult tertiary-hospital studies integrating longitudinal clinical surveillance with environmental and HCW-hand microbiology at unit level remain scarce, particularly in resource-constrained settings. We therefore combined three years of IPC-reviewed HAI surveillance with an epidemiologically directed cross-sectional survey of high-touch surfaces and HCW hands. Our objectives were to describe institutional and unit-level HAI and MDRO burden, identify environmental and hand contamination in selected units, and use converging signals to inform transparent operational IPC prioritisation. This integration addresses a practical gap: how routine and targeted surveillance can guide unit-level action without treating environmental concordance as proof of transmission. National surveillance was used only for qualitative contextualisation.

## Methods

### Study design and setting

This two-component observational study was conducted in an anonymised tertiary referral hospital in Ulaanbaatar, Mongolia. The retrospective component used laboratory-positive records reviewed and classified by the hospital IPC service as part of routine HAI/MDRO surveillance from 28 December 2022 to 31 December 2025. The study team received the IPC-validated dataset and performed secondary analyses of case counts, organism composition, unit distribution, and temporal patterns. Units for cross-sectional sampling were selected before microbiological results were known, using approximately 30 months of available surveillance data, clinical burden, patient acuity, invasive-care density, and feasibility. Sampling was undertaken from 16 June to 3 July 2025 in Anaesthesiology-Intensive Care (ICU), Isolation, Stroke, Surgery Unit I, Surgery Unit II, and the Sterilization Unit.

## Case definitions and denominators

The IPC service classified laboratory-positive records using institutional criteria aligned with CDC/NHSN HAI definitions: the admission day was counted as day 1, infections meeting criteria on or after day 3 were classified as healthcare-associated, and surgical-site infections used the relevant 30- or 90-day surveillance window [20]. Present-on-admission or community-onset infections, asymptomatic colonisation, duplicate records, and records with insufficient classification data were excluded during routine IPC review. For case-level analyses, each IPC-validated MDRO-positive HAI record was counted once. For isolate-level summaries, all clinically reported pathogenic organisms from the same microbiology result were retained separately; therefore, polymicrobial records could contribute more than one organism. Admissions and patient-days were obtained from hospital statistics. Incidence density was standardised to total patient-days because unit-level device-days were not consistently available. The surveillance system retained the admitting unit after transfer, which could shift apparent burden toward admitting services.

## Environmental and hand sampling

High-touch surfaces were sampled from patient-zone, staff-contact, and shared-equipment categories using a sterile 10 x 10 cm template and a pre-moistened sterile nylon-flocked swab applied in horizontal, vertical, and diagonal strokes with rotation. Swabs were transported in Amies medium at 2–8°C and processed within two hours. Field-negative controls were sterile swabs opened at the sampling site without surface contact and processed identically. Hand imprints were collected immediately after patient care and before hand hygiene by pressing the fingertips and palm of the dominant hand onto nutrient agar for five seconds. They were interpreted as point-in-time transient contamination rather than persistent colonisation. Participation was voluntary and based on written informed consent.

## Microbiology

For the cross-sectional survey, primary culture used 5% sheep-blood agar, MacConkey agar, Endo agar, nutrient agar, mannitol-salt agar, and CHROMagar MRSA, incubated at 37°C and read at 18–24 h; CHROMagar MRSA was extended to 48 h when required. Representative morphotypes were subcultured to purity. Species identification was performed by MALDI-TOF MS (VITEK MS), and antimicrobial susceptibility testing by VITEK 2, interpreted according to CLSI M100 (35th edition, 2025) [21]. For phenotypic classification, MRSA, ESBL-producing Enterobacterales, carbapenem-resistant Enterobacterales (CRE), and carbapenem-resistant *A. baumannii* (CRAB) were assigned from species identification and available susceptibility results. Formal MDR/XDR/PDR classification was not assigned because the routine susceptibility panels did not consistently cover all antimicrobial categories required by Magiorakos et al. [22]. In retrospective isolate-level descriptions, *A. baumannii* and *P. aeruginosa* were retained as species labels when a standardised resistance phenotype was not available; accordingly,

these labels were not treated as phenotype-confirmed MDRO categories in the primary temporal analysis.

## Statistical analysis and national contextualisation

Burden, organism composition, and unit-level distribution were summarised descriptively. Monthly IPC-classified MDRO-positive records from January 2023 to December 2025 were modelled using a conservative phenotype-based definition: MRSA, ESBL-producing Enterobacterales, CRE, and VRE when reported. Negative-binomial regression was used because of marked overdispersion, with the natural logarithm of monthly patient-days as an offset; results are reported as incidence-rate ratios (IRRs) with 95% confidence intervals (CIs). Data completeness was checked against the official infection register. Six 2023 months had implausibly low counts, and an incomplete February–June 2025 microbiology export was recovered from a secondary extract that matched the register. The complete-month model, excluding the six data-incomplete months, was treated as the primary interpretable temporal analysis; the full 36-month model and the 2024–2025 restriction were sensitivity analyses. Unit-level prioritisation descriptively juxtaposed clinical-burden categories used during the interim surveillance review with surface MDRO positivity and MRSA-positive HCW hand imprints. Priority tiers reflected convergence of operational signals and were intended to guide IPC verification, not to provide a validated risk score or infer transmission. For national contextualisation, annual organism–antibiotic resistance proportions were extracted from the ARMOR-MN 2019–2023 consolidated report [14] and visualised by the authors in R version 4.5.2. Only organism–antibiotic–year combinations with at least 30 tested isolates were displayed (Additional file 1: Figure S1); the corresponding machine-readable data are provided in Additional file 2. Institutional and national findings were compared qualitatively because the data sources differed in denominator and unit of analysis.

## Use of generative artificial intelligence

ChatGPT (OpenAI) was used to support English-language editing, structural revision, and journal formatting during manuscript preparation; it was not used to generate the study data. All authors reviewed and revised the resulting text and take full responsibility for the manuscript.

## Ethics

The study was conducted in accordance with the ethical principles set out in UNESCO's Universal Declaration on Bioethics and Human Rights (2005). The retrospective component used anonymised routine surveillance and laboratory records without direct patient contact; individual consent was not required. The cross-sectional component was approved by the study hospital's Research Department (approval no. 2025/06/05/01; 5 June 2025), and participating HCWs provided written informed consent. All data were anonymised and reported in aggregate.

# Results

## Institutional and unit-level HAI and MDRO burden

During the surveillance period, the hospital recorded 43,922 admissions and 275,922 patient-days. A total of 2,458 HAIs were recorded, corresponding to an incidence density of 8.91 per 1,000 patient-days and a cumulative incidence of 5.60 per 100 admissions (Table 1). There were 1,819 IPC-classified MDRO-positive HAI cases. The associated records yielded 1,999 clinically reported organisms. MDRO-positive HAI burden was concentrated in General Surgery (632 cases, 34.7%) and Anaesthesiology-ICU (529, 29.1%), followed by Other wards (264, 14.5%) and the Emergency Department (241, 13.2%); together, these services accounted for 91.6% of all MDRO-positive HAI cases (Fig. 1). General Surgery was recorded as an aggregate service in retrospective surveillance, whereas Surgery Units I and II were sampled separately by ward location.

Table 1  
Institutional HAI and MDRO burden, December 2022–  
December 2025.

| Indicator                                      | Value  |
|------------------------------------------------|--------|
| Admissions                                     | 43922  |
| Patient-days                                   | 275922 |
| Total HAIs                                     | 2458   |
| HAI incidence density (per 1,000 patient-days) | 8.91   |
| HAI cumulative incidence (per 100 admissions)  | 5.60   |
| MDRO-positive HAI cases                        | 1819   |
| Organisms from MDRO-positive HAI records       | 1999   |

MDRO, multidrug-resistant organism. Each IPC-validated MDRO-positive HAI record was counted once in case-level analyses. All clinically reported pathogenic organisms from the same microbiology result were retained in isolate-level summaries; therefore, case and organism counts differ. Incidence density was standardised to total patient-days because unit-level device-days were not consistently recorded.

Among organisms recovered from MDRO-positive HAI records, pus/wound specimens (699, 35.0%) and the heterogeneous 'other' category (610, 30.5%) predominated, followed by urine (286, 14.3%) and sputum (170, 8.5%). The most frequent organism/phenotype groups were MRSA (646, 32.3%), ESBL-producing *Escherichia coli* (468, 23.4%), *Acinetobacter baumannii* (295, 14.8%), *Pseudomonas aeruginosa* (196, 9.8%), and CRE (135, 6.8%) (Table 2). The species-only labels *A. baumannii* and *P. aeruginosa* represent organisms recovered from IPC-classified MDRO-positive records and should not be interpreted as uniformly phenotype-confirmed MDRO categories. The strongest specimen-organism intersection was pus/wound-MRSA (412 isolates; 58.9% of pus/wound isolates and 63.8% of all MRSA). Urine was

dominated by ESBL-producing *E. coli* (171 isolates), whereas sputum was dominated by *A. baumannii* (56 isolates). In General Surgery, MRSA (309) and ESBL-producing *E. coli* (262) predominated; in the ICU, *A. baumannii* (237), MRSA (120), and CRE (59) were most frequent.

Table 2  
Specimen types and leading organisms from MDRO-positive HAI records.

| Category                                                  | n   | %    |
|-----------------------------------------------------------|-----|------|
| <b>Specimen type</b>                                      |     |      |
| Pus / wound                                               | 699 | 35.0 |
| Other                                                     | 610 | 30.5 |
| Urine                                                     | 286 | 14.3 |
| Sputum                                                    | 170 | 8.5  |
| Bile                                                      | 96  | 4.8  |
| Peritoneal fluid                                          | 64  | 3.2  |
| Swab / smear                                              | 20  | 1.0  |
| Blood                                                     | 20  | 1.0  |
| <b>Leading organism/phenotype groups</b>                  |     |      |
| Methicillin-resistant <i>Staphylococcus aureus</i> (MRSA) | 646 | 32.3 |
| ESBL-producing <i>Escherichia coli</i>                    | 468 | 23.4 |
| <i>Acinetobacter baumannii</i>                            | 295 | 14.8 |
| <i>Pseudomonas aeruginosa</i>                             | 196 | 9.8  |
| Carbapenem-resistant Enterobacterales (CRE)               | 135 | 6.8  |
| Mixed / multiple                                          | 89  | 4.5  |
| ESBL-producing <i>Klebsiella</i> spp.                     | 62  | 3.1  |
| <i>Stenotrophomonas maltophilia</i>                       | 27  | 1.4  |

ESBL, extended-spectrum beta-lactamase. *Acinetobacter baumannii* and *Pseudomonas aeruginosa* were retained as species-level labels when standardised resistance phenotypes were unavailable; they were not included in the conservative temporal definition.

Across the 30 data-complete months, conservative-definition MDRO incidence averaged 7.5 records per 1,000 patient-days (range 4.9–12.0). Six 2023 months contained implausibly low official counts; a separate February–June 2025 export gap was recovered from a secondary extract. In the complete-

month model, there was no evidence of a temporal increase (IRR per year 1.08, 95% CI 0.98–1.18;  $p = 0.11$ ). The full-period model produced a positive estimate (IRR 1.35, 95% CI 1.28–1.43;  $p < 0.001$ ), but this estimate was driven by the six data-incomplete months. The 2024–2025 restriction was also non-significant (IRR 1.12, 95% CI 0.97–1.31;  $p = 0.13$ ; Table 3; Fig. 2). The temporal findings therefore do not support a robust increase in phenotype-confirmed MDRO-positive HAI records.

Table 3  
Temporal models of monthly phenotype-confirmed MDRO-positive HAI records.

| Specification                    | Months | IRR per year | 95% CI; p              |
|----------------------------------|--------|--------------|------------------------|
| Excluding data-incomplete months | 30     | 1.08         | 0.98–1.18; $p = 0.11$  |
| Full period                      | 36     | 1.35         | 1.28–1.43; $p < 0.001$ |
| 2024–2025 only                   | 24     | 1.12         | 0.97–1.31; $p = 0.13$  |

IRR, incidence-rate ratio; CI, confidence interval. The complete-month model excluding six data-incomplete 2023 months is the primary interpretable analysis. The full-period estimate is shown to demonstrate the effect of incomplete surveillance capture.

## Environmental and hand contamination

All nine field-negative controls showed no growth, arguing against systematic contamination during collection or transport. Of 131 high-touch surfaces, 44 (33.6%) were culture-positive and 22 (16.8%) were MDRO-positive, yielding 65 organisms (mean 1.48 organisms per positive surface). Fourteen of 44 positive surfaces (31.8%) were polymicrobial. Surface MDRO positivity was highest in the Stroke Unit (7/16, 43.8%) and lowest in the Sterilization Unit (0/11) (Fig. 3; Table 4). MRSA was the most frequent environmental MDRO (21/65 organisms, 32.3%), followed by *Pseudomonas aeruginosa*, *Acinetobacter nosocomialis*, and *Staphylococcus hominis* (five each); one CRAB isolate was recovered in the Stroke Unit.

Among 68 HCW hand imprints, 34 (50.0%) showed growth and 23 (33.8%) were MRSA-positive. Because samples were collected after patient care and before hand hygiene, they represent transient point-in-time contamination rather than persistent colonisation. MRSA positivity was highest in Surgery Unit I (10/25, 40.0%) and the ICU (10/27, 37.0%). By profession, positivity was 14/36 (38.9%) among nurses and 7/22 (31.8%) among physicians; comparisons are descriptive because denominators were small and uneven. One *Klebsiella pneumoniae* isolate was recovered from an ICU physician's hand.

Table 4  
Cross-sectional surface and hand sampling outcomes by unit.

| Unit                | Surfaces n | Culture-positive | MDRO-positive | Hand MRSA-positive |
|---------------------|------------|------------------|---------------|--------------------|
| Surgery Unit I      | 24         | 9 (37.5%)        | 8 (33.3%)     | 10/25 (40.0%)      |
| Anaesthesiology–ICU | 22         | 10 (45.5%)       | 3 (13.6%)     | 10/27 (37.0%)      |
| Stroke Unit         | 16         | 10 (62.5%)       | 7 (43.8%)     | 2/6 (33.3%)        |
| Isolation Unit      | 18         | 5 (27.8%)        | 3 (16.7%)     | 1/7 (14.3%)        |
| Surgery Unit II     | 40         | 8 (20.0%)        | 1 (2.5%)      | 0/3 (0.0%)         |
| Sterilization Unit  | 11         | 2 (18.2%)        | 0 (0.0%)      | —                  |
| Total               | 131        | 44 (33.6%)       | 22 (16.8%)    | 23/68 (33.8%)      |

All nine field-negative controls showed no growth. Hand-imprint and surface denominators differ. The 68 hand imprints comprised samples from nurses (n = 36), physicians (n = 22), and residents (n = 10). Hand sampling was a convenience sample with uneven per-unit numbers, limiting between-unit comparison.

## Unit-level integration and operational prioritisation

Integration of the three data sources produced an exploratory four-tier operational prioritisation (Fig. 4). Surgery Unit I and the ICU combined high clinical burden with measurable environmental-surface and HCW-hand contamination and were ranked top priority. The Stroke Unit, which had the highest environmental-surface MDRO positivity (43.8%) despite lower clinical burden, was ranked high priority. The Isolation Unit and Surgery Unit II were assigned to the watch tier, and the Sterilization Unit served as a baseline reference. Because the final retrospective dataset reported General Surgery as an aggregate service while cross-sectional sampling distinguished Surgery Units I and II, the matrix should be interpreted as a descriptive IPC planning tool rather than a validated comparative score. It does not establish transmission or clonal linkage.

## Comparison with national surveillance

The institutional profile was qualitatively consistent with ARMOR-MN national surveillance. National data identified *Staphylococcus aureus* and *Escherichia coli* as dominant organisms and *A. baumannii* as a leading critical-care pathogen [14]. Similarly, MRSA and ESBL-producing *E. coli* predominated overall in this study, while *A. baumannii* was most frequent in the ICU. The cross-sectional recovery of MRSA and one CRAB isolate also occurred against a national background of substantial resistance among these organisms [14]. This contextual comparison is qualitative only because the national report is laboratory-based isolate surveillance, whereas the present study analysed IPC-reviewed HAI records and a targeted cross-sectional survey. The annual organism–antibiotic resistance landscape used for this qualitative contextualisation is provided in Additional file 1: Figure S1.

## Discussion

This single-centre study combined three years of IPC-reviewed surveillance with an epidemiologically directed survey of high-touch surfaces and HCW hands. Three findings are most relevant for practice. First, IPC-classified MDRO-positive HAI burden was concentrated in a small number of services. Second, the organism profile differed by setting, with MRSA and ESBL-producing *E. coli* prominent in surgery and *A. baumannii* prominent in intensive care. Third, triangulation of clinical, surface, and hand signals identified Surgery Unit I, the ICU, and the Stroke Unit as priorities for IPC verification. This integrated profile would not have been visible from clinical or environmental data alone.

Concentration in surgical and intensive-care services is clinically plausible because these settings combine invasive procedures, wound burden, devices, high antimicrobial exposure, and frequent staff-patient contact. The observed organism pattern is consistent with national surveillance and with major healthcare-associated priority pathogens [12, 14, 23]. These findings support unit-specific microbiology review, locally tailored empirical-therapy guidance, and focused IPC audits rather than a uniform response across all wards.

The temporal analysis does not demonstrate a robust increase in phenotype-confirmed MDRO burden. The full-period association was driven by six data-incomplete 2023 months and was not sustained after their exclusion or in the 2024–2025 restriction. The time series should therefore be interpreted as evidence of surveillance-data fragility rather than a confirmed temporal rise. Future prospective surveillance should incorporate standardised case capture, antimicrobial-use data, staffing and occupancy measures, device-days, carriage surveillance, and molecular typing.

The cross-sectional results also require caution. Negative field controls support that observed growth reflected the sampled clinical environment rather than systematic collection or transport contamination. However, surface or hand recovery does not establish patient-to-environment, environment-to-hand, or hand-to-patient transmission [6–11, 24]. Hand imprints were collected after patient care and before hand hygiene; they therefore provide a high-risk process signal, not evidence of persistent carriage. The ICU *Klebsiella pneumoniae* hand isolate should be interpreted as a prompt for IPC verification, not as a transmission chain.

The Stroke Unit finding is operationally important because the highest surface MDRO positivity occurred despite a lower clinical case count. Potential explanations include shared-equipment use, patient dependency, cleaning-cycle effects, or unmeasured colonisation pressure. The appropriate response is targeted verification: observe cleaning practice and disinfectant contact time, review shared-equipment workflows, audit hand hygiene, and repeat sampling after corrective feedback. High-touch selection is a practical proxy, but it does not identify contact-network hubs; future work should combine microbiology with hand-to-surface network methods and genotyping [13].

Recent ICU contact-network work reinforces this distinction: surfaces most relevant to transmission may not be those touched most often, particularly when shared mobile items act as network hubs [13]. Our

study did not measure contact structure, and its single-timepoint surface survey cannot capture the full cleaning-recontamination cycle. These limitations define a practical next step rather than weaken the immediate value of targeted IPC verification.

Operationally, the findings support horizontal IPC foundations across all units, including hand-hygiene performance, environmental-cleaning reliability, and device-care bundles [25–27]. These should be paired with unit-specific microbiology review and antimicrobial stewardship [28]. For Surgery Unit I and the ICU, priorities are hand-hygiene audit and feedback, validation of cleaning for shared equipment, and review of empirical therapy. For the Stroke Unit, environmental cleaning and shared-equipment workflows warrant priority attention. Watch-tier units should undergo repeat review if clinical surveillance or follow-up sampling escalates.

Strengths include the integration of three years of IPC-reviewed laboratory-positive surveillance, prespecified negative-binomial sensitivity analyses, epidemiologically directed environmental sampling, HCW hand sampling with informed consent, and qualitative comparison with contemporary national surveillance. The study separates operational prioritisation from claims of transmission and reports data-completeness limitations transparently.

The limitations are substantial. This observational, single-centre design cannot establish transmission, clonal linkage, or the absence of pseudo-outbreaks. The retrospective dataset included only laboratory-positive, IPC-reviewed records and may not represent the full HAI burden. The temporal series was affected by data-completeness artefacts, while the broad sensitivity definition included species labels without uniformly coded resistance phenotypes. The surveillance system retained the admitting unit after transfer, which may misallocate burden. Surface sampling was a single snapshot, hand sampling was a convenience sample with uneven unit denominators, and baseline hand-hygiene compliance, cleaning audits, staffing ratios, device-days, and nasal MRSA carriage were not systematically measured. Carbapenem resistance was AST-derived, carbapenemase genes were not investigated, formal MDR/XDR/PDR classification was not assigned, and national comparison was qualitative rather than a numerical benchmark. The findings therefore identify priorities for targeted IPC verification, not baseline IPC performance or directional spread.

## Conclusions

In this tertiary hospital, IPC-classified MDRO-positive HAI burden was concentrated in a small number of services and overlapped with environmental-surface and HCW-hand contamination in selected units. This integrated pattern supports risk-stratified IPC: maintain horizontal prevention foundations across the hospital, then intensify targeted cleaning, hand-hygiene, equipment-workflow, microbiology-review, and antimicrobial-stewardship measures in the highest-priority units. Because transmission was not confirmed molecularly, the findings provide operational priorities and a baseline for prospective, genotyping-enabled multi-centre work rather than proof of directional spread.

# Abbreviations

|              |                                                                              |
|--------------|------------------------------------------------------------------------------|
| AMR          | antimicrobial resistance                                                     |
| AST          | antimicrobial susceptibility testing                                         |
| CI           | confidence interval                                                          |
| CRAB         | carbapenem-resistant <i>Acinetobacter baumannii</i>                          |
| CRE          | carbapenem-resistant Enterobacterales                                        |
| ESBL         | extended-spectrum beta-lactamase                                             |
| HAI          | healthcare-associated infection                                              |
| HCW          | healthcare worker                                                            |
| ICU          | intensive care unit                                                          |
| IPC          | infection prevention and control                                             |
| IRR          | incidence-rate ratio                                                         |
| MALDI-TOF MS | matrix-assisted laser desorption/ionisation time-of-flight mass spectrometry |
| MDR          | multidrug-resistant                                                          |
| MDRO         | multidrug-resistant organism                                                 |
| MRSA         | methicillin-resistant <i>Staphylococcus aureus</i>                           |
| PDR          | pandrug-resistant                                                            |
| VRE          | vancomycin-resistant Enterococcus                                            |
| XDR          | extensively drug-resistant.                                                  |

# Declarations

## **Ethics approval and consent to participate**

The study was conducted in accordance with the ethical principles set out in UNESCO's Universal Declaration on Bioethics and Human Rights (2005). Permission to conduct the study was granted by the Deputy Director for Medical Care of the Second State Central Hospital, and ethical approval was obtained from the hospital's Ethics Committee. The retrospective component used anonymised routine surveillance and laboratory records without direct patient contact; therefore, individual informed consent was not required. For the cross-sectional component, participating healthcare workers (HCWs) provided written informed consent prior to sample collection. All study data were anonymised and reported only in aggregate form.

## **Consent for publication**

Not applicable.

## **Competing interests**

OT and GT are affiliated with Medclean LLC, which funded the study. The remaining authors declare that they have no competing interests.

## **Funding**

Medclean LLC provided financial and material support for the study. OT and GT, who are affiliated with Medclean LLC, contributed as investigators as described below. Beyond those investigator contributions, the funding organisation had no additional role in data analysis, interpretation, manuscript preparation, or the decision to submit the manuscript.

## **Author Contribution**

OT conceived and designed the study, developed the methodology, coordinated the project, supervised data collection, performed the statistical analysis, interpreted the findings, prepared the figures and tables, and drafted the manuscript. TD, GT, UT, OM, KB, TT, SEA, AB, and ABa contributed to data collection, study implementation, and critical revision of the manuscript. All authors reviewed, revised, and approved the final version of the manuscript.

## **Acknowledgements**

The authors thank the participating hospital units and microbiology laboratory staff for their cooperation.

# Data Availability

The datasets generated and/or analysed during the current study are not publicly available because they contain information from a tertiary hospital and could compromise institutional confidentiality but are available from the corresponding author on reasonable request.

## References

1. GBD 2021 Antimicrobial Resistance Collaborators. Global burden of bacterial antimicrobial resistance 1990–2021: a systematic analysis with forecasts to 2050. *Lancet*. 2024;404:1199–226. 10.1016/S0140-6736(24)01867-1.
2. Murray CJL, Ikuta KS, Sharara F, et al. Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *Lancet*. 2022;399:629–55.
3. World Health Organization. Global report on infection prevention and control. Geneva: WHO; 2022.
4. Centers for Disease Control and Prevention. COVID-19: U.S. impact on antimicrobial resistance, special report 2022. Atlanta: CDC; 2022.
5. European Centre for Disease Prevention and Control. Antimicrobial resistance surveillance in Europe (EARS-Net), 2019–2023. Stockholm: ECDC; 2024.
6. Kramer A, Schwebke I, Kampf G. How long do nosocomial pathogens persist on inanimate surfaces? A systematic review. *BMC Infect Dis*. 2006;6:130.
7. Otter JA, Yezli S, Salkeld JAG, French GL. Evidence that contaminated surfaces contribute to the transmission of hospital pathogens and an overview of strategies to address contaminated surfaces in hospital settings. *Am J Infect Control*. 2013;41(5 Suppl):S6–11.
8. Huang SS, Datta R, Platt R. Risk of acquiring antibiotic-resistant bacteria from prior room occupants. *Arch Intern Med*. 2006;166:1945–51.
9. Bhalla A, et al. Acquisition of nosocomial pathogens on hands after contact with environmental surfaces near hospitalized patients. *Infect Control Hosp Epidemiol*. 2004;25:164–7.
10. Stiefel U, et al. Contamination of hands with methicillin-resistant *Staphylococcus aureus* after contact with environmental surfaces and after contact with the skin of colonized patients. *Infect Control Hosp Epidemiol*. 2011;32:185–7.
11. Thom KA, et al. Environmental contamination because of multidrug-resistant *Acinetobacter baumannii* surrounding colonized or infected patients. *Am J Infect Control*. 2011;39:711–5.
12. Carroll KC, Pfaller MA, Karlowsky JA, Landry ML, McAdam AJ, Patel R, Pritt BS, editors. Manual of clinical microbiology. 13th ed. Washington, DC: ASM; 2024.
13. Zheng Y, Ni K, Ji S, Lu Q, Hu Y, Wu F, et al. High-touch' surfaces are not always 'high-risk' surfaces in the intensive care unit environment. *J Hosp Infect*. 2026;171:99–106. 10.1016/j.jhin.2026.03.001.
14. National Center for Communicable Diseases (Mongolia). ARMOR-MN Surveillance Network. Consolidated report on antimicrobial resistance, 2019–2023 [in Mongolian]. Ulaanbaatar: NCCD;

2026.

15. Ider B-E, Clements A, Adams J, Whitby M, Muugolog T. Prevalence of hospital-acquired infections and antibiotic use in two tertiary Mongolian hospitals. *J Hosp Infect.* 2010;75:214–9.
16. Ider BE, Baatar O, Rosenthal VD, Khuderchuluun C, Baasanjav B, Donkhim C, et al. Multicenter study of device-associated infection rates in hospitals of Mongolia: findings of the International Nosocomial Infection Control Consortium (INICC). *Am J Infect Control.* 2016;44:327–31. 10.1016/j.ajic.2015.10.010.
17. Baljin B, Gurjav U, Tulgaa K, Baldan G, Gunchin B, Sandag T, et al. High acquisition rate of Gram-negative multidrug-resistant organism colonization during hospitalization: a perspective from a high-endemic setting. *Infect Drug Resist.* 2021;14:3919–27. 10.2147/IDR.S328139.
18. Bataar O, Khuderchuluun C, Lundeg G, Chimeddorj S, Brunauer A, Gradwohl-Matis I, Duenser MW. Rate and pattern of antibiotic resistance in microbiological cultures of sepsis patients in a low-middle-income country's ICU. *Middle East J Anaesthesiol.* 2013;22:293–300.
19. Baljin B, Baldan G, Chimeddorj B, Tulgaa K, Gunchin B, Sandag T, et al. Faecal carriage of Gram-negative multidrug-resistant bacteria among patients hospitalized in two centres in Ulaanbaatar, Mongolia. *PLoS ONE.* 2016;11:e0168146. 10.1371/journal.pone.0168146.
20. Centers for Disease Control and Prevention. NHSN patient safety component manual, ch. 2: identifying HAIs. Atlanta: CDC; 2024.
21. Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing. Volume 100, 35th ed. Wayne, PA: CLSI; 2025.
22. Magiorakos AP, Srinivasan A, Carey RB, et al. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect.* 2012;18:268–81.
23. World Health Organization. WHO bacterial priority pathogens list 2024. Geneva: WHO; 2024.
24. Montoya A, et al. Rates of multidrug-resistant organism contamination on healthcare workers' hands: a systematic review and meta-analysis. *Am J Infect Control.* 2019;47:693–703.
25. Boyce JM, Pittet D. Guideline for hand hygiene in health-care settings. *MMWR Recomm Rep.* 2002;51(RR–16):1–45.
26. World Health Organization. WHO guidelines on hand hygiene in health care. Geneva: WHO; 2009.
27. Yokoe DS, et al. A compendium of strategies to prevent healthcare-associated infections in acute-care hospitals: 2022 updates. *Infect Control Hosp Epidemiol.* 2023;44:1–18.
28. Baur D, Gladstone BP, Burkert F, et al. Effect of antibiotic stewardship on the incidence of infection and colonisation with antibiotic-resistant bacteria: a systematic review and meta-analysis. *Lancet Infect Dis.* 2017;17:990–1001.

## Figures




**Figure 1**

Distribution of MDRO-positive HAI cases across service categories. Four service categories accounted for 91.6% of all 1,819 MDRO-positive HAI cases; General Surgery and Anaesthesiology-ICU alone contributed 63.8%. ‘Other wards’ pools all services other than General Surgery, intensive care, and emergency.




**Figure 2**

Monthly phenotype-confirmed MDRO incidence density and temporal-model sensitivity. (A) Monthly IPC-classified MDRO-positive HAI records per 1,000 patient-days. Filled points denote data-complete months; open points denote six data-incomplete 2023 months. February–June 2025 are shown after recovery from a secondary extract. The solid line is the complete-month negative-binomial trend; the dashed line is the full-period trend. (B) IRRs per year across the three model specifications. The full-period estimate is driven by incomplete surveillance capture.




**Figure 3**

Environmental surface culture and MDRO positivity by unit. Sampling was conducted from 16 June to 3 July 2025. All nine field-negative controls showed no growth. Overall, 44/131 (33.6%) surfaces were culture-positive and 22/131 (16.8%) were MDRO-positive. The Stroke Unit had the highest surface MDRO positivity despite a lower clinical-burden category.




**Figure 4**

Exploratory unit-level IPC prioritisation matrix. The matrix juxtaposes clinical-burden categories, environmental-surface MDRO positivity, and MRSA positivity in HCW hand imprints. Bars are scaled to 0–50%. Clinical categories derive from the interim surveillance review; final retrospective reporting aggregated General Surgery. Tiers indicate operational priorities and do not imply transmission or molecular linkage.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile1FigureS1ARMORMNresistancelandscape.pdf](#)
- [Additionalfile2TableS1ARMORMNheatmapdata.csv](#)