

Spontaneous Excitonic Mass Generation in Graphene Monolayers

Bent Weber

b.weber@ntu.edu.sg

Nanyang Technological University <https://orcid.org/0000-0001-8586-127X>

Yande Que

Nanyang Technological University <https://orcid.org/0000-0002-5267-4985>

Amit Kumar

Nanyang Technological University

Harley Scammell

University of Technology Sydney <https://orcid.org/0000-0001-8969-6064>

Zheng Tong

Nanyang Technological University

Zihao Huang

Nanyang Technological University <https://orcid.org/0009-0005-1434-7851>

Gagandeep Singh

Nanyang Technological University

Michael Fuhrer

Monash University <https://orcid.org/0000-0001-6183-2773>

Oleg Sushkov

University of New South Wales <https://orcid.org/0000-0001-7828-5569>

Physical Sciences - Article

Keywords:

Posted Date: October 22nd, 2026

DOI: <https://doi.org/10.21203/rs.3.rs-11022308/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Additional Declarations: There is **NO** Competing Interest.

Spontaneous Excitonic Mass Generation in Graphene Monolayers

Yande Que,^{1,†} Amit Kumar,^{1,†} Harley D. Scammell,² Zhengjue Tong,¹ Zihao Huang,¹

Gagandeep Singh,¹ Michael S. Fuhrer,³ Oleg P. Sushkov,⁴ and Bent Weber^{1,4,*}

¹*School of Physical and Mathematical Sciences, Nanyang Technological University, Singapore 637371, Singapore*

²*School of Mathematical and Physical Sciences, University of Technology Sydney, Ultimo NSW 2007, Australia*

³*School of Physics and Astronomy, Monash University, Clayton, Victoria 3800, Australia*

⁴*School of Physics, University of New South Wales, Sydney, NSW 2052, Australia*

(Dated: September 13, 2026)

Coulomb interactions among graphene’s massless Dirac fermions are unusually strong, making graphene a unique testbed for correlated quantum phases and interacting field theories. Among the predicted phases is an excitonic insulator, in which strong interactions drive spontaneous electron-hole pairing and condensation into a superfluid ground state. This transition has evaded experimental observation because unscreened long-range Coulomb interactions renormalize and enhance the Dirac velocity, keeping graphene in a weakly coupled semimetallic regime. Here, we overcome this constraint by decorating graphene with nanoscale metals that screen long-range interactions and suppress velocity renormalization. Scanning tunnelling spectroscopy down to 150 mK reveals a hard spectral gap ($2\Delta \approx 14$ meV) that emerges strictly at the Dirac point and is strongly susceptible to doping, temperature, and magnetic field — signatures of a many-body rather than single-particle gap. The gap closes at an anomalously low critical temperature ($\Delta/k_B T_c \geq 100$), consistent with a Berezinskii–Kosterlitz–Thouless transition, and coincides with a charge-density wave at $\vec{q} = \vec{K}' - \vec{K}$ that reflects intervalley excitonic pairing. Together, these results provide strong evidence for spontaneous excitonic mass generation in graphene, a correlated insulating phase long predicted but never before observed.

Graphene has long been viewed as a unique platform to explore interaction-driven quantum phenomena in two dimensions [1, 2]. Its low-energy electronic structure consists of massless Dirac fermions whose linear dispersion and vanishing density of states at the Dirac point amplify the role of Coulomb interactions [3–5]. The strength of the long-range Coulomb interaction can be characterized by an effective dimensionless coupling constant [6],

$$\alpha = \frac{e^2}{4\pi\epsilon_r\epsilon_0\hbar v}, \quad (1)$$

that, analogous to the fine-structure constant $\alpha = 1/137$, makes graphene a condensed-matter analogue of relativistic quantum electrodynamics (QED) [7, 8].

In contrast to the fine-structure constant, the effective interaction strength in graphene is of order unity—enhanced by a Dirac velocity $v \approx 1 \times 10^6$ m/s, 300 times slower than the speed of light, and tunable through the dielectric constant (ϵ_r) of its environment. A strong-coupling regime can be defined for $\alpha_c = 0.9 \sim 1.6$ [8–10], for which theoretical models [11–15] predict that Coulomb interactions give rise to stable electron-hole pairs (excitons). These excitons can spontaneously condense, driving a transition from a semimetal to an excitonic superfluid ground state, exhibiting a spectral mass gap and broken translational symmetry [16–19] (Figure 1).

However, despite two decades of theoretical work, direct experimental evidence for this interaction-driven instability has remained elusive, which is usually attributed to long-range Coulomb interactions renormalising the band dispersion and

enhancing the Dirac velocity near the Dirac point (Figure 1b) [5, 20, 21]. As observed in suspended graphene [22], such enhanced velocity effectively reduces the coupling constant as per Eq. (1) and retains the system in a weakly coupled semimetallic state, in which excitonic pairing is suppressed.

Metallic screening can suppress the interaction-driven velocity enhancement [23, 24], thereby providing an alternative pathway to increasing the effective coupling constant. Indeed, recent experiments on proximity-screened graphene devices have demonstrated that the low-energy electronic structures of graphene can be dramatically altered [25–27], enabling ultrahigh carrier mobility and unconventional quantum Hall responses. However, no evidence for exciton formation and condensation has been reported to date [28].

Here, we follow a different approach. Decorating the graphene surface with nanometre-scale metals as local screening centers, we achieve a two-fold reduction of the Dirac velocity near the Dirac point, enabling access to the strong coupling regime, where $\alpha \simeq 2 > \alpha_c$ [8–10]. Scanning tunnelling microscopy and spectroscopy (STM/STS) down to 150 mK [30] confirms the opening of a spectral gap—as large as $2\Delta = 14$ meV—only at charge neutrality in decorated samples, and strongly suppressed by temperature, carrier density, and magnetic field. In agreement with theoretical predictions, we interpret our findings by the spontaneous formation of excitons and their condensation into a superfluid ground state, described by a Berezinskii–Kosterlitz–Thouless (BKT)-type phase transition. Atomic-resolution STM reveals a charge-density-wave (CDW) order that strictly coincides with the gapped state and melts upon field-effect doping. The period of the observed modulations corresponds well with the wavevector ($\vec{q} = \vec{K}' - \vec{K}$) connecting the two inequivalent Dirac valleys—expected from intervalley excitonic pairing, and providing strong evidence for the presence of an exciton condensate. Our observations thus confirm that controlling

[†] These authors contributed equally to this work.

* b.weber@ntu.edu.sg




Fig. 1. An excitonic condensate in graphene. **a**, Schematic of a graphene-based field-effect device for STM. A gate voltage is applied to the silicon back-gate to tune the carrier density. **b**, Schematic of pristine graphene exhibiting a gapless Dirac spectrum, accompanied by an atomic-resolution STM image and a large-scale topographic image (150 nm $\times$ 150 nm, inset). **c**, Schematic of metal-decorated graphene in the excitonic-condensate state, in which an interaction-induced gap opens at charge neutrality, accompanied by the corresponding atomic-resolution STM image of a clean graphene region in between metals and large-scale topography (150 nm $\times$ 150 nm, inset). **d**, Differential tunnelling conductance (dI/dV) spectra of graphene measured at gate voltages ranging from -60 V to +60 V (curves vertically offset for clarity). The Dirac point energy with respect to the Fermi level (black arrow) shifts in energy with doping as indicated by the dashed red curve. The inset shows an expanded view of the spectra in the n -doped regime, highlighting the systematic shift of the Dirac point with gate voltage. **e**, Dirac point energy E_D , extracted from **d**, as a function of gate voltage, where $|E_D| = e|V_D^*| - \hbar\omega$, with $\hbar\omega \sim 65$ meV corresponds to an inelastic tunnelling suppression [29]. The solid red line shows a fit to $|E_D| = \hbar v \sqrt{\pi\gamma|V_G - V_0|}$ (See Supplementary Information for detail), to extract the Dirac velocity v and charge neutral point at $V_G = V_0$. **f**, High-resolution dI/dV spectra acquired at charge neutrality point, and at different temperatures, exhibiting a hard spectral gap emerging at E_F below $T < 0.5$ K, fingerprint of the exciton condensate.

the dielectric environment can drive graphene into a regime of strong electronic correlations where the excitonic instability becomes energetically favorable.

Figure 1a shows a schematic of the graphene monolayer field-effect devices used in the experiment. A monolayer graphene crystal was exfoliated onto a $\text{SiO}_2/\text{Si}^{++}$ substrate, through which a global back-gate allows for continuous tuning of the charge carrier density from hole (p -type) to electron (n -type) doping. This is confirmed by the gate-voltage-dependent tunnelling spectra in Figures 1d, and 1e allowing us to precisely pinpoint the charge-neutrality condition, where the Fermi level coincides with the Dirac point ($E_D = E_F$). Fits to the Dirac point energy (Figure 1e) as a function of doping (see Supplementary Information for detail) allow for

an extraction of the Dirac velocity in pristine graphene ($v \simeq 1 \times 10^6$ m/s), in excellent agreement with previous reports [29] and theoretical estimates [1]. Nanoscale metals were subsequently deposited onto the graphene top surface including both tungsten (W) and metallic tungsten ditelluride (WTe_2) [31, 32]. In both cases, clusters/islands (Figure 1c, inset) are separated by areas of clean high-quality graphene, as confirmed by the atomic-resolution STM topographic images in Figure 1c, taken after deposition.

Figure 1f shows the key observation of this study—the emergence of a large ($2\Delta \sim 14$ meV) spectral gap—centred at $E = E_F$, and only at charge neutrality ($E_F = E_D$) in metal-decorated graphene. Figures 2 and 3 confirm that this gap is absent in pristine graphene, and is strongly suppressed




Fig. 2. Charge doping dependence of the mass gap. **a, d**, Dirac point energy as a function of gate voltage in pristine (**a**) and W-decorated (**d**) graphene. The solid red lines show fits to extract the Dirac velocity in pristine ($v \approx 1.0 \times 10^6$ m/s) and decorated (0.5×10^6 m/s), respectively (data in **a** same as in Figure 1e). **b, e**, dI/dV intensity maps as functions of gate voltage and energy for pristine (**b**) and W-decorated (**e**) graphene. A mass gap is only observed after decoration. **c, f**, Individual point spectra at different doping extracted from the corresponding maps in (**b**), (**e**). **g**, Dirac velocity in monolayer graphene extracted from fits to gate-dependent tunnelling spectroscopy (open symbols) and Landau level spectroscopy (filled symbols). Triangles and squares denote WTe_2 -decorated and W-decorated graphene samples, respectively (see Figures S3 and S4 of Supplementary Information). **h**, Corresponding coupling constant α calculated from Eq. (1). Graphene is semimetallic for $\alpha < 1.1$, but develops a spectral gap for $\alpha > 1.1$, as resolved by high-resolution spectroscopy in (**e, f**) and Figure S5 in the Supplementary Information.

by field-effect doping (Figure 2), temperature, and magnetic field (Figure 3). Notably, we find that the gap narrows and disappears at a critical temperature of only $T_c \simeq 0.5$ K, giving rise to an exceptionally large gap ratio ($\Delta/k_B T_c \geq 100$)—two orders of magnitude beyond the BCS-like weak coupling limit. Above T_c , the hard gap closes, leaving only a weak V-shaped suppression at E_F , likely a Coulomb gap arising from long-range disorder [31, 33]. The latter persists over a wide range of carrier densities, temperatures, and magnetic fields, and remains largely insensitive to them (see Figure S9 in Supplementary Information).

The emergence of a spectral gap centred at the Fermi level and at charge neutrality suggests the formation of a correlated electronic state. In principle, such a gap could arise from several physical mechanisms, such as exciton condensation [28], a Mott insulating phase [34], or superconductivity [35]. Although the temperature and magnetic-field evolution of the gap resemble that of a superconductor, the observed gapped phase emerges only at charge neutrality, where graphene's

density of states vanishes. This behavior disfavors a superconducting origin, as the gap is tied specifically to charge neutrality rather than to a finite-size Fermi surface. Likewise, the magnetic-field dependence contrasts with that expected for a Mott insulator, whose gap is primarily governed by on-site Coulomb interactions and is generally insensitive to moderate magnetic fields [36].

Instead, the strong sensitivity to carrier density, temperature, and magnetic field is consistent with an excitonic instability. The gap emerges only near charge neutrality, where electron and hole excitations coincide, and is observed to rapidly collapse upon charge doping (Figures 2e and 2f). This is expected when the electron-hole balance is disrupted away from the Dirac point. Increasing temperature similarly destabilises the condensate. Such pronounced susceptibility to doping and temperature is a characteristic feature of excitonic condensation as reported in other candidate systems [28, 31, 37, 38].

An excitonic instability in graphene demands enhanced




Fig. 3. Temperature and magnetic-field stability of the excitonic state. **a**, dI/dV intensity map as a function of energy and temperature. **b**, Spectral gap size as a function of temperature extracted from the data in **(a)**. The solid and dashed lines show simulations based on a 2D XY model of lattice size $N = 16$. **c**, **e**, dI/dV intensity map and representative spectra, respectively, showing the evolution of the spectra with energy and out-of-plane magnetic field. Spectra are vertically offset for clarity. **d**, **f**, Magnetic-field dependence of the gap size extracted from **(c)** and **(e)**. The blue lines are fits to $\Delta(B) = \sqrt{\Delta_0^2 - \frac{1}{4}e\hbar v^2 B}$ (see Supplementary Information), yielding Dirac velocities $v \approx 0.8 \times 10^6$ m/s (WTe_2) and $v \approx 0.4 \times 10^6$ m/s (W) respectively, in excellent agreement with those measured from the gate voltage-dependent spectroscopy (Fig. 2g and Table I).

Coulomb interactions. Their strength is usually quantified by the coupling constant α expressed in Eq. (1) and depends on the Dirac velocity. Figures 2g and 2h plot the Dirac velocity and coupling constant, respectively, measured across different pristine and decorated samples. Dirac velocities were extracted from both gate-dependent conductance and magnetic-field dependent (Landau level) spectroscopic measurements (See Figure S4 in Supplementary Information). Only in the decorated samples, we find that the Dirac velocity is markedly reduced by up to a factor down to $1/2$ ($v \approx 0.5 \times 10^6$ m/s), compared to that of pristine graphene ($v \approx 1.0 \times 10^6$ m/s) [29, 39, 40]. It is also found to scale with metal coverage. We attribute this reduction to enhanced Coulomb screening induced by the metallic decoration, which suppresses the interaction-driven velocity renormalization [24] that ordinarily stabilises a weakly coupled semimetallic ground state. Indeed, as shown in Figure 2h, we observe that the spectral gap emerges only when the effective coupling constant exceeds $\alpha > 1.1$, in close agreement with theoretical estimates of the threshold ($\alpha_c \sim 0.9\text{--}1.6$) beyond which excitonic condensation is expected [8–10].

A striking signature of the excitonic insulating state is the unusually large gap ratio $\Delta/k_B T_c \geq 100$ (Figures 3a and 3b)—indicative of a strongly correlated pairing instability, consistent with the enhanced coupling constant for the decorated samples. In a 2D condensate, the transition would be

expected to be of the BKT type [41], satisfying $k_B T_{\text{BKT}} = \frac{\pi}{2} \rho(T_{\text{BKT}})$ [42], where $\rho(T_{\text{BKT}})$ is the condensate stiffness. As discussed in the Supplementary Information, within mean-field theory the stiffness scales linearly with the gap, $\rho(T) \approx \Delta(T)/16\pi$, so that the temperature dependence of the spectral gap follows directly from $\rho(T)$. Indeed, taking $\rho(T)$ from established Monte Carlo results for a finite size of $N \times N$ clusters in a 2D XY model [43], we find excellent agreement with the experimental data for an effective cluster size $N \approx 16$ (solid blue line in Figure 3b). This size agrees remarkably well with the estimate $N = R/l_s$ that follows from comparing the separation between metallic islands, $R \sim 10\text{--}30$ nm with the effective screening length $l_s \sim 1$ nm (See Supplementary Information for details).

Similar to the effect of temperature, Figures 3c and 3e illustrate how the spectral gap is suppressed by an out-of-plane magnetic field, and with a critical field that depends strongly on metal coverage. The suppression is also highly anisotropic, with an in-plane magnetic field only weakly affecting the gap (see Supplementary Information for details). We note that conventional theoretical studies have established that an out-of-plane magnetic field is generally expected to enhance excitonic instability through magnetic catalysis of long-range Coulomb interactions [6, 44, 45]. Our observations show the opposite trend. The key distinction is that in our samples, the long-range attraction is strongly screened by the metallic dec-




Fig. 4. **Intervalley excitonic order as reflected in gate-tunable charge density modulations.** **a-c**, STM topographic images of decorated graphene acquired at gate voltages of $V_G = -80$ V, 50 V, and 80 V. **d-f**, Corresponding FFT maps of the topographic images shown in **a-c**. **g**, Line profiles of the FFT intensity along the marked directions in **d-f**, showing the evolution of the relative intensity at the $\vec{K}/\vec{K}'$ points as a function of gate voltage. The FFT intensities were normalized to the Bragg peak intensity. **h**, Intensity ratio between the FFT peaks at $\vec{K}$ and $\vec{G}$ as a function of gate voltage, showing a sharp cusp at the onset of the gapped (insulating) regime. Dashed lines are guide to the eye.

oration, leaving only shorter wavelength components.

The pronounced magnetic-field anisotropy further points to an orbital origin of the suppression. In graphene, the in-plane electronic motion couples efficiently to an out-of-plane magnetic field through the orbital moment, whereas the orbital response to an in-plane field is strongly suppressed in strictly 2D electronic systems.

As our fits to the magnetic field dependence show (solid blue lines in Figures 3e and 3f), the observed gap suppression can be modelled *quantitatively* considering that an out-of-plane magnetic field suppresses excitonic pairing when the magnetic length becomes shorter than exciton radius ($l_B < a_B$) (see Supplementary Information for detail). In this case, the gap can be shown to follow $\Delta(B) = \sqrt{\Delta_0^2 - \frac{1}{4}e\hbar v^2 B}$ depending on the Dirac velocity v and the gap magnitude $\Delta_0(B = 0)$ as the only free parameters. Despite a difference with observed critical magnetic field by a factor of nearly

6, our fits (solid blue lines in Figures 3e and 3f) reproduce the measured suppression well in both data. Indeed, the extracted Dirac velocities $v = 0.42 \pm 0.02 \times 10^6$ m/s (W) and $v = 0.77 \pm 0.08 \times 10^6$ m/s (WTe₂), setting the scale of the critical field, are in excellent agreement with those measured in gate-voltage dependent spectroscopy respectively, as summarised in Table I, demonstrating remarkable internal consistency of the model.

A hallmark of many correlated electronic phases is the emergence of translation-symmetry-breaking charge orders. For an excitonic condensate, we would expect pairing of electrons and holes at finite momentum across graphene's inequivalent $\vec{K}$ and $\vec{K}'$ valleys [46], giving rise to a CDW modulation with wavevector $\vec{q} = \vec{K}' - \vec{K}$. Figures 4a-c demonstrate the presence of the expected charge modulations—only in the gapped phases near charge neutrality—and further confirmed by corresponding Fourier components (Figures 4d-f). The

TABLE I. **Measured Dirac velocity across different samples.** Dirac velocity v for pristine, WTe₂- and W-decorated graphene, extracted independently from gate-dependent STS (Gate), Landau-level spectroscopy (LLs), and fits to the magnetic-field dependence of the spectral gap size (Mag.). The close agreement across values obtained using different measurements provides a consistent determination of the Dirac velocity. All velocity values are given in units of 10^6 m/s.

Sample	Coverage (%)	Gate	LLs	Mag.
Pristine-G	0	1.04 ± 0.02	—	—
WTe ₂ -Gr	18	0.93 ± 0.05	—	—
	75	0.78 ± 0.01	0.53 ± 0.03	0.77 ± 0.08
	85	0.71 ± 0.09	—	—
W-Gr	37	0.46 ± 0.01	—	0.42 ± 0.02

amplitude of these components is rapidly suppressed upon carrier doping (Figures 4g and 4h), coinciding with the appearance of the spectral gap. No such charge order was observed in pristine graphene, indicating that the charge modulations are intrinsically linked to the gapped state.

A minimal symmetry-based description of the intervalley excitonic state follows directly from the low-energy Dirac theory (see Supplementary Information). Starting from $H_0 = v(k_x\sigma_x\tau_z + k_y\sigma_y)$, the simplest mass term that (i) gaps the spectrum, (ii) preserves lattice symmetries, and (iii) carries momentum $\vec{q} = \vec{K}' - \vec{K}$ is $\delta H = \Delta\sigma_x\tau_x$. This term encodes intervalley coherence and naturally generates a charge density modulation at $\vec{q}$, exactly as observed in experiments.

While other mechanisms such as intervalley scattering or Kekulé order [47, 48] can in principle give rise to similar modulations, these would be expected to persist over a wide range of carrier density. In stark contrast, here doping acts to destabilise the condensate once the chemical potential exceeds the pairing scale and the modulation subsides. The simplest mean-field treatment would lead us to expect an energy scale $\mu_c \sim \Delta_0$ for this transition. Experimentally, we observe stability up to $|\mu| \approx 70$ meV $\sim 10\Delta_0$ (see Figures 2e and 2f), substantially larger than this bare expectation. As detailed in the Supplementary Information, this can be reconciled as a phonon mode at the K point ($\Omega_0 \approx 150$ meV) couples to the identical $\sigma_x\tau_x$ structure, thereby reinforcing the same intervalley excitonic channel. For $2|\mu| \lesssim \Omega_0$, this interaction provides an additional attractive contribution, leading to a natural scale $\mu_c \sim \Omega_0/2 \approx 85$ meV, consistent with the experimental observations. So, while the phonon does not constitute the primary pairing mechanism, it enhances and stabilises the intervalley excitonic condensate under doping.

Conclusion

In summary, we have presented strong evidence for an excitonic condensate in graphene stabilised by enhanced Coulomb interactions. Our tunnelling spectroscopy reveals a hard spectral gap with a remarkably large gap-to-critical-temperature ratio $\Delta/k_B T_c \geq 100$, gate-tunable density-wave order at $\vec{q} = \vec{K}' - \vec{K}$, and strong susceptibility to carrier doping and magnetic fields. These findings collectively provide compelling evidence for spontaneous excitonic mass generation

in graphene, long sought in theory but elusive in experimental demonstrations.

Methods

Device fabrication. Graphene monolayers were mechanically exfoliated onto a SiO₂(300 nm)/Si substrate using Scotch tape and identified by optical contrast, Raman spectroscopy, and atomic force microscopy (Figure S1 in the Supplementary Information). The freshly exfoliated graphene was annealed in ultra-high vacuum (UHV) at 400 °C with a slow ramp rate of ~ 2 °C/min to avoid cracking. Electrical contacts to the graphene were fabricated by indium micro-soldering [49], while the heavily doped Si substrate served as the back-gate electrode.

MBE deposition. Both WTe₂ monolayer islands and W clusters were deposited in UHV (below 1×10^{-10} mbar) by molecular-beam epitaxy (MBE). Prior to deposition, the substrates were degassed in UHV at 180 °C for 30 min. WTe₂ islands were grown by co-deposition of high-purity W (99.998%) and Te (99.999%) with a flux ratio of 1:280 with the substrate held at 160 °C for 1–3 h, yielding 20–80% monolayer coverage, consistent with previous reports [31, 32, 50, 51]. W clusters were obtained by direct deposition of W (99.998%) onto graphene held at room temperature. After deposition, all samples were directly transferred to the STM stage under UHV without exposure to air, ensuring pristine measurement conditions.

Scanning tunnelling microscopy/spectroscopy. Scanning tunnelling microscopy and spectroscopy (STM/STS) were performed in a customized ultra-low temperature STM (USM1600, Unisoku) operating under UHV conditions ($< 1 \times 10^{-10}$ mbar), with an effective electronic temperature below 150 mK [30] at a base temperature of 40 mK. The effective junction temperature was calibrated by analyzing the spectral broadening of the layered superconductor 1T-PdTe₂ (see Section SVIII of the Supplementary Information). Chemically etched PtIr tips were calibrated against the Shockley surface state of Au(111) ahead of spectroscopic measurement. Unless otherwise specified, STS measurement was performed at base temperature using a standard lock-in technique with a modulation amplitude of $V_{ac} = 5$ mV for large-energy-scale spectra, and 0.1 mV for high-resolution spectra, both at a frequency of 726 Hz.

Acknowledgment

This work was supported by the Singapore Ministry of Education (MOE) Academic Research Fund Tier 3 grant (MOE-MOET32023-0003) Quantum Geometric Advantage and a Singapore National Research Foundation (NRF) Grant (No. NRF-F-CRP-2024-0012). MSF, OPS, and BW acknowledge support from the FLEET Centre of Excellence, ARC Grant No. CE170100039. We acknowledge very useful discussions with Jürgen Smet and Alexander R. Hamilton. BW acknowledges a Visiting Fellowship at the University of New South Wales (UNSW).

Author Contributions. YQ performed the STM/STS ex-

periments with assistance from AK, ZT, ZH, and GS. AK and YQ prepared the samples. HDS and OPS developed the theoretical model. YQ, BW, HDS, OPS and MSF analysed the data. YQ wrote the manuscript with the help of BW. BW conceived, supervised, and coordinated the project. All authors discussed and provided input and edits to the manuscript.

Correspondence and requests for materials should be addressed to b.weber@ntu.edu.sg

Competing financial interests

The authors declare no competing financial interests.

- [1] A. H. Castro Neto, F. Guinea, N. M. R. Peres, K. S. Novoselov, and A. K. Geim, The electronic properties of graphene, *Reviews of Modern Physics* **81**, 109 (2009).
- [2] V. N. Kotov, B. Uchoa, V. M. Pereira, F. Guinea, and A. H. Castro Neto, Electron-electron interactions in graphene: Current status and perspectives, *Reviews of Modern Physics* **84**, 1067 (2012).
- [3] I. F. Herbut, Interactions and phase transitions on graphene honeycomb lattice, *Physical Review Letters* **97**, 146401 (2006).
- [4] I. F. Herbut, V. Juričić, and B. Roy, Theory of interacting electrons on the honeycomb lattice, *Physical Review B* **79**, 085116 (2009).
- [5] H.-K. Tang, J. Leaw, J. Rodrigues, I. Herbut, P. Sengupta, F. Asaad, and S. Adam, The role of electron-electron interactions in two-dimensional Dirac fermions, *Science* **361**, 570 (2018).
- [6] D. V. Khveshchenko, Ghost excitonic insulator transition in layered graphite, *Physical Review Letters* **87**, 246802 (2001).
- [7] J. E. Drut and T. A. Lähde, Critical exponents of the semimetal-insulator transition in graphene: A Monte Carlo study, *Physical Review B* **79**, 241405 (2009).
- [8] J. E. Drut and T. A. Lähde, Is graphene in vacuum an insulator?, *Physical Review Letters* **102**, 026802 (2009).
- [9] O. Gamayun, E. Gorbar, and V. Gusynin, Supercritical coulomb center and excitonic instability in graphene, *Physical Review B* **80**, 165429 (2009).
- [10] O. Gamayun, E. Gorbar, and V. Gusynin, Gap generation and semimetal-insulator phase transition in graphene, *Physical Review B* **81**, 075429 (2010).
- [11] D. Son, Quantum critical point in graphene approached in the limit of infinitely strong coulomb interaction, *Physical Review B* **75**, 235423 (2007).
- [12] D. Khveshchenko, Massive Dirac fermions in single-layer graphene, *Journal of Physics: Condensed Matter* **21**, 075303 (2009).
- [13] G.-Z. Liu, W. Li, and G. Cheng, Interaction and excitonic insulating transition in graphene, *Physical Review B* **79**, 205429 (2009).
- [14] J. E. Drut and T. A. Lähde, Lattice field theory simulations of graphene, *Physical Review B* **79**, 165425 (2009).
- [15] J.-R. Wang and G.-Z. Liu, Eliashberg theory of excitonic insulating transition in graphene, *Journal of Physics: Condensed Matter* **23**, 155602 (2011).
- [16] N. F. Mott, The transition to the metallic state, *Philosophical Magazine* **6**, 287 (1961).
- [17] D. Jérôme, T. M. Rice, and W. Kohn, Excitonic insulator, *Physical Review* **158**, 462 (1967).
- [18] B. I. Halperin and T. M. Rice, Possible anomalies at a semimetal-semiconductor transition, *Reviews of Modern Physics* **40**, 755 (1968).
- [19] W. Kohn and D. Sherrington, Two kinds of bosons and bose condensates, *Reviews of Modern Physics* **42**, 1 (1970).
- [20] C. Hwang, D. A. Siegel, S.-K. Mo, W. Regan, A. Ismach, Y. Zhang, A. Zettl, and A. Lanzara, Fermi velocity engineering in graphene by substrate modification, *Scientific Reports* **2**, 590 (2012).
- [21] T. Stauber, P. Parida, M. Trushin, M. V. Ulybyshev, D. L. Boyda, and J. Schliemann, Interacting electrons in graphene: Fermi velocity renormalization and optical response, *Physical Review Letters* **118**, 266801 (2017).
- [22] D. C. Elias, R. V. Gorbachev, A. S. Mayorov, S. V. Morozov, A. A. Zhukov, P. Blake, L. A. Ponomarenko, I. V. Grigorieva, K. S. Novoselov, F. Guinea, and G. A. K., Dirac cones reshaped by interaction effects in suspended graphene, *Nature Physics* **7**, 701 (2011).
- [23] A. Raoux, M. Polini, R. Asgari, A. R. Hamilton, R. Fazio, and A. H. MacDonald, Velocity-modulation control of electron-wave propagation in graphene, *Physical Review B* **81**, 073407 (2010).
- [24] D. A. Siegel, C.-H. Park, C. Hwang, J. Deslippe, A. V. Fedorov, S. G. Louie, and A. Lanzara, Many-body interactions in quasi-freestanding graphene, *Proceedings of the National Academy of Sciences* **108**, 11365 (2011).
- [25] M. Kim, S. Xu, A. Berdyugin, A. Principi, S. Slizovskiy, N. Xin, P. Kumaravel, W. Kuang, M. Hamer, R. Krishna Kumar, R. V. Gorbachev, K. Watanabe, T. Taniguchi, I. V. Grigorieva, V. I. Fal'ko, M. Polini, and A. K. Geim, Control of electron-electron interaction in graphene by proximity screening, *Nature Communications* **11**, 2339 (2020).
- [26] D. Domaretskiy, Z. Wu, V. H. Nguyen, N. Hayward, I. Babich, X. Li, E. Nguyen, J. Barrier, K. Indykiewicz, W. Wang, R. V. Gorbachev, N. Xin, K. Watanabe, T. Taniguchi, L. Hague, V. I. Fal'ko, I. V. Grigorieva, L. A. Ponomarenko, A. I. Berdyugin, and A. K. Geim, Proximity screening greatly enhances electronic quality of graphene, *Nature* **644**, 646 (2025).
- [27] A. S. Mayorov, P. Wang, X. Yue, B. Wu, J. He, D. Zhang, F. Lian, S. Jiang, J. Huang, Z. Wang, G. Qian, W. Kenji, T. Takashi, D. Renjun, W. Rui, W. Beigeng, W. Lei, N. K. S., and Y. Geliang, Quantum Hall effect at 0.002 T in graphene,

- Nature Communications* **17**, 2003 (2026).
- [28] Y. Que, C. Rebanal, L. Watson, M. Fuhrer, M. Papaj, B. Weber, and I. Di Bernardo, Excitonic order in quantum materials: fingerprints, platforms and opportunities, arXiv [10.48550/arXiv.2603.24211](https://arxiv.org/abs/10.48550/arXiv.2603.24211) (2026).
- [29] Y. Zhang, V. W. Brar, F. Wang, C. Girit, Y. Yayon, M. Panlasigui, A. Zettl, and M. F. Crommie, Giant phonon-induced conductance in scanning tunnelling spectroscopy of gate-tunable graphene, *Nature Physics* **4**, 627 (2008).
- [30] Y. Que, A. Kumar, M. S. Lodge, Z. Tong, M. Lai Kar Fai, W. Tao, Z. Cui, R. Shivajirao, J. Jia, S. E. Lee, and B. Weber, Performance benchmarking of an ultra-low vibration laboratory to host a commercial millikelvin scanning tunnelling microscope, *Nanotechnology* **34**, 455704 (2023).
- [31] Y. Que, Y.-H. Chan, J. Jia, A. Das, Z. Tong, Y.-T. Chang, Z. Cui, A. Kumar, G. Singh, S. Mukherjee, H. Lin, and B. Weber, A gate-tunable ambipolar quantum phase transition in a topological excitonic insulator, *Advanced Materials* **36**, 2309356 (2024).
- [32] L. Watson, J. Ripoll, Z. Tong, A. Kumar, Y. Que, Y.-H. Chan, H. Lin, S. Mukherjee, M. Garnica, M. T. Edmonds, P. Michal, M. Papaj, A. L. Vazquez de Parga, B. Weber, I. Di Bernardo, and M. S. Fuhrer, Observation of the charge density wave excitonic order parameter in topological insulator monolayer WTe_2 , *ACS Nano* **19**, 32374–32381 (2025).
- [33] Y. H. Song, Z. Y. Jia, D. Zhang, X. Y. Zhu, Z. Q. Shi, H. Wang, L. Zhu, Q. Q. Yuan, H. Zhang, D. Y. Xing, and S. C. Li, Observation of coulomb gap in the quantum spin Hall candidate single-layer $1\text{T}'\text{-WTe}_2$, *Nature Communications* **9**, 4071 (2018).
- [34] V. V. Deshpande, B. Chandra, R. Caldwell, D. S. Novikov, J. Hone, and M. Bockrath, Mott insulating state in ultraclean carbon nanotubes, *Science* **323**, 106 (2009).
- [35] T. T. Heikkilä, Surprising superconductivity of graphene, *Science* **375**, 719 (2022).
- [36] Y. H. Matsuda, D. Nakamura, A. Ikeda, S. Takeyama, Y. Suga, H. Nakahara, and Y. Muraoka, Magnetic-field-induced insulator–metal transition in W-doped VO_2 at 500 T, *Nature Communications* **11**, 3591 (2020).
- [37] Y. Jia, P. Wang, C. L. Chiu, Z. Song, G. Yu, B. Jäck, S. Lei, S. Klemen, F. A. Cevallos, M. Onyszczyk, N. Fishchenko, X. Liu, G. Farahi, F. Xie, Y. Xu, K. Watanabe, T. Taniguchi, B. A. Bernevig, R. J. Cava, L. M. Schoop, A. Yazdani, and S. Wu, Evidence for a monolayer excitonic insulator, *Nature Physics* **18**, 87 (2022).
- [38] Q. Gao, Y.-H. Chan, Y. Wang, H. Zhang, P. Jinxi, S. Cui, Y. Yang, Z. Liu, D. Shen, Z. Sun, J. Jiang, T. C. Chiang, and P. Chen, Evidence of high-temperature exciton condensation in a two-dimensional semimetal, *Nature Communications* **14**, 994 (2023).
- [39] Z. Jiang, E. A. Henriksen, L. C. Tung, Y.-J. Wang, M. E. Schwartz, M. Y. Han, P. Kim, and H. L. Stormer, Infrared spectroscopy of Landau levels of graphene, *Physical Review Letters* **98**, 197403 (2007).
- [40] A. Luican, G. Li, and E. Y. Andrei, Quantized Landau level spectrum and its density dependence in graphene, *Physical Review B* **83**, 041405 (2011).
- [41] J. M. Kosterlitz and D. J. Thouless, Ordering, metastability and phase transitions in two-dimensional systems, *Journal of Physics C: Solid State Physics* **6**, 1181 (1973).
- [42] D. R. Nelson and J. M. Kosterlitz, Universal jump in the superfluid density of two-dimensional superfluids, *Physical Review Letters* **39**, 1201 (1977).
- [43] Y.-D. Hsieh, Y.-J. Kao, and A. W. Sandvik, Finite-size scaling method for the Berezinskii–Kosterlitz–Thouless transition, *Journal of Statistical Mechanics: Theory and Experiment* **2013**, P09001 (2013).
- [44] E. V. Gorbar, V. P. Gusynin, V. A. Miransky, and I. A. Shovkovy, Magnetic field driven metal–insulator phase transition in planar systems, *Physical Review B* **66**, 045108 (2002).
- [45] V. P. Gusynin, V. A. Miransky, S. G. Sharapov, and I. A. Shovkovy, Excitonic gap, phase transition, and quantum Hall effect in graphene, *Physical Review B* **74**, 195429 (2006).
- [46] H. D. Scammell and O. P. Sushkov, Exciton condensation in biased bilayer graphene, *Physical Review Research* **5**, 043176 (2023).
- [47] C. Gutiérrez, C.-J. Kim, L. Brown, T. Schiros, D. Nordlund, E. B. Lochocki, K. M. Shen, J. Park, and A. N. Pasupathy, Imaging chiral symmetry breaking from Kekulé bond order in graphene, *Nature Physics* **12**, 950 (2016).
- [48] C. Bao, H. Zhang, X. Wu, S. Zhou, Q. Li, P. Yu, J. Li, W. Duan, and S. Zhou, Coexistence of extended flat band and Kekulé order in Li-intercalated graphene, *Physical Review B* **105**, L161106 (2022).
- [49] Ç. Ö. Girit and A. Zettl, Soldering to a single atomic layer, *Applied Physics Letters* **91**, 193512 (2007).
- [50] W. Tao, Z. J. Tong, A. Das, D.-Q. Ho, Y. Sato, M. Haze, J. Jia, Y. Que, F. Bussolotti, K. E. J. Goh, B. Wang, H. Lin, A. Bansil, S. Mukherjee, Y. Hasegawa, and B. Weber, Multiband superconductivity in strongly hybridized $1\text{T}'\text{-WTe}_2/\text{NbSe}_2$, *Physical Review B* **105**, 094512 (2022).
- [51] J. Jia, E. Marcellina, A. Das, M. S. Lodge, B. Wang, D.-Q. Ho, R. Biswas, T. A. Pham, W. Tao, C.-Y. Huang, H. Lin, A. Bansil, S. Mukherjee, and B. Weber, Tuning the many-body interactions in a helical Luttinger liquid, *Nature Communications* **13**, 6046 (2022).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [ElinGSI.pdf](#)