

Supplementary Information for

Myometrial contraction-activated endometrial PIEZO2 safeguards fertility under maternal stress

Jin-Wen Kang^{1,2†}, Yu Zhang^{1†}, Hui-Xia Li¹, He Zou¹, Jing-Yi Yang¹, Xiao-Qing Huang¹, Jia-Ying Yu¹,
Zi-Wei Ai³, Ying Zhang³, Rui Zhang¹, Ying-Nan Liu¹, Zeng-Ming Yang^{4*}, Bin Guo^{5*}, Hong-Lu Diao^{3*},
Ren-Wei Su^{1*}

Affiliations:

¹College of Veterinary Medicine, South China Agricultural University, Guangzhou, China

²Reproductive Sciences Center, Division of Developmental Biology, Cincinnati Children's Hospital Medical Center, Cincinnati, OH 45229, USA

³Reproductive Medicine Center of Renmin Hospital, Biomedical Engineering College, Hubei University of Medicine, Shiyan, Hubei, China

⁴College of Animal Science, Guizhou University, Guiyang, China

⁵College of Veterinary Medicine, Jilin University, Changchun, China

†These authors contributed equally to this work

***Correspondence**

Ren-Wei Su, 483 Wushan Road, Guangzhou, Guangdong, China, 510642

Email: renweisu@scau.edu.cn

Hong-Lu Diao, PhD.

39 Chaoyang Middle Road, Shiyan, Hubei, China, 442000

Email: hldiao@hbmh.edu.cn

27 Bin Guo, PhD.

28 5333 Xi'an Road, Changchun, Jilin, China, 130062

29 Email: guobin79@jlu.edu.cn

30 Zeng-Ming Yang, PhD.

31 2708 Huaxi District, Guiyang, Guizhou, China, 550025

32 Email: yangzm@gzu.edu.cn

33

34


35 **The file includes:**

36


37 Materials and Methods

38 Extended Data Figs. 1 to 7




39 Extended Data Tables 1 to 4




40 **Extended data Fig. 1. Generation and characterization of uterine-specific *Piezo2* conditional**
41 **knockout mice.** (A) *Piezo2* mRNA expression in non-pregnant uterus (NP-U), uterus at 3.5 dpc of
42 pregnancy (D4-U), and dorsal root ganglion (DRG), quantified by RT-qPCR and normalized to *Rpl19*, n
43 = 3 biological replicates. One-way ANOVA with Tukey's post hoc test. (B) Schematic of the conditional
44 *Piezo2* knockout strategy. *Piezo2^{ff}* mice carrying loxP sites flanking exons 43–45 were crossed with
45 *Pgr^{Cre/+}* transgenic mice to generate uterine-specific *Piezo2* conditional knockout (*Piezo2^{d/d}*) females. (C)
46 Immunohistochemistry detecting PIEZO2 in uterine cross-sections at 3.5 dpc from *Piezo2^{ff}* and *Piezo2^{d/d}*
47 mice, confirming the loss of PIEZO2 protein expression in the endometrial stroma of knockout mice. S,
48 stroma; LE, luminal epithelium. Scale bar, 100 μ m and 20 μ m. (D) Quantitative RT-PCR confirming
49 efficient *Piezo2* deletion in *Piezo2^{d/d}* uteri, with *Rpl19* serving as an internal control. Statistical
50 significance determined using a two-sided Student's t-test. (E) Western blot analysis of PIEZO2
51 expression in uterine tissues at 3.5 dpc in *Piezo2^{d/d}* and *Piezo2^{ff}* ($n = 5$ each) mice. Expression was
52 normalized to α -tubulin with the quantification shown to the right (two-sided Student's t-test). (F)
53 Representative uterine images at 4.5 dpc from *Piezo2^{ff}* and *Piezo2^{d/d}* females housed under quiet (45 ± 5
54 dB) or noisy (65 ± 5 dB) acoustic conditions. Red arrowheads indicate implantation sites visualized by
55 Chicago Sky Blue dye injection. Morphologically normal blastocysts were recovered from the uterine
56 lumen of *Piezo2^{d/d}* mice under noisy conditions (right panel). Scale bars, 5 mm and 10 μ m. (G)
57 Quantification of implantation site number per mouse under quiet and noisy conditions. *Piezo2^{d/d}* mice
58 displayed normal implantation under quiet conditions but complete implantation failure under noisy
59 conditions (two-way ANOVA with Tukey's post hoc test). (H) Schematic of the modified Single
60 Prolonged Stress (SPS) protocol. SPS consists of sequential restraint stress (1.5 h), forced swim (10 min),
61 and diethyl ether exposure until loss of consciousness, with this protocol administered daily between 0.5
62 and 3.5 dpc (9:00–11:00 AM). (I) Serum levels of corticosterone (Cort), epinephrine (EPI), and
63 noradrenaline (NA) in *Piezo2^{ff}* and *Piezo2^{d/d}* mice with or without SPS treatment, confirming successful
64 stress induction. SPS significantly elevated corticosterone in both genotypes ($n=5$, two-way ANOVA with
65 Tukey's post hoc test). (J) Dose–response analysis of stress sensitivity. Mice were subjected to various
66 intensities of SPS by adjusting the exposure durations. *Piezo2^{d/d}Piezo2-cKO* mice displayed implantation
67 failure at lower stress intensities than controls. Fractions indicate number of mice with implantation sites
68 out of the total. Quantitative data are presented as mean $\pm$ SEM. ns, $P > 0.05$; * $P < 0.05$; ** $P < 0.01$; *** P
69 < 0.001 .




70 **Extended data Fig. 2. Ovarian function in response to stress and differential regulation in**
71 ***Piezo2/Piezo1*.** (A) Representative images of blastocysts recovered at 3.5 dpc from *Piezo2^{ff/ff}* and *Piezo2^{d/d}*
72 mice with or without SPS treatment. Blastocyst morphology was comparable across all groups. Scale bar,
73 10 μ m. (B) Quantification of blastocyst number per mouse at 3.5 dpc across all experimental groups. No
74 significant differences were observed (n=5, two-way ANOVA with Tukey's post hoc test). (C) Serum
75 estradiol (E2) and progesterone (P4) levels at 3.5 dpc in *Piezo2^{ff/ff}* and *Piezo2^{d/d}* mice with or without SPS
76 treatment. No significant differences were observed between genotypes irrespective of stress conditions
77 (n = 3, two-way ANOVA with Tukey's post hoc test). (D) Single-molecule fluorescence *in situ*
78 hybridization (smFISH) of *Piezo2* mRNA in uterine sections from *Piezo2^{ff/ff}* and *Piezo2^{d/d}* mice with or
79 without SPS treatment. SPS selectively upregulated *Piezo2* expression in the endometrial stroma of
80 *Piezo2^{ff/ff}* mice. Arrows indicate *Piezo2*-positive cells. Scale bar, 100 μ m. (E) Quantitative RT-PCR
81 analysis of *Piezo2* (left), *Piezo2* exons 17–18 (middle, upstream of the floxed region), and *Piezo1* (right)
82 expression in *Piezo2^{ff/ff}* and *Piezo2^{d/d}* uteri with or without SPS treatment. SPS increased *Piezo2*
83 transcription only when functional PIEZO2 protein was present (left, middle panels). n=3 or 4. Statistical
84 significance was determined via two-way ANOVA with Tukey's post hoc test. (F) Representative uterine
85 images and quantification of implantation sites in *Piezo1^{ff/ff}* (control) and *Piezo2^{d/d}* (*Pgr^{Cre/+} Piezo1^{ff/ff}*) mice
86 with or without SPS treatment. Unlike *Piezo2*-cKO mice, *Piezo1*-cKO mice showed stable implantation
87 that was unaffected by SPS (two-way ANOVA with Tukey's post hoc test). Scale bar, 5 mm. Quantitative
88 data are presented as mean $\pm$ SEM. ns, $P > 0.05$; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

A**B****C**


89 **Extended data Fig. 3. Exogenous progesterone supplementation normalizes gene expression in SPS-**
90 **Piezo2-deficient mice. (A)** Schematic of the progesterone (P4) supplementation protocol. Exogenous
91 progesterone (P4) was administered subcutaneously to SPS-*Piezo2^{d/d}* mice at 2.5, 2.75, 3.5, and 3.75 dpc
92 during the peri-implantation period. The SPS protocol was followed daily from 0.5–3.5 dpc and
93 administered from 9:00–11:00 a.m. **(B)** Quantification of implantation site number (left) and implantation
94 site weight (right) at 4.5 dpc in SPS-*Piezo2^{ff}*, SPS-*Piezo2^{d/d}*, and SPS-*Piezo2^{d/d}+P4* mice ($n = 10$ per
95 group). **(C)** Quantitative RT-PCR analysis of estrogen-responsive genes (*Esr1*, *Muc1*, *Muc4*, *Ltf*) and
96 progesterone-responsive genes (*Pgr*, *Hoxa10*, *Hand2*, *Areg*) in non-SPS and SPS-treated *Piezo2^{ff}*,
97 *Piezo2^{d/d}*, and *Piezo2^{d/d}+P4* uteri at 3.5 dpc. Expression was normalized to *Rpl19* ($n = 4$ or 3 per group).
98 Quantitative data are presented as mean $\pm$ SEM. Statistical significance was determined using a two-way
99 ANOVA with Tukey's post hoc test. ns, $P > 0.05$; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.




100 **Extended data Fig. 4. NOS-mediated paracrine regulation of uterine contractility in PIEZO2-**
101 **deficient endometrium. (A)** Quantitative RT-PCR analysis of *nNos*, *iNos*, and *eNos* expression in
102 *Piezo2^{ff}* and *Piezo2^{d/d}* uteri with or without SPS treatment. Expression was normalized to *Rpl19* ($n = 3$
103 per group), with NOS isoforms significantly upregulated in *Piezo2^{d/d}* endometrium. **(B)** Nitric oxide (NO)
104 levels in endometrial tissue from *Piezo2^{ff}* and *Piezo2^{d/d}* mice treated with vehicle or NOS inhibitor (L-
105 NAME HCl). Injections were administered daily from 0.5 to 3.5 dpc, with NOS levels assessed on 3.5
106 dpc ($n = 3$ per group). **(C and D)** Representative *ex vivo* tension traces from *Piezo2^{ff}* (top) and *Piezo2^{d/d}*
107 (bottom) uteri before and after L-NAME HCl treatment **(C)**, and quantification of contraction frequency
108 and amplitude following L-NAME HCl treatment **(D)** collected at 3.5 dpc ($n = 5$ per group). **(E)**
109 Representative uterine images at 4.5 dpc from *Piezo2^{ff}* and *Piezo2^{d/d}* mice treated with L-NAME HCl
110 daily for 4 days. Implantation sites were visualized by intravenous injection of Chicago Sky Blue dye
111 (blue spots) and indicated with red arrowheads. Scale bar, 5 mm. **(F, G)** Histological analysis (H&E
112 staining) **(F)** and immunohistochemistry detecting COX2 **(G)** in uterine sections at 4.5 dpc following L-
113 NAME HCl treatment. Scale bar, 100 μ m. Quantitative data are presented as mean $\pm$ SEM. Statistical
114 significance was determined using a two-way ANOVA with Tukey's post hoc test. ns, $P > 0.05$; * $P <$
115 0.05; ** $P < 0.01$; *** $P < 0.001$.



116 **Extended data Fig. 5. scRNA-seq-based characterization of uterine cell populations and stromal**
117 **paracrine signaling.** (A) UMAP visualization of cell clusters identified by scRNA-seq in SPS-*Piezo2^{ff/ff}*
118 (10,678 cells) and SPS-*Piezo2^{d/d}* (8,199 cells) uteri at 3.5 dpc. Each point represents a single cell and is
119 colored according to its assigned cell cluster ($n = 14$). (B) Stacked bar plot showing the distribution of
120 each cell type identified by scRNA-seq in SPS-*Piezo2^{ff/ff}* versus SPS-*Piezo2^{d/d}* uteri. Stromal fibroblasts
121 were the largest population in both genotypes but were reduced in PIEZO2-deficient uteri relative to
122 controls (62.94% vs. 55.56%). (C) UMAP visualization of uterine stromal cell (uSC) subclusters after
123 fibroblast sub-clustering. Dashed circle indicates the uterine stromal cells. (D) Violin plots showing
124 collagen and extracellular matrix-associated gene expression in mesenchymal fibroblast in subcluster 2
125 from SPS-*Piezo2^{ff/ff}* (Ctrl, blue) and SPS-*Piezo2^{d/d}* (KO, pink) uteri. Statistical significance was determined
126 by Wilcoxon rank-sum test with Bonferroni correction. (E) Violin plots showing *Piezo2* expression levels
127 across the six fibroblast subclusters. (F) Chord diagram of predicted ligand–receptor interactions among
128 fibroblast subclusters generated with CellPhoneDB. Node size represents the relative abundance of
129 predicted interactions for each subcluster, and line thickness indicates the relative interaction strength. (G)
130 Dot plots showing the top five most significant ligand–receptor pairs for each PIEZO2-expressing
131 fibroblast subcluster compared to all other subclusters. Dot size represents the $-\log_{10}(P \text{ value})$, while the
132 color gradient represents the communication probability ($\log_2$ mean expression of the ligand–receptor
133 pair). Blue shading highlights targeted paracrine signaling directed toward the ECM-producing subcluster
134 2. (H) Quantitative RT-PCR analysis of *Igfl* expression in *Piezo2^{ff/ff}* and *Piezo2^{d/d}* uteri with or without
135 SPS treatment. Expression was normalized to *Rpl19* ($n = 3$ per group, two-way ANOVA with Tukey's
136 post hoc test). (I) Quantitative RT-PCR analysis of collagen gene expression (*Colla1*, *Colla2*, *Col4a1*)
137 in *Piezo2^{ff/ff}* and *Piezo2^{d/d}* cells supplemented with IGF1. Cells were cultured with IGF1 (100 ng/mL) or
138 vehicle for 24 h prior to RNA extraction. Expression was normalized to *Rpl19* ($n = 3$ per group, two-way
139 ANOVA with Tukey's post hoc test). Quantified data are presented as mean $\pm$ SEM. ns, $P > 0.05$; * $P <$
140 0.05; ** $P < 0.01$; *** $P < 0.001$.



141 **Extended data Fig. 6. Hippo–YAP pathway validation and Pgr promoter analysis.** (A) Quantification
142 of p-FAK (Tyr397) and FAK protein levels normalized to α -tubulin, corresponding to the Western blot
143 shown in Fig. 5A. $n = 3$ biological replicates. (B) Quantification of p-CaMKII, p-FAK (Tyr397), and FAK
144 protein levels normalized to β -actin, corresponding to the Western blot shown in Fig. 5B. $n = 3$ biological
145 replicates. (C) Quantification of p-CaMKII, p-FAK (Tyr397), and FAK protein levels normalized to β -
146 actin, corresponding to the Western blot shown in Fig. 5C. $n = 3$ biological replicates. (D) Quantification
147 of p-FAK (Tyr397) and FAK protein levels normalized to β -actin, corresponding to the Western blot
148 shown in Fig. 5E. $n = 3$ biological replicates. (E) Western blot analysis of p-YAP (Ser127) and p-TAZ
149 (Ser89) protein levels, along with total YAP and TAZ, in mouse endometrial stromal cells (MESC)s and
150 human endometrial stromal cells (T-HESC)s. (F) Quantification of p-YAP (Ser127) and YAP protein
151 levels normalized to α -tubulin, corresponding to the Western blot shown in Fig. 5G. $n = 3$ biological
152 replicates. (G) Quantification of p-YAP (Ser127) and YAP protein levels normalized to α -tubulin,
153 corresponding to the Western blot shown in Fig. 5I. $n = 3$ biological replicates. (H) Western blot analysis
154 of p-YAP (Ser127) and total YAP in primary mouse endometrial stromal cells (MESC)s from *Piezo2^{ff}*
155 and *Piezo2^{d/d}* mice treated with TDI. Cells were cultured with COL-I coating, TDI (no COL-I), or COL-
156 I+TDI, with a control cultured without COL-I coating or treatment (Vehicle). Cultures were maintained
157 for 48 h and then lysed. Relative expression was determined by normalizing to α -tubulin and based on n
158 $= 3$ per group (two-way ANOVA with Tukey's post hoc test, graph on right). Data are presented as mean
159 $\pm$ SEM. ns, $P > 0.05$; *** $P < 0.001$. (I) Schematic of the *Pgr* promoter region showing the predicted
160 TEAD-binding element (TRE) located at positions -138 to -127 relative to the transcription start site
161 (TSS), as identified using the JASPAR transcription factor binding site database (<https://jaspar.elixir.eu>).
162 The wild-type sequence (AGGTGGGAAG) and mutant sequence (AGGATTTAAG) used in luciferase
163 reporter assays are indicated. (J) Schematic of the TDI-011536 (TDI) treatment protocol for *in vivo* rescue
164 experiments. TDI ($3 \mu\text{M}$) was administered daily from 1.5 to 3.5 dpc, and implantation was assessed at
165 4.5 dpc. Quantitative data are presented as mean $\pm$ SEM. Statistical significance was determined using a
166 two-way ANOVA with Tukey's post hoc test. ns, $P > 0.05$; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.



167 **Extended data Fig. 7. Human endometrial stromal cell signaling cascade and PGR isoform-specific**
168 **rescue. (A)** Serum levels of cortisol (CORT), epinephrine (EPI), and noradrenaline (NA) in fertile controls
169 (Ctrl) and patients with recurrent implantation failure (RIF). Individual data points are displayed for Ctrl
170 and RIF endometrial samples ($n = 8$ per group, two-sided Student's t-test). (B) Quantification of p-
171 CaMKII, p-FAK (Tyr397), FAK, p-YAP (Ser127), and YAP protein levels normalized to β -actin,
172 corresponding to the Western blot shown in Fig. 6E. $n = 3$ biological replicates. (C, D) Western blot
173 analysis evaluating p-CaMKII (Thr286), p-FAK (Tyr397), total FAK, p-YAP (Ser127), and total YAP in
174 T-HESCs. PIEZO2-KD and control cells were cultured on COL-I, or without COL-1 (Vehicle), with or
175 without BAPTA-AM (C) or with or without KN93 (D). Lysates were obtained at 48 h post-treatment, and
176 expression was normalized to β -actin ($n = 3$ independent experiments, two-way ANOVA with Tukey's
177 post hoc test), with graphical quantification shown on the right. (E) Quantitative RT-PCR analysis of *PR*
178 (total PGR) and *PRB* (PGR-B isoform) expression in T-HESCs. PIEZO2-KD and control cells were
179 cultured as non-decidualized cells (Vehicle), decidualized (Dec) cells, or Dec cells with a vector
180 overexpressing PGR-A (PRA+Dec) or PGR-B (PRB+Dec). After 48 h, RNA was extracted and expression
181 was normalized to *GAPDH* ($n = 3$ per group, two-way ANOVA with Tukey's post hoc test). Quantitative
182 data are presented as mean $\pm$ SEM. ns, $P > 0.05$; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

183 **Extended data Table 1. Fertility assessment of *Piezo2^{ff}* and *Piezo2^{d/d}* females**

Genotype	Females	Pups	Litters	Litters/Female	Pups/Female	Pups/Litter	Weight/Pup	Gestation Days
<i>Piezo2^{ff}</i>	8	351	45	5.63±0.183	43.88±1.747	7.80±0.122	1.37±0.002	19.13±0.072
<i>Piezo2^{d/d}</i>	8	250	42	5.25±0.164	31.25±1.461	5.95±0.125	1.16±0.005	21.08±0.150
<i>P</i> value	/	/	/	0.1489	7.23 × 10⁻⁵	4.31 × 10⁻⁸	3.00 × 10⁻¹⁵	4.94 × 10⁻⁷

184 Data are presented as mean ± SEM. Statistical comparisons were performed using a two-sided Student's *t*-test. Bold values indicate significant
185 differences between genotypes.

Extended data Table 2. Patient clinical characteristics

Category	ID	Age	BMI (Kg/m ²)	Fertility history	Pathology	Time of embryo transfer	Sample collection phase	Embryo transfer day	Numbers of live births	Embryo quality grade	Endocrine metabolic abnormality
Pregnant (n=18)	B1	33	22.24	Yes	mechanical obstruction of fallopian	1	HRT cycle mid- secretory phase (LHP+7)	LHP+5 (Next cycle after sample collection)	1	I-II	Without
	B2	32	23.24								
	B3	39	23.07								
	B4	31	21.56								
	B5	32	19.47								
	B6	35	21.41								
	B7	34	22.50								
	B8	29	22.18								
	S1	31	22.31								
	S2	36	23.26								
	S3	27	19.47								
	S4	33	20.14								
	S5	38	22.27								
	S6	30	22.47								
	S7	31	23.14								
	S8	32	21.57								
	S9	34	22.21								
	S10	30	21.63								
RIF (n=18)	B9	35	21.88	No	Unexplained	4	HRT cycle mid- secretory phase (LHP+7)	LHP+5 (Next cycle after sample collection)	0	I-II	Without
	B10	31	20.21			3					
	B11	38	21.99			3					
	B12	27	20.07			4					
	B13	30	19.10			3					
	B14	34	22.67			2					
	B15	36	20.70			3					
	B16	34	23.75			4					
	S11	26	19.65			3					
	S12	29	20.83			3					
	S13	30	25.61			2					
	S14	29	29.69			2					
	S15	32	29.24			5					
	S16	29	23.71			3					

	S17	35	27.06	2
	S18	30	24.00	2
	S19	32	22.37	2
	S20	31	26.84	4
<i>p</i> -value		0.3016	0.2752	<0.0001

Gene	Primer application	Forward sequence	Reverse sequence
<i>mPiezo2</i>	qPCR	5'-CCCGGAACAGTAGCCAGAAA-3'	5'-TTCCCAAAGGCCCCAGAATCC-3'
<i>mPiezo2-</i>	qPCR	5'-ACTCGTCTGCATCCTACACC-3'	5'-GGTACACGCGACCATTGACT-3'
<i>E17/18</i>	qPCR	5'-GCTGGCTTCTGGAGGTGACTTG-3'	5'-GGTACACGCGACCATTGACT-3'
<i>mPiezo1</i>			
<i>mPgr</i>	qPCR	5'-GCAATGGAAGGGCAGCATAAC-3'	5'-CTTACGACCTCCAAGGACCA-3'
<i>mEsr1</i>	qPCR	5'-GGAAGCTCCTGTTTGCTCCT-3'	5'-AACCGACTTGACGTAGCCAG-3'
<i>mRpl19</i>	qPCR	5'-TCATGGAGCACATCCACAAGCTGA-3'	5'-CGCTTTCGTGCTTCCTTGGTCTTA-3'
<i>mLif</i>	qPCR	5'-ATTGAGCTTGACCTGGAGGC-3'	5'-GGTGGAGCTGTATCGGATGG-3'
<i>mNos1</i>	qPCR	5'-CCTTCACAGGGGATGGAACC-3'	5'-AGATCGACAGCTTTGGTGGG-3'
<i>mNos2</i>	qPCR	5'-AGAGCCACAGTCCTCTTGC-3'	5'-CTGGTCCATGCAGACAACCT-3'
<i>mNos3</i>	qPCR	5'-ATTGGCATGAGGGACCTGTG-3'	5'-GGGATGAGGTTGTCCTGGTG-3'
<i>mLtf</i>	qPCR	5'-CCCTACAACTGCGACCTGT-3'	5'-CACGACTGCTACCGCATAGT-3'
<i>mMuc1</i>	qPCR	5'-TCACCCCAGTTGTCTGTTGG-3'	5'-AGTAGTTGGAGCTGGGGTCT-3'
<i>mMuc4</i>	qPCR	5'-ATGGCTACAAAGGCTACCACC-3'	5'-CCTCACTACATGGGGACACAC-3'
<i>mHand2</i>	qPCR	5'-CCAGCTACATCGCCTACCTC-3'	5'-GGTCTTCTTGATCTCCGCCT-3'
<i>mHoxa10</i>	qPCR	5'-TCCAAAGGCGAAAATGCAGC-3'	5'-TCTGGTGCTTCGTGTAAGGG-3'
<i>mAreg</i>	qPCR	5'-ACAGCGAGGATGACAAGGAC-3'	5'-GCCAATAGCTGCGAGGATGA-3'
<i>mIhh</i>	qPCR	5'-TCTGTCTGTTCCCTGCTGCTG-3'	5'-TTGTAGGCAAGAGGCACGAG-3'
<i>mBmp2</i>	qPCR	5'-AAGCCAAACACAAACAGCGG-3'	5'-CGATCCAGTCATTCCACCCC-3'
<i>mPrl8a2</i>	qPCR	5'-GGTGCATGGATCACTCCTGT-3'	5'-ACCCACGTAAGGTCATCATGG-3'
<i>mPrl3c1</i>	qPCR	5'-ACTGCCACACGATATGACCG-3'	5'-TGTGACGAGAAGAGGAAAGCA-3'
<i>mColla1</i>	qPCR	5'-AGCACGTCTGGTTTGGAGAG-3'	5'-GACATTAGGCGCAGGAAGGT-3'
<i>mColla2</i>	qPCR	5'-CTAGCCAACCGTGCTTCTCA-3'	5'-TCTCCTCATCCAGGTACGCA-3'
<i>mCol3a1</i>	qPCR	5'-ACGTAAGCACTGGTGGACAG-3'	5'-CAGGAGGGCCATAGCTGAAC-3'
<i>mCol4a1</i>	qPCR	5'-TCATTAGCAGGTGTGCGGTT-3'	5'-GTTAGGGCACTGCGGAATCT-3'
<i>mIl1b</i>	qPCR	5'-ATGCCACCTTTTGACAGTGAT-3'	5'-AGCCCTTCATCTTTTGGGGT-3'
<i>mIl6</i>	qPCR	5'-CCCCAATTTCCAATGCTCTCC-3'	5'-GGATGGTCTTGGTCCTTAGCC-3'
<i>mTnfa</i>	qPCR	5'-GATCGGTCCCCAAAGGGATG-3'	5'-CCACTTGGTGGTTTGTGAGTG-3'
<i>mIgfl</i>	qPCR	5'-TGGGGAAAATCAGCAGCCTT-3'	5'-CAGAGCGCCAGGTAGAAGAG-3'
<i>hGAPDH</i>	qPCR	5'-GGCAAATTCCATGGCACCG-3'	5'-AGCATCGCCCCACTTGATTT-3'
<i>hPIEZO2</i>	qPCR	5'-AAGGCAGTCCACCCTTGATG-3'	5'-TGGGCTCGCTTGATGCTAAA-3'
<i>hPGR-B</i>	qPCR	5'-GGAATGGGCTGTACCGAGAG-3'	5'-CGGCTCCTTTATCTCCCGAC-3'
<i>hPGR</i>	qPCR	5'-CCGGGCACTGAGTGTTGAAT-3'	5'-GAAGGGGTTTCACCATCCCT-3'

<i>hCOL1A1</i>	qPCR	5'-GGGAGATGCTGGTCCTGCT-3'	5'-GCACCATCATTTCCACGAGC-3'
<i>hIGFBP1</i>	qPCR	5'-TGTGACGAGAAGAGGAAAGCA-3'	5'-CTCCTGATGTCTCCTGTGCC-3'
<i>hFOXO1</i>	qPCR	5'-TGGAGATCGACCCGGACTTC-3'	5'-CCGAGTTGGACTGGCTAAACT-3'
<i>hPRL</i>	qPCR	5'-GGGGTTCATTACCAAGGCCA-3'	5'-TTGGGCTTGCTCCTTGTCTT-3'
<i>hIGF1</i>	qPCR	5'-ATGTATTGCGCACCCCTCAA-3'	5'-GCACTCCCTCTACTTGCGTT-3'
<i>siPIEZO2</i>	RNA interference	5'-GCTTGAGGATCTTGGCTTA-3'	
<i>Padlock</i>	sm-ISH	5'-TCTTTATGGTAAACGCCTTTGTTCCCTTTTACGACCTCAATG	
<i>Probe-PIEZO2</i>		CTGCTGCTGTACTACTCTTGACATAGATCTGTAGAGTCT-3'	
<i>Oligo-dT</i>	sm-ISH	5'-AAGCAGTGGTATCAACGCAGAGTACTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTVN-3'	
<i>RCA primer</i>	sm-ISH	5'-ACAGCAGCAGCATTGAGG*T*C-3' Note: *= thiophosphate modifications	
<i>HRP labeled detection probe</i>	sm-ISH	5'- HRP-CCTCAATGCTGCTGCTGTACTAC-3'	
<i>Pgr Promoter-WT</i>	Dual-Luciferase Reporter Assay	5'- GCTTTTTATAGGGTGGGGCTGGCATGCTTCTTAATTCAGTACAAATTCTGATATGTTT AGATTTCTTCTCAAAGCATCTGATATTCAGGTGGAATCAGACTAGTGAGTAGCTGG AATCCAACGCCAGAGATTAGATCTAGCCAGTGATTGGCTAGGGAGGGGCTTTGGG CGGGCCTTCCTAGAGCGCCAACGCTTGCTAGAAAGCTATGGAGCCAGTCTAGACTGT CACTATCAGTCTTTGTAGTATTTACGGGTGCGAGGATTC-3'	
<i>Pgr Promoter-Mut</i>	Dual-Luciferase Reporter Assay	5'- GCTTTTTATAGGGTGGGGCTGGCATGCTTCTTAATTCAGTACAAATTCTGATATGTTT AGATTTCTTCTCAAAGCATCTGATGAGTAAGGTGGAATCAGACTAGTGAGTAGCTGG AATCCAACGCCAGAGATTAGATCTAGCCAGTGATTGGCTAGGGAGGGGCTTTGGG CGGGCCTTCCTAGAGCGCCAACGCTTGCTAGAAAGCTATGGAGCCAGTCTAGACTGT CACTATCAGTCTTTGTAGTATTTACGGGTGCGAGGATTC-3'	
<i>Piezo2-WT</i>	Genotyping	5'-ACTTAGATGGGGCAGGTGCT-3'	5'-ACTTCCCTACCCACCCATTC-3'
<i>Piezo2-flox</i>	Genotyping	5'-GCCGCTCTAGAACTAGTGGA-3'	5'-ATCTACCACGGGGGCTCTCTC-3'
<i>Pgr-WT</i>	Genotyping	5'-TATACCGATCTCCCTGGACG-3'	5'-CCCAAAGAGACACCAGGAAG-3'
<i>Pgr-Cre</i>	Genotyping	Same as <i>Pgr-WT</i>	5'-ACTTAGATGGGGCAGGTGCT-3'

Extended data Table 4. List of antibodies

Antibody	Type	Manufacturer	Catalog no.	Application	Dilution
Ki67	Rabbit IgG	Santa Cruz	SC-23900	IHC	1:200
Cox2	Rabbit IgG	Cell Signaling	4842	IHC	1:200
Hand2	Rabbit IgG	Abcam	ab200040	IHC	1:200
Muc1	Rabbit IgG	Cell Signaling	16564S	IHC	1:200
Pgr	Rabbit IgG	Invitrogen	MA5-16393	IHC	1:200
ER α	Rabbit IgG	Santa Cruz	sc7207	IHC	1:200
		Biotechnology			
Biotinylated secondary	Goat anti-	Vactorlabs	BA-1000	IHC	1:200
	Rabbit IgG				
Collagen I	Rabbit IgG	Abways	ab138492	IF	1:100
Fluorescent secondaries	Goat anti-	Jackson	711-225-152	IF	1:200
	Rabbit IgG				
Piezo2	Rabbit IgG	NOVUS	NBP1-78624	WB	1:1000
FAK	Rabbit IgG	PTM BIO	PTM-5717	WB	1:1000
p-FAK	Rabbit IgG	Cell Signaling	8556T	WB	1:1000
p-CaMKII	Rabbit IgG	Affinity Biosciences	AF3493	WB	1:1000
α -Tubulin	Rabbit IgG	Cell Signaling	2144	WB	1:1000
YAP/TAZ	Rabbit IgG	Cell Signaling	93622S	WB	1:1000
p-YAP	Rabbit IgG	Cell Signaling	4911S	WB	1:1000
β -actin	Mouse IgG	Abmart	M20011XS	WB	1:1000
Streptavidin–HRP conjugate	N/A	Invitrogen	434323	IHC	1:1000
GAPDH	Mouse IgG	Abmart	M20050XS	WB	1:1000

IHC, immunohistochemistry; IF, immunofluorescence; WB, western blot.