

Supporting Information

Cooperative NHC/Lewis Acid Catalysis for Stereoselective Strain-Release Geminal Difunctionalization of Bicyclo[1.1.0]butanes to 1,1,3,3-Tetrasubstituted Cyclobutanes

Sayali G. Jagtap,^{ac} Gaurang J. Bhatt,^a Rajesh G. Gonnade^{bc} and Santosh B. Mhaske^{*ac}

^aOrganic Chemistry Division, CSIR-National Chemical Laboratory (CSIR-NCL), Dr. Homi Bhabha Road, Pune-411008 (India).

^bPhysical and Materials Chemistry Division, CSIR-National Chemical Laboratory (CSIR-NCL), Dr. Homi Bhabha Road, Pune-411008 (India).

^cAcademy of Scientific and Innovative Research (AcSIR), Ghaziabad-201002 (India).

*E-mail: sb.mhaske.ncl@csir.res.in

Contents

1.	General Information	S1
2.	Experimental Section	S2
3.	Mechanistic Studies	S6
4.	Gram Scale Experiment	S8
5.	Synthetic Transformations	S9
6.	Characterization Data of Compounds	S11
7.	References	S25
8.	X-ray crystallographic data of compound 3g and 4g	S26
9.	Copies of ¹ H, ¹³ C NMR and HRMS Spectra	S37

1. General Information


All reagents were used as received from commercial sources without additional purification. Solvents were purified and dried by standard techniques. All glassware were stored in an oven or flame-dried (60-70 °C) for at least 3 h before use. Room temperature (rt) corresponds to the 25-35 °C, when the experiments were performed. Catalytic reactions were performed under an inert atmosphere of argon using schlenk tube. Thin layer chromatography (TLC) was performed using silica gel precoated aluminium foil, which was visualized with UV light at 254 nm or under iodine / 2,4-DNP / KMnO₄ charring. Column chromatography was performed with flash silica-gel (230-400 mesh). ¹H NMR (400, or 500 MHz), ¹³C{¹H} NMR (100.6, or 126 MHz) spectra were recorded on the NMR spectrometer. The flip angle of the DEPT spectra are 135 degrees. Deuterated chloroform was used as solvent, and chemical shift values (δ) are reported in parts per million relative to the residual signal of the chloroform [δ 7.27 for ¹H (chloroform-*d*), δ 77.0 for ¹³C{¹H} (chloroform-*d*)]. Abbreviations used in the NMR data: br, broad. s, singlet. br s, broad singlet. d, doublet. t, triplet. q, quartet. quint, quintet. sext, sextet. m, multiplet. Coupling constants (*J*) are reported in hertz. Melting points were measured using a melting point apparatus in open glass capillaries. High-resolution mass spectrometry (HRMS) was performed on QTOF/ESI-TOF mass spectrometer.

Abbreviations: NHC - N-Heterocyclic carbene, THF - Tetrahydrofuran, DCM -Dichloromethane, ACN: Acetonitrile, DCM: Dichloromethane, DCE: Dichloroethane, DMF: *N*, *N*-Dimethylformamide, EtOAc: Ethyl Acetate, NHC: N-Heterocyclic carbene, DIPEA- *N*, *N*-Diisopropylethylamine, DBU- 1,8-Diazabicyclo[5.4.0]undec-7-ene, rt - room temperature.

2. Experimental section


2.1. Preparation of starting materials

The α,β -unsaturated aldehydes **1a-b**, **1d-g**, **1k-1l**, **1o** were purchased from commercial sources. All other α,β -unsaturated aldehydes **1c**, **1h**, **1i-j**, **1m-n** and **1p** were synthesized by following the literature procedure.¹⁻³ BCBs (**2a-2s**) are known in the literature and were prepared according to corresponding literature procedures and all the spectral data are in agreement with the reported values⁴⁻⁸. The achiral N-heterocyclic carbene precursor NHCs (**I-K**) was prepared following the literature procedure.⁹⁻¹¹ The chiral NHCs (**A-H**) and achiral NHC **L** were purchased from commercial sources.



2.2. Protocol optimization:


2.2. A) General procedure for the optimization of the reaction condition for methyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4d**):**



An oven-dried schlenk tube equipped with a magnetic stirring bar was charged with a NHC and base. The reaction vessel was evacuated and backfilled with Argon three times. Under an argon atmosphere, α,β -unsaturated aldehyde **1d** and methyl 3-phenylbicyclo[1.1.0]butane-1-carboxylate (**2a**) were added followed by addition of dry solvent. The reaction vessel was evacuated and backfilled with argon three times. The schlenk tube was placed in an ice bath and the reaction mixture was stirred for 5 minutes. After this, Lewis acid was added and the tube was tightly closed with a glass stopper. The resulting mixture was kept for stirring at rt for completion of the reaction. Upon completion of the reaction as observed by TLC, the reaction mixture was directly subjected to column chromatography using ethyl acetate:pet ether over 230-400 flash silica gel to afford the corresponding product **4d**.

Various permutations and combinations attempted are provided in the Table S1-S4.

Table S1: Screening of different NHCs:^a




Entry	Variation of NHCs from Standard Reaction Condition	Yield 3d(%) ^b	Yield 4d(%) ^b
1.	NHC A (20 mol%)	trace	76%
2.	NHC B (20 mol%)	5%	80%
3.	NHC C (20 mol%)	trace	78%
4.	NHC D (20 mol%)	trace	74%
5.	NHC E (20 mol%)	21%	58%
6.	NHC F (20 mol%)	trace	trace

7.	NHC G (20 mol%)	36%	29%
8.	NHC H (20 mol%)	40%	20%
9.	NHC I (20 mol%)	14%	53%
10.	NHC J (20 mol%)	N.R.	N.R.
11.	NHC K (20 mol%)	28%	43%
12.	NHC L (20 mol%)	N.R.	N.R.
13.	NHC A (10 mol%)	-	85%
14.	NHC B (10 mol%)	7%	71%
15.	NHC C (10 mol%)	45%	33%

^aStandard Reaction conditions: **1d** (1 equiv., 0.16 mmol), **2a** (2 equiv., 0.32 mmol), NHC-A (10 mol%, 0.016 mmol), K₂CO₃ (40 mol%, 0.064 mmol), Sc(OTf)₃ (30 mol%, 0.045 mmol), THF (0.2 mL), 0°C-rt, 15 min. ^bIsolated yield. N.R.: No reaction


Table S2: Screening of different Lewis acids:^a



Entry	Variation of Lewis acids from Standard Reaction Condition	Yield 3d(%) ^b	Yield 4d(%) ^b
1.	Sc(OTf)₃ (30 mol%)	-	85%
2.	Sc(OTf) ₃ (20 mol%)	20%	57%
3.	Bi(OTf) ₃ (30 mol%)	N.R.	N.R.
4.	Cu(OTf) ₂ (30 mol%)	N.R.	N.R.
5.	AgOTf (30 mol%)	N.R.	N.R.
6.	Yb(OTf) ₃ (30 mol%)	N.R.	N.R.

^aStandard Reaction conditions: **1d** (1 equiv., 0.16 mmol), **2a** (2 equiv., 0.32 mmol), NHC-A (10 mol%, 0.016 mmol), K₂CO₃ (40 mol%, 0.064 mmol), Sc(OTf)₃ (30 mol%, 0.045 mmol), THF (0.2 mL), 0°C-rt, 15 min. ^bIsolated yield. N.R.: No reaction

Table S3: Screening of different bases:^a




Entry	Variation of bases from Standard Reaction Condition	Yield 3d(%) ^b	Yield 3c(%) ^b
1.	K₂CO₃ (40 mol%)	-	85%
2.	K ₃ PO ₄ (40 mol%)	N.R.	N.R.

3.	Cs ₂ CO ₃ (40 mol%)	-	24%
4.	DBU (40 mol%)	-	18%
5.	DIPEA (40 mol%)	trace	trace
6.	Na ₂ CO ₃ (40 mol%)	63%	19%

^aStandard Reaction conditions: **1d** (1 equiv., 0.16 mmol), **2a** (2 equiv., 0.32 mmol), NHC-A (10 mol%, 0.016 mmol), K₂CO₃ (40 mol%, 0.064 mmol), Sc(OTf)₃ (30 mol%, 0.045 mmol), THF (0.2 mL), 0°C-rt, 15 min. ^bIsolated yield. N.R.: No reaction


Table S4: Screening of different solvents:^a



Entry	Variation of solvents from Standard Reaction Condition	Yield 3d(%) ^b	Yield 3c(%) ^b
1.	THF (0.2 mL)	-	85%
2.	Toluene (0.2 mL)	trace	49.4%
3.	DCM (0.2 mL)	N.R.	N.R.
4.	Chloroform (0.2 mL)	10%	48%
5.	ACN (0.2 mL)	89%	-
6.	1,4-dioxane (0.2 mL)	trace	47%
7.	THF (0.5 mL)	12%	70%
8.	THF (0.7 mL)	56%	27%

^aStandard Reaction conditions: **1d** (1 equiv., 0.16 mmol), **2a** (2 equiv., 0.32 mmol), NHC-A (10 mol%, 0.016 mmol), K₂CO₃ (40 mol%, 0.064 mmol), Sc(OTf)₃ (30 mol%, 0.045 mmol), THF (0.2 mL), 0°C-rt, 15 min. ^bIsolated yield. N.R.: No reaction


2.2. B) General optimized procedure for the synthesis of polyfunctionalised cyclobutane derivatives 4



An oven-dried schlenk tube equipped with a magnetic stirring bar was charged with the NHC precatalyst-A (10 mol%) and K₂CO₃ (40 mol%). The reaction vessel was evacuated and backfilled with argon three times. Under an argon atmosphere, α,β -unsaturated aldehyde **1** (1 equiv.) and bicyclobutane derivative **2** (2 equiv.) were added followed by addition of THF (0.2 mL). The reaction vessel was evacuated and backfilled with argon three times. The schlenk tube was placed in an ice bath and the reaction mixture was stirred for 5 minutes. After this, Sc(OTf)₃ (30 mol%) was added

and the tube was tightly closed with a glass stopper. The resulting mixture was kept for stirring at rt for completion of the reaction. Upon completion of the reaction as observed by TLC, the reaction mixture was directly subjected to column chromatography using ethyl acetate:pet ether over 230-400 flash silica gel to afford the desired products **4** in good to excellent yields.


2.2. C) Typical procedure for the synthesis of representative product **4d**



An oven-dried schlenk tube equipped with a magnetic stirring bar was charged with NHC precatalyst-**A** (6.7 mg, 0.016 mmol, 10 mol%) and K_2CO_3 (8.8 mg, 0.064 mmol, 40 mol%). The reaction vessel was evacuated and backfilled with argon three times. Under an argon atmosphere, (E)-3-(4-Chlorophenyl)acrylaldehyde (**1d**, 27 mg, 0.16 mmol, 1 equiv.), methyl 3-phenylbicyclo[1.1.0]butane-1-carboxylate (**2a**, 60 mg, 0.32 mmol, 2 equiv.) were added followed by addition of dry THF (0.2 mL). The schlenk tube was placed in an ice bath and the reaction mixture was stirred for 5 minutes. After this, Sc(OTf)_3 (24 mg, 0.048 mmol, 30 mol%) was added and the tube was tightly closed with a glass stopper. The resulting mixture was kept for stirring at rt for 15 min. Upon completion of the reaction as observed by TLC, the reaction mixture was directly subjected to column chromatography using ethyl acetate:pet ether (1:4) over 230-400 flash silica gel to afford the methyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (**4d**) as a yellow oil with 85% yield (48 mg, >19:1 dr).

3. Mechanistic studies and control experiments


3.1 Reaction with saturated aldehyde



An oven-dried schlenk tube equipped with a magnetic stirring bar was charged with NHC precatalyst-**A** (6.7 mg, 0.016 mmol, 10 mol%) and K_2CO_3 (8.8 mg, 0.064 mmol, 40 mol%). The reaction vessel was evacuated and backfilled with argon three times. Under an argon atmosphere, 3-phenylpropanal (**1a'**, 21 mg, 0.16 mmol, 1 equiv.), methyl 3-phenylbicyclo[1.1.0]butane-1-carboxylate (**2a**, 60 mg,


0.32 mmol, 2 equiv.) were added followed by addition of dry THF (0.2 mL). The schlenk tube was placed in an ice bath and the reaction mixture was stirred for 5 minutes. After this, Sc(OTf)₃ (24 mg, 0.048 mmol, 30 mol%) was added and the tube was tightly closed with a glass stopper. The resulting mixture was stirred at room temperature for 24 h. After monitoring the reaction for different time intervals, no formation of **3a'** and/or **4a'** was observed.

3.2 Reactivity of Intermediate under standard reaction conditions



An oven-dried schlenk tube equipped with a magnetic stirring bar was charged with NHC precatalyst-A (3.5 mg, 0.008 mmol, 10 mol%) and K₂CO₃ (4.6 mg, 0.033 mmol, 40 mol%). The reaction vessel was evacuated and backfilled with argon three times. Under an argon atmosphere, methyl (E)-3-(4-chlorostyryl)-1-phenyl-2-oxabicyclo[2.1.1]hexane-4-carboxylate (**3d**, 30 mg, 0.085 mmol, 1 equiv.) was added followed by addition of dry THF (0.2 mL). The schlenk tube was placed in an ice bath and the reaction mixture was stirred for 5 minutes. After this, Sc(OTf)₃ (12.5 mg, 0.025 mmol, 30 mol%) was added and the tube was tightly closed with a glass stopper. The resulting mixture was kept for stirring at rt for 15 min. Upon completion of the reaction as observed by TLC, the reaction mixture was directly subjected to column chromatography using ethyl acetate:pet ether (1:4) over 230-400 flash silica gel to afford the methyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (**4d**) as a yellow solid with 93% yield (28 mg, >19:1 dr).


3.3 Reactivity of Intermediate under Standard Reaction Conditions in the Absence of NHC precatalyst



An oven-dried schlenk tube equipped with a magnetic stirring bar was charged with K₂CO₃ (7.8 mg, 0.056 mmol, 40 mol%). The reaction vessel was evacuated and backfilled with argon three times. Under an argon atmosphere, methyl (E)-3-(4-chlorostyryl)-1-phenyl-2-oxabicyclo[2.1.1]hexane-4-

carboxylate (**3d**, 50 mg, 0.14 mmol, 1 equiv.) was added followed by addition of dry THF (0.5 mL). The schlenk tube was placed in an ice bath and the reaction mixture was stirred for 5 minutes. After this, Sc(OTf)₃ (21 mg, 0.042 mmol, 30 mol%) was added and the tube was tightly closed with a glass stopper. The resulting mixture was stirred at room temperature for 24 h. After monitoring the reaction for different time intervals, no formation of **4d** was observed.


3.4 Reactivity of Intermediate under Standard Reaction Conditions in the Absence of Sc(OTf)₃



An oven-dried schlenk tube equipped with a magnetic stirring bar was charged with NHC precatalyst-A (5.9 mg, 0.014 mmol, 10 mol%) and K₂CO₃ (7.8 mg, 0.056 mmol, 40 mol%). The reaction vessel was evacuated and backfilled with argon three times. Under an argon atmosphere, methyl (E)-3-(4-chlorostyryl)-1-phenyl-2-oxabicyclo[2.1.1]hexane-4-carboxylate (**3d**, 50 mg, 0.14 mmol, 1 equiv.) was added followed by addition of dry THF (0.5 mL). The resulting mixture was stirred at room temperature for 24 h. After monitoring the reaction for different time intervals, no formation of **4d** was observed.


4. Gram Scale Experiments

4.1. Gram Scale synthesis of Methyl (1*s*,3*s*)-3-(4-(*tert*-butyl)phenyl)-3-((*E*)-4-chlorostyryl)-1-formylcyclobutane-1-carboxylate (**4x**)



reaction mixture was directly subjected to column chromatography using ethyl acetate:pet ether (1:19) over 230-400 flash silica gel to afford the Methyl (1s,3s)-3-(4-(tert-butyl)phenyl)-3-((E)-4-chlorostyryl)-1-formylcyclobutane-1-carboxylate (**4x**) as a white solid with 89% yield (1.1 g, >19:1 dr).


4.2. Gram Scale synthesis of Methyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-phenyl cyclobutane-1-carboxylate (**4d**)



A 25mL round bottom flask was charged with NHC precatalyst-A (251 mg, 0.6 mmol, 10 mol%) and K_2CO_3 (331 mg, 1.2 mmol, 40 mol%). The reaction vessel was evacuated and backfilled with argon three times. Under an argon atmosphere, (E)-3-(4-Chlorophenyl)acrylaldehyde (**1d**, 1 g, 6 mmol, 1 equiv.), methyl 3-phenylbicyclo[1.1.0]butane-1-carboxylate (**2a**, 2.26 g, 12 mmol, 2 equiv.) were added followed by addition of dry THF (4 mL). The reaction vessel was placed in an ice bath and the reaction mixture was stirred for 5 minutes. After this, $Sc(OTf)_3$ (886 mg, 0.9 mmol, 30 mol%) was added and the resulting mixture was kept for stirring at rt for 15 min. The reaction mixture was then directly subjected to column chromatography using ethyl acetate:pet ether (1:19) over 230-400 flash silica gel to afford the Methyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-phenyl cyclo-butane-1-carboxylate (**4d**) as a yellow solid. (1.51 g, 71% yield, >19:1 dr).

5. Synthetic Transformations


5.1 Synthesis of compound 5



To a solution of methyl (1s,3s)-3-(4-(tert-butyl)phenyl)-3-((E)-4-chlorostyryl)-1-formylcyclobutane-1-carboxylate (**4x**, 100 mg, 0.24 mmol) in EtOAc (8 mL), 10% Pd/C (25.9 mg, 0.024 mmol, 0.1 equiv.) was added and the resulting mixture was stirred under H_2 atmosphere for 1 h. After complete consumption of the starting material, the reaction mixture was filtered through a pad of celite and concentrated under reduced pressure. The reaction mixture was then subjected to column chromatography using ethyl acetate:pet ether (1: 9) over 230-400 flash silica gel to afford the methyl

(1*r*,3*s*)-3-(4-(*tert*-butyl)phenyl)-3-(4-chlorophenethyl)-1-formylcyclobutane-1-carboxylate (**5**) as a colourless oil. (85 mg, 85% yield).


5.2 Synthesis of compound 6



To the solution of Methyl (1*s*,3*s*)-3-((*E*)-4-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (**4d**, 30 mg, 0.085 mmol, 1 equiv.) in dry THF (0.5 mL) at 0 °C was added NaBH₄ (3.2 mg, 0.085 mmol, 1.0 equiv.). The reaction mixture was warmed to rt and stirred for 1 h. after completion of reaction as monitored by TLC, the resulting reaction mixture was quenched with brine (5 mL) and extracted with ethyl acetate (10 mL × 3). The combined organic layer was dried over anhydrous Na₂SO₄, and concentrated in vacuo. The obtained crude product was purified by flash chromatography (20% EtOAc in petroleum ether) to yield methyl (1*s*,3*s*)-3-((*E*)-4-chlorostyryl)-1-(hydroxymethyl)-3-phenylcyclobutane-1-carboxylate (**6**, 29 mg, 96% yield) as a yellow oil.


6. Characterization Data

Methyl (1*s*,3*s*)-1-formyl-3-phenyl-3-((*E*)-styryl)cyclobutane-1-carboxylate (4a)




Yellow oil, 88% (45 mg) isolated yield, $R_f = 0.6$ in 20% EtOAc: Pet Ether. $^1\text{H NMR}$ (400MHz, $\text{CHLOROFORM-}d$): δ ppm 9.82 (s, 1 H), 7.38 - 7.33 (m, 2 H), 7.29 (t, $J = 2.2$ Hz, 2 H), 7.26 - 7.22 (m, 5 H), 7.21 - 7.16 (m, 1 H), 6.38 (d, $J = 15.9$ Hz, 1 H), 6.10 (d, $J = 16.0$ Hz, 1 H), 3.68 (s, 3 H), 3.17 - 3.07 (m, 4 H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101MHz, $\text{CHLOROFORM-}d$): δ ppm 196.3, 171.3, 146.4, 136.7, 136.4, 128.5, 128.4, 127.4, 126.3, 126.0, 52.8, 52.7, 43.6, 37.7. **HRMS (ESI-TOF) m/z :** $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{21}\text{H}_{21}\text{O}_3$ 321.1485; Found 321.1485.

Methyl (1*s*,3*s*)-3-((*E*)-4-fluorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4b)



Yellow oil, 85% (46 mg) isolated yield, $R_f = 0.4$ in 20% EtOAc: Pet Ether. $^1\text{H NMR}$ (400MHz, $\text{CHLOROFORM-}d$): δ ppm 9.84 (s, 1 H), 7.40 - 7.35 (m, 2 H), 7.25 (m, 5 H), 6.96 (t, $J = 8.6$ Hz, 2 H), 6.31 (d, $J = 15.9$ Hz, 1 H), 6.05 (d, $J = 15.9$ Hz, 1 H), 3.70 (s, 3 H), 3.17 - 3.08 (m, 4 H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101MHz, $\text{CHLOROFORM-}d$): δ ppm 196.3, 171.3, 162.2 (d, $J_{\text{C-F}} = 246.4$ Hz), 146.3, 136.3, 136.3, 132.8 (d, $J_{\text{C-F}} = 3.81$ Hz), 128.5, 127.8 (d, $J_{\text{C-F}} = 8.39$ Hz), 127.3, 126.4, 126.0, 115.3 (d, $J_{\text{C-F}} = 22.13$ Hz), 52.7, 43.5, 37.7. **HRMS (ESI-TOF) m/z :** $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{21}\text{H}_{20}\text{O}_3\text{F}$ 339.1391; Found 339.1388.


Methyl (1*s*,3*s*)-3-((*E*)-3,4-difluorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4c)



Yellow oil, 79% (45 mg) isolated yield, $R_f = 0.41$ in 20% EtOAc: Pet Ether. $^1\text{H NMR}$ (400MHz, $\text{CHLOROFORM-}d$): δ ppm 9.84 (s, 1 H), 7.40 - 7.36 (m, 2 H), 7.26 - 7.22 (m, 3 H), 7.11 - 7.06 (m,


1 H), 7.06 - 7.01 (m, 1 H), 6.99 - 6.95 (m, 1 H), 6.31 (d, $J = 16.0$ Hz, 1 H), 5.99 (d, $J = 16.0$ Hz, 1 H), 3.71 (s, 3 H), 3.17 - 3.08 (m, 4 H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101MHz, CHLOROFORM- d): δ ppm 196.2, 171.2, 151.6, 150.2 (dd, $J_{\text{C-F}} = 247.2$, 13.0 Hz, 149.6 (dd, $J_{\text{C-F}} = 248.7$, 12.2 Hz), 146.0, 137.6 (d, $J_{\text{C-F}} = 2.3$ Hz), 134.0 (dd, $J_{\text{C-F}} = 6.1$, 3.8 Hz), 128.6, 126.5 (2C), 126.0, 122.5 (dd, $J_{\text{C-F}} = 6.1$, 3.0 Hz), 117.1 (d, $J_{\text{C-F}} = 17.5$ Hz), 114.6 (d, $J_{\text{C-F}} = 17.5$ Hz), 52.8, 52.7, 43.5, 37.5. HRMS (ESI-TOF) m/z : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{21}\text{H}_{19}\text{O}_3\text{F}_2$ 357.1297; Found 357.1290.

Methyl (1s,3s)-3-((*E*)-4-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4d)




Yellow solid, mp 56-58 °C, 85% (48 mg) isolated yield, $R_f = 0.3$ in 10% EtOAc: Pet Ether. ^1H NMR (400MHz, CHLOROFORM- d): δ ppm 9.86 (s, 1 H), 7.43 - 7.36 (m, 2 H), 7.28 (d, $J = 7.1$ Hz, 3 H), 7.26 - 7.21 (m, 4 H), 6.39 (d, $J = 16.0$ Hz, 1 H), 6.05 (d, $J = 15.9$ Hz, 1 H), 3.72 (s, 3 H), 3.20 - 3.09 (m, 4 H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101MHz, CHLOROFORM- d): δ ppm 196.3, 171.3, 146.1, 137.2, 135.2, 133.0, 128.6, 128.5, 127.5, 127.3, 126.4, 126.0, 52.8, 52.7, 43.6, 37.6. HRMS (ESI-TOF) m/z : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{21}\text{H}_{20}\text{O}_3^{35}\text{Cl}$ 355.1095; Found 355.1096.

Methyl (1s,3s)-3-((*E*)-3-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4e)




Colorless oil, 69% (39 mg) isolated yield, $R_f = 0.6$ in 10% EtOAc: Pet Ether. ^1H NMR (400MHz, CHLOROFORM- d): δ ppm 9.83 (s, 1 H), 7.40 - 7.35 (m, 2 H), 7.27 - 7.22 (m, 4 H), 7.22 - 7.11 (m, 3 H), 6.40 (d, $J = 15.9$ Hz, 1 H), 6.03 (d, $J = 16.0$ Hz, 1 H), 3.70 (s, 3 H), 3.18 - 3.07 (m, 4 H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101MHz, CHLOROFORM- d): δ ppm 196.2, 171.3, 146.0, 138.6, 138.0, 134.4, 129.7, 128.6, 127.4, 127.2, 126.5, 126.3, 126.0, 124.6, 52.8, 52.7, 43.6, 37.5. HRMS (ESI-TOF) m/z : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{21}\text{H}_{20}\text{O}_3^{35}\text{Cl}$ 355.1095; Found 355.1092.

Methyl (1s,3s)-3-((E)-2-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4f)




Yellow oil, 20% (11 mg) isolated yield, R_f = 0.41 in 20% EtOAc: Pet Ether. **^1H NMR** (400MHz, $\text{CHLOROFORM-}d$): δ ppm 9.83 (s, 1 H), 7.44 - 7.35 (m, 3 H), 7.33 - 7.27 (m, 3 H), 7.26 - 7.21 (m, 1 H), 7.16 (m, 2 H), 6.38 (d, J = 15.9 Hz, 1 H), 6.01 (d, J = 15.9 Hz, 1 H), 3.70 (s, 3 H), 3.17 - 3.07 (m, 4 H). **$^{13}\text{C}\{^1\text{H}\}$ NMR** (101MHz, $\text{CHLOROFORM-}d$): δ ppm 196.3, 171.3, 146.1, 139.1, 134.9, 133.1, 129.5, 128.6, 128.5, 126.9, 126.8, 126.5, 126.0, 124.9, 52.8, 52.8, 43.9, 37.6. **HRMS (ESI-TOF) m/z** : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{21}\text{H}_{20}\text{O}_3^{35}\text{Cl}$ 355.1095; Found 355.1095.

Methyl (1s,3s)-3-((E)-4-bromostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4g)




White solid, mp 91-93 °C, 82% (52 mg) isolated yield, R_f = 0.6 in 20% EtOAc: Pet Ether. **^1H NMR** (400MHz, $\text{CHLOROFORM-}d$): δ ppm 9.83 (s, 1 H), 7.41 - 7.35 (m, 4 H), 7.24 (d, J = 8.1 Hz, 2 H), 7.15 (d, J = 8.5 Hz, 2 H), 6.38 (d, J = 15.9 Hz, 1 H), 6.01 (d, J = 15.9 Hz, 1 H), 3.70 (s, 3 H), 3.17 - 3.07 (m, 4 H). **$^{13}\text{C}\{^1\text{H}\}$ NMR** (101MHz, $\text{CHLOROFORM-}d$): δ ppm 196.3, 171.3, 146.1, 137.3, 135.7, 131.5, 128.5, 127.9, 127.3, 126.4, 126.0, 121.2, 52.8, 52.7, 43.6, 37.6. **HRMS (ESI-TOF) m/z** : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{21}\text{H}_{20}\text{O}_3^{81}\text{Br}$ 401.0570; Found 401.0562.

Methyl (1s,3s)-1-formyl-3-((E)-4-methylstyryl)-3-phenylcyclobutane-1-carboxylate (4h)




Colorless oil, 68% (36 mg) isolated yield, $R_f = 0.3$ in 10% EtOAc: Pet Ether. **^1H NMR** (400MHz, CHLOROFORM-*d*): δ ppm 9.82 (s, 1 H), 7.37 - 7.32 (m, 2 H), 7.25 - 7.21 (m, 3 H), 7.17 (d, $J = 8.0$ Hz, 2 H), 7.06 (d, $J = 8.0$ Hz, 2 H), 6.32 (d, $J = 15.9$ Hz, 1 H), 6.06 (d, $J = 16.0$ Hz, 1 H), 3.68 (s, 3 H), 3.11 (s, 4 H), 2.30 (s, 3 H). **$^{13}\text{C}\{^1\text{H}\}$ NMR** (101MHz, CHLOROFORM-*d*): δ ppm 196.4, 171.4, 146.5, 137.3, 135.4, 133.9, 129.2, 128.4, 128.3, 126.3, 126.2, 126.1, 52.8, 52.7, 43.6, 37.8, 21.1. **HRMS (ESI-TOF) m/z :** $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{22}\text{H}_{23}\text{O}_3$ 335.1642; Found 335.1637.

Methyl (1*s*,3*s*)-1-formyl-3-((*E*)-3-methylstyryl)-3-phenylcyclobutane-1-carboxylate (4i)




Yellow oil, 73% (39 mg) isolated yield, $R_f = 0.35$ in 10% EtOAc: Pet Ether. **^1H NMR** (400MHz, CHLOROFORM-*d*): δ ppm 9.84 (s, 1 H), 7.40 - 7.34 (m, 2 H), 7.27 (s, 1 H), 7.27 - 7.21 (m, 2 H), 7.20 - 7.15 (m, 1 H), 7.14 - 7.08 (m, 2 H), 7.02 (d, $J = 7.4$ Hz, 1 H), 6.38 (d, $J = 16.0$ Hz, 1 H), 6.09 (d, $J = 16.0$ Hz, 1 H), 3.71 (s, 3 H), 3.20 - 3.09 (m, 4 H), 2.32 (s, 3 H). **$^{13}\text{C}\{^1\text{H}\}$ NMR** (101MHz, CHLOROFORM-*d*): δ ppm 196.4, 171.3, 146.4, 138.0, 136.6, 136.2, 128.5, 128.5, 128.4, 128.2, 127.0, 126.3, 126.0, 123.5, 52.8, 52.7, 43.6, 37.7, 21.3. **HRMS (ESI-TOF) m/z :** $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{22}\text{H}_{23}\text{O}_3$ 335.1642; Found 335.1637.

Methyl 4-((*E*)-2-((1*s*,3*s*)-3-formyl-3-(methoxycarbonyl)-1-phenylcyclobutyl)vinyl) benzoate (4j)




Yellow oil, 70% (42 mg) isolated yield, $R_f = 0.26$ in 20% EtOAc: Pet Ether. **^1H NMR** (400MHz, CHLOROFORM-*d*): δ ppm 9.84 (s, 1 H), 7.93 (d, $J = 8.4$ Hz, 2 H), 7.40 - 7.36 (m, 2 H), 7.34 (d, $J = 8.4$ Hz, 2 H), 7.25 (d, $J = 8.0$ Hz, 3 H), 6.51 (d, $J = 16.0$ Hz, 1 H), 6.11 (d, $J = 16.0$ Hz, 1 H), 3.90 (s, 3 H), 3.70 (s, 3 H), 3.18 - 3.10 (m, 4 H). **$^{13}\text{C}\{^1\text{H}\}$ NMR** (101MHz, CHLOROFORM-*d*): δ ppm 196.2, 171.3, 166.9, 145.9, 141.2, 139.2, 129.8, 128.9, 128.6, 127.6, 126.5, 126.2, 126.1, 52.8, 52.7, 52.0, 43.7, 37.5. **HRMS (ESI-TOF) m/z :** $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{23}\text{H}_{23}\text{O}_5$ 379.1540; Found 379.1524.

Methyl (*E*)-3-(4-cyanostyryl)-1-phenyl-2-oxabicyclo[2.1.1]hexane-4-carboxylate (3k)




Colourless oil, 85% (47 mg) isolated yield, R_f = 0.5 in 20% EtOAc: Pet Ether. **^1H NMR** (400MHz, CHCl_3 - d): δ ppm 7.61 (d, J = 8.4 Hz, 2 H), 7.52 - 7.44 (m, 4 H), 7.44 - 7.35 (m, 3 H), 6.84 (d, J = 15.9 Hz, 1 H), 6.53 (dd, J = 5.5, 15.9 Hz, 1 H), 4.98 (dd, J = 1.1, 5.6 Hz, 1 H), 3.78 (s, 3 H), 2.56 (d, J = 7.1 Hz, 1 H), 2.46 - 2.39 (m, 2 H), 2.34 - 2.30 (m, 1 H). **$^{13}\text{C}\{^1\text{H}\}$ NMR** (101MHz, CHCl_3 - d): δ ppm 169.9, 140.9, 136.6, 132.3, 131.9, 130.8, 128.4, 128.4, 127.1, 126.0, 118.8, 110.9, 86.9, 79.0, 53.2, 52.0, 49.7, 41.6. **HRMS (ESI-TOF) m/z** : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{23}\text{H}_{23}\text{O}_5$ 346.1438; Found 346.1426.

Methyl (*E*)-3-(4-nitrostyryl)-1-phenyl-2-oxabicyclo[2.1.1]hexane-4-carboxylate (3l)



Yellow oil, 93% (54 mg) isolated yield, R_f = 0.5 in 20% EtOAc: Pet Ether. **^1H NMR** (400MHz, CHCl_3 - d): δ ppm 8.28 - 8.10 (m, J = 8.6 Hz, 2 H), 7.58 - 7.51 (m, J = 8.8 Hz, 2 H), 7.50 - 7.44 (m, 2 H), 7.44 - 7.34 (m, 3 H), 6.90 (d, J = 15.9 Hz, 1 H), 6.60 (dd, J = 5.4, 15.8 Hz, 1 H), 3.79 (s, 3 H), 2.57 (d, J = 7.1 Hz, 1 H), 2.49 - 2.40 (m, 2 H), 2.36 - 2.30 (m, 1 H). **$^{13}\text{C}\{^1\text{H}\}$ NMR** (101MHz, CHCl_3 - d): δ ppm 169.9, 147.0, 142.9, 136.6, 131.8, 131.5, 128.4, 128.4, 127.2, 126.0, 123.9, 87.0, 79.0, 53.2, 52.1, 49.7, 41.7. **HRMS (ESI-TOF) m/z** : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{21}\text{H}_{20}\text{O}_5\text{N}$ 366.1336; Found 366.1324.


Methyl (1*s*,3*s*)-1-formyl-3-((*E*)-2-methoxystyryl)-3-phenylcyclobutane-1-carboxylate (4m)



Yellow oil, 66% (35 mg) isolated yield, R_f = 0.23 in 10% EtOAc: Pet Ether. **^1H NMR** (400MHz, CHCl_3 - d): δ ppm 9.84 (s, 1 H), 7.39 - 7.32 (m, 3 H), 7.29 (d, J = 1.4 Hz, 2 H), 7.25 - 7.16


(m, 2 H), 6.91 - 6.80 (m, 2 H), 6.56 (d, $J = 16.1$ Hz, 1 H), 6.36 (d, $J = 16.1$ Hz, 1 H), 3.81 (s, 3 H), 3.70 (s, 3 H), 3.19 - 3.10 (m, 4 H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101MHz, CHLOROFORM- d): δ ppm 196.5, 171.4, 156.6, 146.7, 136.6, 128.5, 128.5, 126.6, 126.2, 126.0, 125.8, 123.1, 120.6, 110.9, 55.5, 52.9, 52.7, 44.0, 37.9. HRMS (ESI-TOF) m/z : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{22}\text{H}_{23}\text{O}_4$ 351.1591; Found 351.1586.

Methyl (1*s*,3*s*)-1-formyl-3-((*E*)-2-(naphthalen-2-yl)vinyl)-3-phenylcyclobutane-1-carboxylate (4n)




Yellow solid, mp 125-127 °C, 49% (29 mg) isolated yield, $R_f = 0.5$ in 10% EtOAc: Pet Ether. ^1H NMR (400MHz, CHLOROFORM- d): δ ppm 9.88 (s, 1 H), 7.79 - 7.73 (m, 3 H), 7.64 (s, 1 H), 7.53 (dd, $J = 1.6, 8.6$ Hz, 1 H), 7.46 - 7.38 (m, 4 H), 7.33 - 7.27 (m, 3 H), 6.53 (d, $J = 16.0$ Hz, 1 H), 6.26 (d, $J = 15.9$ Hz, 1 H), 3.71 (s, 3 H), 3.19 (s, 4 H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101MHz, CHLOROFORM- d): δ ppm 196.4, 171.4, 146.3, 136.9, 134.1, 133.5, 132.9, 128.6, 128.5, 128.1, 127.9, 127.6, 126.4, 126.2, 126.2, 126.1, 125.7, 123.5, 52.8, 52.7, 43.7, 37.8. HRMS (ESI-TOF) m/z : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{22}\text{H}_{23}\text{O}_4$ 351.1591; Found 351.1586.

Methyl (1*s*,3*s*)-1-formyl-3-phenyl-3-((*E*)-2-(thiophen-2-yl)vinyl)cyclobutane-1-carboxylate (4o)




Yellow oil, 48% (25 mg) isolated yield, $R_f = 0.2$ in 10% EtOAc: Pet Ether. ^1H NMR (400MHz, CHLOROFORM- d): δ ppm 9.83 (s, 1 H), 7.39 - 7.35 (m, 2 H), 7.27 - 7.23 (m, 3 H), 7.11 (d, $J = 4.9$ Hz, 1 H), 6.93 - 6.89 (m, 1 H), 6.86 (d, $J = 2.4$ Hz, 1 H), 6.29 - 6.15 (m, 2 H), 3.69 (s, 3 H), 3.17 - 3.05 (m, 4 H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101MHz, CHLOROFORM- d): δ ppm 196.3, 171.3, 146.0, 141.8, 136.2, 128.5, 127.3, 126.4, 126.1, 125.6, 124.0, 122.0, 52.7 (2C), 43.5, 37.7. HRMS (ESI-TOF) m/z : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{19}\text{H}_{19}\text{O}_3\text{S}$ 327.1049; Found 351.1045

Methyl (*E*)-3-(2-cyclohexylvinyl)-1-phenyl-2-oxabicyclo[2.1.1]hexane-4-carboxylate (3p)




Colorless oil, 33% (17 mg) isolated yield, R_f = 0.43 in 20% EtOAc: Pet Ether. $^1\text{H NMR}$ (400MHz, $\text{CHLOROFORM-}d$): δ ppm 7.47 (d, J = 7.1 Hz, 2 H), 7.38 (t, J = 7.6 Hz, 2 H), 7.31 - 7.28 (m, 1 H), 6.54 (dd, J = 10.8, 15.2 Hz, 1 H), 5.97 (d, J = 15.3 Hz, 1 H), 5.85 (d, J = 10.8 Hz, 1 H), 3.61 (s, 3 H), 3.28 - 3.22 (m, 2 H), 2.68 - 2.62 (m, 2 H), 2.34 - 2.28 (m, 2 H), 2.20 - 2.09 (m, 3 H), 1.58 (br. s., 6 H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101MHz, $\text{CHLOROFORM-}d$): δ ppm 175.5, 145.2, 144.3, 132.2, 128.5, 127.5, 126.2, 125.1, 121.1, 73.0, 52.2, 46.0 (2C), 42.8, 37.3, 29.3, 28.4, 27.6, 26.7. **HRMS (ESI-TOF) m/z :** $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{21}\text{H}_{27}\text{O}_3$ 327.1955; Found 327.1943.

***Tert*-butyl (1*s*,3*s*)-3-((*E*)-4-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4q)**



Yellow oil, 74% (47 mg) isolated yield, R_f = 0.4 in 5% EtOAc: Pet Ether. $^1\text{H NMR}$ (400MHz, $\text{CHLOROFORM-}d$): δ ppm 9.79 (s, 1 H), 7.39 - 7.35 (m, 2 H), 7.28 (br. s., 1 H), 7.25 (d, J = 5.9 Hz, 2 H), 7.23 - 7.19 (m, 4 H), 6.36 (d, J = 15.9 Hz, 1 H), 5.99 (d, J = 15.9 Hz, 1 H), 3.10 - 3.01 (m, 4 H), 1.35 (s, 9 H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101MHz, $\text{CHLOROFORM-}d$): δ ppm 197.1, 170.0, 146.0, 137.8, 135.3, 133.0, 128.6, 128.5, 127.5, 127.1, 126.4, 126.3, 82.5, 53.5, 43.5, 37.8, 27.8. **HRMS (ESI-TOF) m/z :** $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{24}\text{H}_{25}\text{O}_3^{35}\text{Cl}$ 397.1565; Found 397.1561.


Benzyl (1*s*,3*s*)-3-((*E*)-4-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4r)



Colorless oil, 79% (54 mg) isolated yield, R_f = 0.6 in 10% EtOAc: Pet Ether. $^1\text{H NMR}$ (400MHz, $\text{CHLOROFORM-}d$): δ ppm 9.76 (s, 1 H), 7.30 - 7.22 (m, 5 H), 7.19 - 7.14 (m, 4 H), 7.14 - 7.11 (m, 5 H), 6.29 (d, J = 16.0 Hz, 1 H), 5.93 (d, J = 15.9 Hz, 1 H), 5.03 (s, 2 H), 3.09 - 3.00 (m, 4 H).

$^{13}\text{C}\{^1\text{H}\}$ NMR (101MHz, CHLOROFORM-*d*): δ ppm 196.3, 170.7, 145.9, 137.3, 135.2, 135.0, 133.0, 128.6 (2C), 128.5, 128.4, 128.0, 127.5, 127.2, 126.4, 126.1, 67.4, 52.7, 43.5, 37.7. **HRMS (ESI-TOF) m/z:** $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{27}\text{H}_{24}\text{O}_3^{35}\text{Cl}$ 431.1408; Found 431.1406.


(1*s*,3*s*)-3-((*E*)-4-chlorostyryl)-1-formyl-*N*-methoxy-*N*-methyl-3-phenylcyclobutane-1-carboxamide (4*s*)



Yellow solid, mp 102-104 °C, 49% (30 mg) isolated yield, R_f = 0.4 in 30% EtOAc: Pet Ether. **^1H NMR** (400MHz, CHLOROFORM-*d*): δ ppm 9.77 (s, 1 H), 7.38 - 7.32 (m, 3 H), 7.24 (s, 5 H), 7.22 (s, 1 H), 6.47 (d, J = 16.0 Hz, 1 H), 6.11 (d, J = 15.9 Hz, 1 H), 3.59 (s, 3 H), 3.24 - 3.19 (m, 2 H), 3.18 (s, 3 H), 3.10 - 3.05 (m, 2 H). **$^{13}\text{C}\{^1\text{H}\}$ NMR** (101MHz, CHLOROFORM-*d*): δ ppm 196.4, 146.9, 137.4, 135.3, 133.0, 128.6, 128.5, 128.3, 127.5, 127.1, 126.3, 125.9, 61.2, 52.9, 43.2, 37.7, 32.9. **HRMS (ESI-TOF) m/z:** $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{22}\text{H}_{23}\text{O}_3\text{N}^{35}\text{Cl}$ 384.1361; Found 384.1353.


(The reaction mixture was stirred at 50 °C for 10 h)

(*E*)-(3-(4-chlorostyryl)-1-phenyl-2-oxabicyclo[2.1.1]hexan-4-yl)(naphthalen-2-yl) methanone (3*t*)




White solid, mp 140-142 °C, 49% (35 mg) isolated yield, R_f = 0.4 in 40% EtOAc: Pet Ether. **^1H NMR** (400MHz, CHLOROFORM-*d*): δ ppm 8.35 (s, 1 H), 7.96 - 7.86 (m, 4 H), 7.67 - 7.53 (m, 2 H), 7.44 (d, J = 7.4 Hz, 2 H), 7.32 (t, J = 7.6 Hz, 2 H), 7.25 - 7.20 (m, 3 H), 7.17 - 7.12 (m, 2 H), 6.35 (d, J = 15.8 Hz, 1 H), 6.06 (dd, J = 6.8, 15.8 Hz, 1 H), 5.17 (d, J = 6.8 Hz, 1 H), 3.59 (d, J = 13.9 Hz, 1 H), 3.38 - 3.27 (m, 2 H), 2.84 (dd, J = 3.5, 14.0 Hz, 1 H). **$^{13}\text{C}\{^1\text{H}\}$ NMR** (101MHz, CHLOROFORM-*d*): δ ppm 200.9, 145.3, 135.4, 134.4, 133.7, 132.3, 132.0, 131.6, 131.0, 129.6, 128.7, 128.7, 128.5, 128.4, 127.9, 127.8, 127.5, 127.2, 126.9, 125.0, 124.8, 74.8, 72.3, 51.4, 47.0, 40.4. **HRMS (ESI-TOF) m/z:** $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{30}\text{H}_{24}\text{O}_2^{35}\text{Cl}$ 451.1459; Found 451.1443.

Methyl (1*s*,3*s*)-3-((*E*)-4-chlorostyryl)-3-(4-fluorophenyl)-1-formylcyclobutane-1-carboxylate (4u)




Yellow oil, 79% (47 mg) isolated yield, $R_f = 0.5$ in 20% EtOAc: Pet Ether. $^1\text{H NMR}$ (400MHz, $\text{CHLOROFORM-}d$): δ ppm 9.83 (s, 1 H), 7.26 - 7.19 (m, 6 H), 7.08 - 7.03 (m, 2 H), 6.35 (d, $J = 15.9$ Hz, 1 H), 5.99 (d, $J = 15.9$ Hz, 1 H), 3.71 (s, 3 H), 3.09 (s, 4 H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101MHz, $\text{CHLOROFORM-}d$): δ ppm 196.1, 171.3, 161.3 (d, $J_{\text{C-F}} = 246.4$ Hz), 141.7 (d, $J_{\text{C-F}} = 3.8$ Hz), 137.1, 135.0, 133.1, 128.6, 127.8 (d, $J_{\text{C-F}} = 7.6$ Hz), 127.5, 127.4, 115.4, 115.2, 52.8, 52.4, 43.1, 37.7. **HRMS (ESI-TOF) m/z:** $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{21}\text{H}_{19}\text{O}_3^{35}\text{ClF}$ 373.1001; Found 373.0997.

Methyl (1*s*,3*s*)-3-(4-chlorophenyl)-3-((*E*)-4-chlorostyryl)-1-formylcyclobutane-1-carboxylate (4v)




Yellow oil, 56% (35 mg) isolated yield, $R_f = 0.4$ in 20% EtOAc: Pet Ether. $^1\text{H NMR}$ (400MHz, $\text{CHLOROFORM-}d$): δ ppm 9.83 (s, 1 H), 7.34 (d, $J = 8.5$ Hz, 2 H), 7.25 - 7.20 (m, 4 H), 7.19 - 7.17 (m, 2 H), 6.34 (d, $J = 16.0$ Hz, 1 H), 5.99 (d, $J = 16.0$ Hz, 1 H), 3.71 (s, 3 H), 3.09 (s, 4 H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101MHz, $\text{CHLOROFORM-}d$): δ ppm 196.1, 171.2, 144.5, 136.7, 135.0, 133.2, 132.3, 128.7, 128.6, 127.6 (2C), 127.5, 52.9, 52.4, 43.1, 37.6. **HRMS (ESI-TOF) m/z:** $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{21}\text{H}_{19}\text{O}_3^{35}\text{Cl}_2$ 389.0706; Found 389.0700.

Methyl(1*s*,3*s*)-3-(4-bromophenyl)-3-((*E*)-4-chlorostyryl)-1-formylcyclobutane-1-carboxylate (4w)




Yellow oil, 81% (56 mg) isolated yield, $R_f = 0.3$ in 20% EtOAc: Pet Ether. **^1H NMR** (400MHz, CHLOROFORM- d): δ ppm 9.82 (s, 1 H), 7.54 - 7.47 (m, $J = 8.5$ Hz, 2 H), 7.25 - 7.18 (m, 4 H), 7.15 - 7.10 (m, 2 H), 6.33 (d, $J = 16.0$ Hz, 1 H), 5.99 (d, $J = 16.0$ Hz, 1 H), 3.71 (s, 3 H), 3.08 (s, 4 H). **$^{13}\text{C}\{^1\text{H}\}$ NMR** (101MHz, CHLOROFORM- d): δ ppm 196.0, 171.2, 145.1, 136.6, 135.0, 133.2, 131.6, 128.6, 128.0, 127.6, 127.5, 120.4, 52.8, 52.4, 43.2, 37.6. **HRMS (ESI-TOF) m/z :** $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{21}\text{H}_{19}\text{O}_3^{35}\text{Cl}^{81}\text{Br}$ 435.0180; Found 435.0173.

Methyl (1s,3s)-3-(4-(*tert*-butyl)phenyl)-3-((*E*)-4-chlorostyryl)-1-formylcyclobutane-1-carboxylate (4x)



White solid, mp 124-126 °C, 93% (61 mg) isolated yield, $R_f = 0.32$ in 10% EtOAc: Pet Ether. **^1H NMR** (400MHz, CHLOROFORM- d): δ ppm 9.82 (s, 1 H), 7.38 (d, $J = 8.4$ Hz, 2 H), 7.23 (s, 4 H), 7.17 (d, $J = 8.5$ Hz, 2 H), 6.35 (d, $J = 16.0$ Hz, 1 H), 6.10 (d, $J = 16.0$ Hz, 1 H), 3.70 (s, 3 H), 3.15 - 3.07 (m, 4 H), 1.33 (s, 9 H). **$^{13}\text{C}\{^1\text{H}\}$ NMR** (101MHz, CHLOROFORM- d): δ ppm 196.3, 171.3, 149.2, 143.1, 137.1, 135.3, 133.0, 128.6, 127.5, 127.0, 125.6, 125.4, 52.8, 52.7, 43.2, 37.6, 34.4, 31.3. **HRMS (ESI-TOF) m/z :** $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{25}\text{H}_{28}\text{O}_3^{35}\text{Cl}$ 411.1721; Found 411.1719.


Methyl (1s,3s)-3-((*E*)-4-chlorostyryl)-1-formyl-3-(4-(trifluoromethyl)phenyl)cyclobutane-1-carboxylate (4y)



Yellow oil, 42% (28 mg) isolated yield, $R_f = 0.5$ in 20% EtOAc: Pet Ether. **^1H NMR** (400MHz, CHLOROFORM- d): δ ppm 9.84 (s, 1 H), 7.66 - 7.62 (m, $J = 8.4$ Hz, 2 H), 7.38 - 7.35 (m, $J = 8.3$ Hz, 2 H), 7.22 (q, $J = 8.6$ Hz, 4 H), 6.34 (d, $J = 16.0$ Hz, 1 H), 6.01 (d, $J = 16.0$ Hz, 1 H), 3.72 (s, 3 H), 3.17 - 3.09 (m, 4 H). **$^{13}\text{C}\{^1\text{H}\}$ NMR** (101MHz, CHLOROFORM- d): δ ppm 196.9, 171.2, 150.2, 136.2, 134.8, 133.4, 128.8 (q, $J_{\text{C-F}} = 32.4$ Hz), 128.7, 128.0, 127.6, 126.5, 125.6 (q, $J_{\text{C-F}} = 3.8$ Hz),


124.1 (q, $J_{C-F} = 272.4$ Hz), 52.9, 52.4, 43.5, 37.5. **HRMS (ESI-TOF) m/z:** $[M+H]^+$ Calcd for $C_{22}H_{19}^{35}ClF_3O_3$ 423.0969; Found 423.0969.

Methyl (1*s*,3*s*)-3-((*E*)-4-chlorostyryl)-1-formyl-3-(3-methoxyphenyl)cyclobutane-1-carboxylate (4z)




Colorless oil, 82% (50 mg) isolated yield, $R_f = 0.5$ in 20% EtOAc: Pet Ether. **1H NMR** (400MHz, $CHCl_3$ - d): δ ppm 9.79 (s, 1 H), 7.26 - 7.22 (m, 1 H), 7.22 - 7.15 (m, 4 H), 6.79 (d, $J = 7.6$ Hz, 1 H), 6.77 - 6.71 (m, 2 H), 6.31 (d, $J = 15.9$ Hz, 1 H), 6.02 (d, $J = 16.0$ Hz, 1 H), 3.78 (s, 3 H), 3.67 (s, 3 H), 3.15 - 3.00 (m, 4 H). **$^{13}C\{^1H\}$ NMR** (101MHz, $CHCl_3$ - d): δ ppm 196.3, 171.3, 159.7, 147.9, 136.9, 135.2, 133.0, 129.6, 128.6, 127.5, 127.3, 118.4, 112.4, 111.3, 55.2, 52.8, 52.6, 43.6, 37.6. **HRMS (ESI-TOF) m/z:** $[M+H]^+$ Calcd for $C_{22}H_{22}O_4^{35}Cl$ 385.1201; Found 385.1196.

Methyl(1*s*,3*s*)-3-([1,1'-biphenyl]-4-yl)-3-((*E*)-4-chlorostyryl)-1-formylcyclobutane-1-carboxylate (4aa)




White solid, mp 104-106 °C, 93% (64 mg) isolated yield, $R_f = 0.24$ in 10% EtOAc: Pet Ether. **1H NMR** (400MHz, $CHCl_3$ - d): δ ppm 9.86 (s, 1 H), 7.62 (d, $J = 8.1$ Hz, 4 H), 7.46 (t, $J = 7.6$ Hz, 2 H), 7.40 - 7.32 (m, 3 H), 7.25 (s, 4 H), 6.41 (d, $J = 16.0$ Hz, 1 H), 6.11 (d, $J = 16.0$ Hz, 1 H), 3.73 (s, 3 H), 3.23 - 3.11 (m, 4 H). **$^{13}C\{^1H\}$ NMR** (101MHz, $CHCl_3$ - d): δ ppm 196.2, 171.3, 145.1, 140.6, 139.3, 137.0, 135.2, 133.0, 128.8, 128.6, 127.5, 127.3, 127.3, 127.2, 127.0, 126.5, 52.8, 52.7, 43.4, 37.6. **HRMS (ESI-TOF) m/z:** $[M+H]^+$ Calcd for $C_{27}H_{24}^{35}ClO_3$ 431.1408; Found 431.1405.

Methyl (1*s*,3*s*)-3-((*E*)-4-chlorostyryl)-1-formyl-3-(naphthalen-1-yl)cyclobutane-1-carboxylate (4ab)




Yellow oil, 74% (48 mg) isolated yield, R_f = 0.5 in 20% EtOAc: Pet Ether. ^1H NMR (400MHz, CHLOROFORM-*d*): δ ppm 9.97 (s, 1 H), 7.92 - 7.88 (m, 1 H), 7.81 (d, J = 8.1 Hz, 1 H), 7.68 (d, J = 8.3 Hz, 1 H), 7.54 - 7.49 (m, 1 H), 7.48 - 7.41 (m, 3 H), 7.20 - 7.12 (m, 4 H), 6.67 (d, J = 16.0 Hz, 1 H), 5.94 (d, J = 15.9 Hz, 1 H), 3.69 (s, 3 H), 3.42 - 3.34 (m, 4 H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101MHz, CHLOROFORM-*d*): δ ppm 196.6, 171.2, 141.8, 136.7, 135.2, 134.4, 132.9, 130.3, 129.1, 128.5, 128.1, 127.8, 127.5, 125.6, 125.5(2C), 125.1, 124.3, 52.9, 52.8, 43.9, 39.0. HRMS (ESI-TOF) m/z : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{25}\text{H}_{22}\text{O}_3^{35}\text{Cl}$ 405.1252; Found 405.1248.

(4*R*,8*R*,9*R*,10*S*,13*S*,17*S*)-10,13-dimethyl-17-((*S*)-6-methylheptan-2-yl)-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1*H*-cyclopenta[*a*]phenanthren-4-yl (1*s*,3*S*)-3-((*E*)-4-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4ac)




White solid, mp 164-166 °C, 94% (106 mg) isolated yield, R_f = 0.44 in 5% EtOAc: Pet Ether. ^1H NMR (400MHz, CHLOROFORM-*d*): δ ppm 9.83 (s, 1 H), 7.40 - 7.34 (m, 2 H), 7.27 (s, 1 H), 7.26 - 7.15 (m, 6 H), 6.38 (d, J = 16.0 Hz, 1 H), 6.01 (d, J = 15.9 Hz, 1 H), 5.34 (d, J = 4.4 Hz, 1 H), 4.63 - 4.54 (m, 1 H), 3.14 - 3.05 (m, 4 H), 2.18 (d, J = 7.1 Hz, 2 H), 2.04 - 1.93 (m, 2 H), 1.87 - 1.79 (m, 2 H), 1.76 - 1.70 (m, 1 H), 1.57 - 1.31 (m, 11 H), 1.19 - 0.99 (m, 10 H), 0.98 (s, 3 H), 0.92 (d, J = 6.5 Hz, 3 H), 0.88 (d, J = 1.5 Hz, 3 H), 0.87 (d, J = 1.5 Hz, 3 H), 0.68 (s, 3 H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101MHz, CHLOROFORM-*d*): δ ppm 196.8, 170.4, 146.0, 139.2, 137.6, 135.3, 133.0, 128.6, 128.5, 127.5, 127.1, 126.4, 126.3, 122.9, 75.5, 56.6, 56.1, 52.8, 49.9, 43.5, 42.3, 39.7, 39.5, 37.8, 37.7, 36.8, 36.5, 36.2, 35.8, 31.9, 31.8, 28.2, 28.0, 27.4, 24.2, 23.8, 22.8, 22.5, 21.0, 19.2, 18.7, 11.8.

**(1*S*,2*R*,4*R*)-2-isopropyl-4-methylcyclohexyl
cyclobutane-1-carboxylate (4ad)**



Yellow oil, 90% (69 mg) isolated yield, $R_f = 0.6$ in 10% EtOAc: Pet Ether. ^1H NMR (400MHz, CHLOROFORM-*d*): δ ppm 9.83 (s, 1 H), 7.40 - 7.35 (m, 2 H), 7.28 (br. s., 1 H), 7.26 - 7.24 (m, 2 H), 7.22 (d, $J = 3.8$ Hz, 4 H), 6.38 (d, $J = 15.9$ Hz, 1 H), 6.01 (d, $J = 15.9$ Hz, 1 H), 4.70-4.63 (m, 1 H), 3.16 - 3.06 (m, 4 H), 1.83 (d, $J = 11.6$ Hz, 1 H), 1.71 - 1.65 (m, 2 H), 1.49 - 1.39 (m, 1 H), 1.37 - 1.23 (m, 2 H), 1.06 - 0.96 (m, 1 H), 0.93 - 0.89 (m, 2 H), 0.88 (d, $J = 6.6$ Hz, 3 H), 0.83 (d, $J = 7.0$ Hz, 3 H), 0.66 (d, $J = 6.9$ Hz, 3 H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101MHz, CHLOROFORM-*d*): δ ppm 196.7, 170.4, 145.8, 137.7, 135.2, 133.0, 128.6, 128.5, 127.5, 127.1, 126.4, 126.2, 75.9, 52.9, 46.7, 43.5, 40.4, 37.7, 34.0, 31.3, 26.1, 23.1, 21.9, 20.7, 15.9. HRMS (ESI-TOF) m/z : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{30}\text{H}_{36}^{35}\text{ClO}_3$ 479.2347; Found 479.2347.


**5-(2,4-dimethylphenoxy)-2,2-dimethylpentyl
cyclobutane-1-carboxylate (4ae)**



Yellow oil, 45% (40 mg) isolated yield, $R_f = 0.29$ in 10% EtOAc: Pet Ether. ^1H NMR (400MHz, CHLOROFORM-*d*): δ ppm 9.87 (s, 1 H), 7.39 - 7.34 (m, 2 H), 7.27 - 7.20 (m, 7 H), 7.03 (d, $J = 7.4$ Hz, 1 H), 6.69 (d, $J = 7.5$ Hz, 1 H), 6.63 (s, 1 H), 6.39 (d, $J = 16.0$ Hz, 1 H), 6.02 (d, $J = 16.0$ Hz, 1 H), 3.91 - 3.85 (m, 4 H), 3.19 - 3.09 (m, 4 H), 2.34 (s, 3 H), 2.20 (s, 3 H), 1.75 - 1.67 (m, 2 H), 1.38 - 1.33 (m, 2 H), 0.88 (s, 6 H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101MHz, CHLOROFORM-*d*): δ ppm 196.3, 170.9, 156.9, 146.1, 137.4, 136.4, 135.2, 133.0, 130.3, 128.6, 128.5, 127.5, 127.2, 126.4, 126.1, 123.5,


120.7, 111.9, 73.4, 68.1, 52.6, 43.5, 37.6, 35.2, 33.7, 24.0, 21.4, 15.8. **HRMS (ESI-TOF) m/z:** $[M+H]^+$ Calcd for $C_{35}H_{39}^{35}ClO_4$ 559.2610; Found 559.2589.

Methyl (1*r*,3*s*)-3-(4-(*tert*-butyl)phenyl)-3-(4-chlorophenethyl)-1-formylcyclobutane-1-carboxylate (5)



Colourless oil, 85% (85 mg) isolated yield, R_f = 0.5 in 10% EtOAc: Pet Ether. **1H NMR** (400MHz, $CHCl_3$ - d): δ ppm 9.84 (s, 1 H), 7.38 (d, J = 8.4 Hz, 2 H), 7.21 - 7.17 (m, J = 8.4 Hz, 2 H), 7.14 - 7.10 (m, J = 8.5 Hz, 2 H), 6.97 (d, J = 8.4 Hz, 2 H), 3.69 (s, 3 H), 2.94 (d, J = 13.1 Hz, 2 H), 2.80 (d, J = 13.3 Hz, 2 H), 2.28 - 2.21 (m, 2 H), 1.96 - 1.89 (m, 2 H), 1.35 (s, 9 H). **$^{13}C\{^1H\}$ NMR** (101MHz, $CHCl_3$ - d): δ ppm 196.6, 171.5, 148.8, 143.9, 140.3, 131.3, 129.6, 128.3, 125.3, 125.1, 52.6, 52.6, 44.7, 40.8, 36.9, 34.3, 31.3, 30.0. **HRMS (ESI-TOF) m/z:** $[M+H]^+$ Calcd for $C_{25}H_{30}^{35}ClO_3$ 413.1878; Found 413.1870.

Methyl (1*s*,3*s*)-3-((*E*)-4-chlorostyryl)-1-(hydroxymethyl)-3-phenylcyclobutane-1-carboxylate (6)



Yellow oil, 96% (29 mg) isolated yield, R_f = 0.4 in 40% EtOAc: Pet Ether. **1H NMR** (400MHz, $CHCl_3$ - d): δ ppm 7.40 - 7.34 (m, 2 H), 7.29 (s, 4 H), 7.27 (s, 3 H), 7.25 - 7.21 (m, 1 H), 6.46 (d, J = 16.0 Hz, 1 H), 6.29 (d, J = 16.0 Hz, 1 H), 3.97 (s, 2 H), 3.69 (s, 3 H), 3.07 (d, J = 13.3 Hz, 2 H), 2.68 (d, J = 13.3 Hz, 2 H), 2.36 (br. s., 1 H). **$^{13}C\{^1H\}$ NMR** (101MHz, $CHCl_3$ - d): δ ppm 176.4, 147.7, 138.9, 135.4, 132.9, 128.6, 128.5, 127.5, 126.2, 126.1, 125.9, 67.0, 52.2, 43.2 (2C), 38.2. **HRMS (ESI-TOF) m/z:** $[M+Na]^+$ Calcd for $C_{21}H_{21}^{35}ClO_3Na$ 379.1071; Found 379.1069.

7. References:

- 1) Gately, T. J., Perry, C. J., Weiss, S., Lam, K., Islam, I., Almtiri, M. N., Carta, V., Beran, G. J. O., Al-Kaysi, R. O., & Bardeen, C. J. Using chemical substitution to engineer photomechanical cinnamalmalononitrile crystals. *Crystal Growth & Design* **24**, 9544–9555 (2024).
- 2) Putatunda, S., Alegre-Requena, J. V., Meazza, M., Franc, M., Rohal'ová, D., Vemuri, P., Císařová, I., Herrera, R. P., Rios, R., & Veselý, J.. Proline bulky substituents consecutively act as steric hindrances and directing groups in a Michael/Conia-ene cascade reaction under synergistic catalysis. *Chem. Sci.* **10**, 4107–4115 (2019).
- 3) Nie, R., Shen, X., Wen, Y., & Li, S. Cobalt-Catalyzed Coupling of Conjugated Dienes and Epoxides with CO. *Org. Lett.* **27**, 6238–6243 (2025).
- 4) Deswal, S., Das, R. C., Sarkar, D., & Biju, A. T. Simultaneous Activation of Bicyclobutanes and Indolyl Alcohols with HFIP: Access to Indole-Fused Bicyclo[3.1.1]Heptanes. *Angew. Chem. Int. Ed.* **64**, e202501655. (2025).
- 5) Dhake, K., Woelk, K. J., Becica, J., Un, A., Jenny, S. E., & Leitch, D. C. Beyond Bioisosteres: Divergent Synthesis of Azabicyclohexanes and Cyclobutenyl Amines from Bicyclobutanes. *Angew. Chem. Int. Ed.* **61**, e202204719. (2022).
- 6) Gao, X., Xiao, Y., Peng, Q., Yang, F., Li, D., Wang, G., Qian, Y., & Feng, J. Rapid and Modular Access to All-Carbon Quaternary Center-Containing 2,2-Disubstituted Bicyclo[1.1.1]pentanes via Cyanocarbene Addition to Bicyclo[1.1.0]butanes. *Angew. Chem. Int. Ed.* **64**, e202513768 (2025).
- 7) Yi, L., Makarov, A. S., Maity, B., Kong, D., Alshehri, R., Cavallo, L., Koenigs, R. M., & Rueping, M. Modular Access to sp³-Rich Bicyclo[1.1.1]pentane Bioisosteres via Energy-Transfer-Mediated Carbene Insertion. *Angew. Chem. Int. Ed.* **65**, e18508 (2025).
- 8) Guo, R., Chang, Y., Herter, L., Salome, C., Braley, S. E., Fessard, T. C., & Brown, M. K. Strain-Release [2 π + 2 σ] Cycloadditions for the Synthesis of Bicyclo[2.1.1]hexanes Initiated by Energy Transfer. *J. Am. Chem. Soc.* **144**, 7988–7994 (2022).
- 9) Hans, M.; Lorkowski, J.; Demonceau, A.; Delaude, L. Efficient synthetic protocols for the preparation of common N-heterocyclic carbene precursors. *Beilstein journal of organic chemistry*, **11**, 2318-2325 (2015).
- 10) Enders, D.; Breuer, K.; Kallfass, U.; Balensiefer, T. Preparation and application of 1, 3, 4-triphenyl-4, 5-dihydro-1H-1, 2, 4-triazol-5-ylidene, a stable carbene. *Synthesis*, **2003**, 1292-1295 (2003).
- 11) Lezius, L.; Reimler, J.; Döben, N.; Hamm, M.; Daniliuc, C. G. and Studer, A. Aminoacylation of Alkenes by Cooperative NHC and Photoredox Catalysis. *Synlett*, **35**, 445-450 (2024).

8. X-Ray Crystallographic Data

8.1 X-Ray Crystallographic Data of Compound 3g

A colourless, needle-shaped crystal was mounted on the goniometer. Data for mo_3g were collected from a single crystal in n/a hours at 100(2) K on a Bruker D8 VENTURE Kappa Duo PHOTON II CPAD \k-geometry diffractometer with a micro-focus sealed tube using a multilayer mirrors as monochromator and a Photon II CPAD Area Detector detector. The diffractometer used MoK α radiation ($\lambda = 0.71073$ Å). All data were integrated with SAINT (Bruker,, yielding 39429 reflections of which 4324 were independent and 90.7% were greater than $2\sigma(F_2)$.^[1] A multi-scan absorption correction using SADABS 2016/2 was applied.^[2] The structure was solved by intrinsic phasing method methods with SHELXT 2018/2 and refined by full-matrix least-squares methods against F² using SHELXL-2019/2.^[3,4] All non-hydrogen atoms were refined with anisotropic displacement parameters. All hydrogen atoms were refined isotropic on calculated positions using a riding model with their Uiso values constrained to 1.5 times the Ueq of their pivot atoms for terminal sp³ carbon atoms and 1.2 times for all other carbon atoms. Crystallographic data for the structures reported in this paper have been deposited with the Cambridge Crystallographic Data Centre.^[5] CCDC contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/structures. This report and the CIF file were generated using Final Cif.^[6]

Table 1. Crystal data and structure refinement for mo_3g

CCDC number	2588893
Empirical formula	C ₂₁ H ₁₉ BrN ₀ O ₃
Formula weight	399.27
Temperature [K]	100(2)
Crystal system	monoclinic
Space group (number)	<i>P</i> 2 ₁ / <i>c</i> (14)
<i>a</i> [Å]	16.5799(4)
<i>b</i> [Å]	6.04510(10)
<i>c</i> [Å]	17.4355(4)
α [°]	90
β [°]	95.5860(10)
γ [°]	90
Volume [Å ³]	1739.21(6)
<i>Z</i>	4
ρ_{calc} [gcm ⁻³]	1.525
μ [mm ⁻¹]	2.381
<i>F</i> (000)	816
Crystal size [mm ³]	0.12×0.14×0.33
Crystal colour	colourless
Crystal shape	needle
Radiation	MoK α ($\lambda=0.71073$ Å)

2 θ range [°]	4.94 to 56.72 (0.75 Å)
Index ranges	-21 ≤ h ≤ 22 -8 ≤ k ≤ 8 -23 ≤ l ≤ 23
Reflections collected	39429
Independent reflections	4324 $R_{\text{int}} = 0.0545$ $R_{\text{sigma}} = 0.0319$
Completeness to $\theta = 25.242^\circ$	99.6
Data / Restraints / Parameters	4324 / 0 / 227
Goodness-of-fit on F^2	1.024
Final R indexes [$I \geq 2\sigma(I)$]	$R_1 = 0.0256$ $wR_2 = 0.0629$
Final R indexes [all data]	$R_1 = 0.0298$ $wR_2 = 0.0650$
Largest peak/hole [eÅ ⁻³]	0.46/-0.39




Figure 1. ORTEP view of compound **3g** showing the atom-numbering scheme, Displacement ellipsoids are drawn at the 50% probability level and H atoms are shown as small spheres with arbitrary radii. The compound crystallizes in the monoclinic space group $P2_1/c$ containing one molecule in the asymmetric unit.

Table 2. Bond lengths and angles for mo_3g

Atom–Atom	Length [Å]
Br1–C10	1.9030(15)
O1–C3	1.4558(18)
O1–C4	1.4633(18)
O2–C20	1.2037(19)
O3–C20	1.3415(18)
O3–C21	1.4538(19)
C1–C20	1.491(2)
C1–C19	1.547(2)
C1–C4	1.552(2)
C1–C2	1.556(2)
C1–C3	2.014(2)
C2–C3	1.550(2)
C3–C13	1.490(2)
C3–C19	1.551(2)
C4–C5	1.489(2)
C5–C6	1.338(2)
C6–C7	1.466(2)
C7–C8	1.396(2)
C7–C12	1.400(2)
C8–C9	1.386(2)
C9–C10	1.382(2)
C10–C11	1.387(2)
C11–C12	1.389(2)
C13–C18	1.388(2)
C13–C14	1.396(2)
C14–C15	1.389(2)
C15–C16	1.390(2)
C16–C17	1.387(2)
C17–C18	1.397(2)
Atom–Atom–Atom	Angle [°]
C3–O1–C4	102.23(11)
C20–O3–C21	114.84(13)
C20–C1–C19	120.74(13)
C20–C1–C4	117.03(12)
C19–C1–C4	101.42(12)
C20–C1–C2	125.25(12)
C19–C1–C2	87.13(11)
C4–C1–C2	99.14(12)
C20–C1–C3	164.81(12)
C19–C1–C3	49.53(8)
C4–C1–C3	77.96(9)
C2–C1–C3	49.45(8)
C3–C2–C1	80.85(10)
O1–C3–C13	112.02(12)
O1–C3–C2	102.96(12)

C13–C3–C2	125.10(13)
O1–C3–C19	101.78(11)
C13–C3–C19	123.11(13)
C2–C3–C19	87.20(11)
O1–C3–C1	80.88(9)
C13–C3–C1	167.01(12)
C2–C3–C1	49.70(8)
C19–C3–C1	49.37(8)
O1–C4–C5	111.65(12)
O1–C4–C1	98.81(11)
C5–C4–C1	116.61(12)
C6–C5–C4	121.11(14)
C5–C6–C7	127.84(15)
C8–C7–C12	118.07(14)
C8–C7–C6	118.25(14)
C12–C7–C6	123.66(14)
C9–C8–C7	121.70(15)
C10–C9–C8	118.65(14)
C9–C10–C11	121.60(14)
C9–C10–Br1	118.54(12)
C11–C10–Br1	119.85(12)
C10–C11–C12	118.93(15)
C11–C12–C7	121.05(14)
C18–C13–C14	119.20(14)
C18–C13–C3	120.30(14)
C14–C13–C3	120.42(14)
C15–C14–C13	120.22(15)
C14–C15–C16	120.38(15)
C17–C16–C15	119.72(15)
C16–C17–C18	119.89(15)
C13–C18–C17	120.59(15)
C1–C19–C3	81.10(11)
O2–C20–O3	123.52(14)
O2–C20–C1	124.21(14)
O3–C20–C1	112.26(13)

Bonds and angles to hydrogen atoms were omitted.

Table 3. Torsion angles for mo_3g

Atom–Atom–Atom–Atom	Torsion Angle [°]
C20–C1–C2–C3	–161.92(15)
C19–C1–C2–C3	–35.67(10)
C4–C1–C2–C3	65.45(11)
C4–O1–C3–C13	179.13(12)
C4–O1–C3–C2	42.29(13)
C4–O1–C3–C19	–47.53(13)
C4–O1–C3–C1	–2.54(10)
C1–C2–C3–O1	–65.88(11)

C1–C2–C3–C13	164.92(15)
C1–C2–C3–C19	35.57(10)
C3–O1–C4–C5	126.60(13)
C3–O1–C4–C1	3.30(13)
C20–C1–C4–O1	175.07(12)
C19–C1–C4–O1	41.55(13)
C2–C1–C4–O1	–47.35(12)
C3–C1–C4–O1	–2.38(9)
C20–C1–C4–C5	55.39(18)
C19–C1–C4–C5	–78.13(15)
C2–C1–C4–C5	–167.02(13)
C3–C1–C4–C5	–122.06(13)
O1–C4–C5–C6	124.42(15)
C1–C4–C5–C6	–123.06(16)
C4–C5–C6–C7	–176.93(14)
C5–C6–C7–C8	167.01(16)
C5–C6–C7–C12	–11.2(3)
C12–C7–C8–C9	0.4(2)
C6–C7–C8–C9	–177.83(15)
C7–C8–C9–C10	–0.6(2)
C8–C9–C10–C11	0.3(2)
C8–C9–C10–Br1	–178.67(12)
C9–C10–C11–C12	0.2(2)
Br1–C10–C11–C12	179.15(12)
C10–C11–C12–C7	–0.4(2)
C8–C7–C12–C11	0.1(2)
C6–C7–C12–C11	178.25(15)
O1–C3–C13–C18	–138.99(14)
C2–C3–C13–C18	–13.5(2)
C19–C3–C13–C18	99.23(18)
C1–C3–C13–C18	48.4(6)
O1–C3–C13–C14	44.29(19)
C2–C3–C13–C14	169.73(15)
C19–C3–C13–C14	–77.49(19)
C1–C3–C13–C14	–128.3(5)
C18–C13–C14–C15	–0.8(2)
C3–C13–C14–C15	175.98(15)
C13–C14–C15–C16	0.4(2)
C14–C15–C16–C17	0.4(2)
C15–C16–C17–C18	–0.9(2)
C14–C13–C18–C17	0.4(2)
C3–C13–C18–C17	–176.40(15)
C16–C17–C18–C13	0.5(2)
C20–C1–C19–C3	165.60(14)
C4–C1–C19–C3	–63.12(11)
C2–C1–C19–C3	35.62(10)
O1–C3–C19–C1	66.89(11)
C13–C3–C19–C1	–166.72(14)

C2–C3–C19–C1	–35.77(10)
C21–O3–C20–O2	–2.3(2)
C21–O3–C20–C1	177.22(13)
C19–C1–C20–O2	38.9(2)
C4–C1–C20–O2	–85.30(19)
C2–C1–C20–O2	149.33(16)
C3–C1–C20–O2	85.1(5)
C19–C1–C20–O3	–140.62(14)
C4–C1–C20–O3	95.17(15)
C2–C1–C20–O3	–30.2(2)
C3–C1–C20–O3	–94.4(5)

Bonds and angles to hydrogen atoms were omitted.

Table 4. Hydrogen bonds for mo_3g

D–H---A [Å]	d(D–H) [Å]	d(H---A) [Å]	d(D---A) [Å]	<(DHA) [°]
C5–H5---O3 ^{#1}	0.95	2.65	3.5676(19)	161.9
C21–H21B---O2 ^{#2}	0.98	2.52	3.488(2)	170.7

Symmetry transformations used to generate equivalent atoms:

#1: +X, -1+Y, +Z; #2: +X, 1+Y, +Z;

5.2 X-Ray Crystallographic Data of Compound 4g

A colourless, cube-shaped crystal was mounted on the goniometer. Data for cu_4g were collected from a single crystal in n/a hours at 100(2) K on a Bruker D8 VENTURE Kappa Duo PHOTON II CPAD λ -geometry diffractometer with a micro-focus sealed tube using a multilayer mirrors as monochromator and a Photon II CPAD Area Detector detector. The diffractometer used CuK α radiation ($\lambda = 1.54178$ Å). All data were integrated with SAINT V8.41, yielding 261745 reflections of which 7571 were independent and 88.3% were greater than $2\sigma(F^2)$.^[1] A Multi-Scan absorption correction using TWINABS Bruker was applied.^[2] The structure was solved by Intrinsic Phasing methods with SHELXT 2018/2 and refined by full-matrix least-squares methods against F^2 using SHELXL-2019/2.^[3,4] All non-hydrogen atoms were refined with anisotropic displacement parameters. All hydrogen atoms were refined isotropic on calculated positions using a riding model with their U_{iso} values constrained to 1.5 times the U_{eq} of their pivot atoms for terminal sp³ carbon atoms and 1.2 times for all other carbon atoms. Crystallographic data for the structures reported in this paper have been deposited with the Cambridge Crystallographic Data Centre.^[5] CCDC ?????? contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/structures. This report and the CIF file were generated using FinalCif.^[6]

Table 5: Crystal data and structure refinement for cu_4g

CCDC number	2588894
Empirical formula	C ₂₁ H ₁₉ BrO ₃
Formula weight	399.27
Temperature [K]	100(2)
Crystal system	triclinic
Space group (number)	$P\bar{1}$ (2)
<i>a</i> [Å]	11.2175(5)
<i>b</i> [Å]	11.6447(5)
<i>c</i> [Å]	15.3731(6)
α [°]	78.532(2)
β [°]	81.760(2)
γ [°]	65.099(2)
Volume [Å ³]	1781.04(13)
<i>Z</i>	4
ρ_{calc} [gcm ⁻³]	1.489
μ [mm ⁻¹]	3.282
<i>F</i> (000)	816
Crystal size [mm ³]	0.16×0.18×0.22
Crystal colour	colourless
Crystal shape	cube
Radiation	CuK α (λ =1.54178 Å)
2 θ range [°]	5.88 to 158.31 (0.78 Å)
Index ranges	$-13 \leq h \leq 14$ $-14 \leq k \leq 14$ $0 \leq l \leq 19$
Reflections collected	261745
Independent reflections	7571 $R_{\text{int}} = 0.0995$ $R_{\text{sigma}} = 0.0287$
Completeness to $\theta = 67.679^\circ$	99.7
Data / Restraints / Parameters	7571 / 0 / 462
Goodness-of-fit on F^2	1.029
Final <i>R</i> indexes [$I \geq 2\sigma(I)$]	$R_1 = 0.0426$ $wR_2 = 0.1115$
Final <i>R</i> indexes [all data]	$R_1 = 0.0503$ $wR_2 = 0.1205$
Largest peak/hole [eÅ ⁻³]	0.61/−0.68




Figure 2. ORTEP view of compound **4g** showing the atom-numbering scheme, Displacement ellipsoids are drawn at the 50% probability level and H atoms are shown as small spheres with arbitrary radii. The compound crystallizes in the triclinic space group $P\bar{1}$ containing two molecules in the asymmetric unit, of which only molecule labelled with A suffix is shown. The Bromine atom also showed thermal disorder over two positions with occupancies 95% and 5%.

Refinement details for cu_4g

Refined as a 2-component twin.

Table 6. Bond lengths and angles for cu_4g

Atom–Atom	Length [Å]
Br1A–C10A	1.900(3)
Br2A–C10A	1.876(12)
O1A–C19A	1.202(4)
O2A–C20A	1.201(4)
O3A–C20A	1.341(3)
O3A–C21A	1.448(3)
C1A–C19A	1.510(4)
C1A–C20A	1.517(4)
C1A–C3A	1.559(4)
C1A–C2A	1.561(4)
C2A–C4A	1.565(4)
C3A–C4A	1.558(4)
C4A–C5A	1.509(4)
C4A–C13A	1.519(4)
C5A–C6A	1.329(4)
C6A–C7A	1.473(4)

C7A–C8A	1.391(4)
C7A–C12A	1.406(4)
C8A–C9A	1.386(4)
C9A–C10A	1.388(4)
C10A–C11A	1.382(4)
C11A–C12A	1.392(4)
C13A–C18A	1.392(4)
C13A–C14A	1.400(4)
C14A–C15A	1.391(4)
C15A–C16A	1.390(4)
C16A–C17A	1.388(4)
C17A–C18A	1.393(4)
Br1B–C10B	1.901(3)
Br2B–C10B	1.928(9)
O1B–C19B	1.193(4)
O2B–C20B	1.204(4)
O3B–C20B	1.338(4)
O3B–C21B	1.448(3)
C1B–C20B	1.507(4)
C1B–C19B	1.511(4)
C1B–C3B	1.559(4)
C1B–C2B	1.559(4)
C2B–C4B	1.573(4)
C3B–C4B	1.567(4)
C4B–C13B	1.512(4)
C4B–C5B	1.521(4)
C5B–C6B	1.320(4)
C6B–C7B	1.481(4)
C7B–C8B	1.392(4)
C7B–C12B	1.393(4)
C8B–C9B	1.398(4)
C9B–C10B	1.379(4)
C10B–C11B	1.383(4)
C11B–C12B	1.382(4)
C13B–C18B	1.394(4)
C13B–C14B	1.394(4)
C14B–C15B	1.391(4)
C15B–C16B	1.390(4)
C16B–C17B	1.391(4)
C17B–C18B	1.392(4)
Atom–Atom–Atom	Angle [°]
C20A–O3A–C21A	115.8(2)
C19A–C1A–C20A	112.4(2)
C19A–C1A–C3A	109.5(2)
C20A–C1A–C3A	116.5(2)
C19A–C1A–C2A	113.8(2)
C20A–C1A–C2A	113.6(2)

C3A–C1A–C2A	89.07(19)
C1A–C2A–C4A	89.89(19)
C4A–C3A–C1A	90.2(2)
C5A–C4A–C13A	105.3(2)
C5A–C4A–C3A	118.1(2)
C13A–C4A–C3A	115.6(2)
C5A–C4A–C2A	112.9(2)
C13A–C4A–C2A	116.0(2)
C3A–C4A–C2A	88.93(19)
C6A–C5A–C4A	127.2(3)
C5A–C6A–C7A	126.4(3)
C8A–C7A–C12A	117.7(3)
C8A–C7A–C6A	118.8(2)
C12A–C7A–C6A	123.3(3)
C9A–C8A–C7A	121.7(3)
C8A–C9A–C10A	119.0(3)
C11A–C10A–C9A	121.4(2)
C11A–C10A–Br2A	122.9(4)
C9A–C10A–Br2A	114.4(4)
C11A–C10A–Br1A	119.4(2)
C9A–C10A–Br1A	119.2(2)
C10A–C11A–C12A	118.7(3)
C11A–C12A–C7A	121.5(3)
C18A–C13A–C14A	119.3(3)
C18A–C13A–C4A	120.1(2)
C14A–C13A–C4A	120.5(2)
C15A–C14A–C13A	120.4(3)
C16A–C15A–C14A	120.2(3)
C17A–C16A–C15A	119.4(3)
C16A–C17A–C18A	120.8(3)
C13A–C18A–C17A	119.9(3)
O1A–C19A–C1A	124.2(3)
O2A–C20A–O3A	124.5(3)
O2A–C20A–C1A	125.6(3)
O3A–C20A–C1A	109.8(2)
C20B–O3B–C21B	116.3(2)
C20B–C1B–C19B	111.6(2)
C20B–C1B–C3B	116.2(2)
C19B–C1B–C3B	110.7(2)
C20B–C1B–C2B	114.7(2)
C19B–C1B–C2B	113.0(2)
C3B–C1B–C2B	89.05(19)
C1B–C2B–C4B	90.3(2)
C1B–C3B–C4B	90.5(2)
C13B–C4B–C5B	111.3(2)
C13B–C4B–C3B	115.8(2)
C5B–C4B–C3B	111.1(2)
C13B–C4B–C2B	115.5(2)

C5B–C4B–C2B	113.0(2)
C3B–C4B–C2B	88.29(19)
C6B–C5B–C4B	124.6(3)
C5B–C6B–C7B	127.3(3)
C8B–C7B–C12B	118.3(3)
C8B–C7B–C6B	123.3(3)
C12B–C7B–C6B	118.3(3)
C7B–C8B–C9B	121.0(3)
C10B–C9B–C8B	118.7(3)
C9B–C10B–C11B	121.6(3)
C9B–C10B–Br1B	119.5(2)
C11B–C10B–Br1B	118.9(2)
C9B–C10B–Br2B	126.3(3)
C11B–C10B–Br2B	110.4(3)
C12B–C11B–C10B	118.8(3)
C11B–C12B–C7B	121.6(3)
C18B–C13B–C14B	119.0(3)
C18B–C13B–C4B	119.6(2)
C14B–C13B–C4B	121.4(2)
C15B–C14B–C13B	120.5(3)
C16B–C15B–C14B	120.0(3)
C15B–C16B–C17B	119.9(3)
C16B–C17B–C18B	119.9(3)
C17B–C18B–C13B	120.7(3)
O1B–C19B–C1B	125.0(3)
O2B–C20B–O3B	124.7(3)
O2B–C20B–C1B	125.7(3)
O3B–C20B–C1B	109.6(2)


Bonds and angles to hydrogen atoms were omitted.

Bibliography


- [1] Bruker, *SAINT*, (Bruker,, Bruker AXS Inc., Madison, Wisconsin, USA.
- [2] L. Krause, R. Herbst-Irmer, G. M. Sheldrick, D. Stalke, *J. Appl. Cryst.* **2015**, 48, 3–10, doi:10.1107/S1600576714022985.
- [3] G. M. Sheldrick, *Acta Cryst.* **2015**, A71, 3–8, doi:10.1107/S2053273314026370.
- [4] G. M. Sheldrick, *Acta Cryst.* **2015**, C71, 3–8, doi:10.1107/S2053229614024218.
- [5] C. R. Groom, I. J. Bruno, M. P. Lightfoot, S. C. Ward, *Acta Cryst.* **2016**, B72, 171–179, doi:10.1107/S2052520616003954.
- [6] D. Kratzert, *FinalCif*, (Bruker Edition), <https://dkratzert.de/finalcif.html>.

9. Copies of ^1H , ^{13}C NMR spectra:

^1H NMR (400 MHz, CDCl_3) spectrum of Methyl (1s,3s)-1-formyl-3-phenyl-3-((E)-styryl)cyclobutane-1-carboxylate (4a)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl (1s,3s)-1-formyl-3-phenyl-3-((E)-styryl)cyclobutane-1-carboxylate (4a)




HRMS Spectrum of Methyl (1s,3s)-1-formyl-3-phenyl-3-((E)-styryl)cyclobutane-1-carboxylate (4a)


SJ-11 #503 RT: 3.86 AV: 1 NL: 3.24E5
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Methyl (1s,3s)-3-((E)-4-fluorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4b)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl (1s,3s)-3-((E)-4-fluorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4b)




HRMS Spectrum of Methyl (1s,3s)-3-((E)-4-fluorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4b)


SJ-2_20260812235019 #348 RT: 1.90 AV: 1 NL: 9.59E7
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Methyl (1s,3s)-3-((E)-3,4-difluorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4c)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl (1*s*,3*s*)-3-((*E*)-3,4-difluorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (**4c**)




HRMS Spectrum of Methyl (1s,3s)-3-((E)-3,4-difluorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4c)


SJ-8 #356 RT: 1.95 AV: 1 NL: 1.18E7
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Methyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4d)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4d)




HRMS Spectrum of Methyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4d)


SJ-22 #532 RT: 4.01 AV: 1 NL: 5.33E5
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Methyl (1s,3s)-3-((E)-3-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4e)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl (1s,3s)-3-((E)-3-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4e)




HRMS Spectrum of Methyl (1s,3s)-3-((E)-3-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4e)


SJ-3 20260812235618 #398 RT: 2.17 AV: 1 NL: 6.36E7
T: FTMS + p ESI Full ms [100.0000-1500.0000]



^1H NMR (400 MHz, CDCl_3) spectrum of Methyl (1s,3s)-3-((E)-2-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4f)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl (1*s*,3*s*)-3-((*E*)-2-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (**4f**)




HRMS spectrum of Methyl (1s,3s)-3-((E)-2-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4f)


SJR-15 #441 RT: 3.51 AV: 1 NL: 4.27E5
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Methyl (1s,3s)-3-((E)-4-bromostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4g)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl (1s,3s)-3-((E)-4-bromostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4g)




HRMS spectrum of Methyl (1s,3s)-3-((E)-4-bromostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4g)


SJR-16 #379 RT: 2.07 AV: 1 NL: 2.12E7
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Methyl (1s,3s)-1-formyl-3-((E)-4-methylstyryl)-3-phenylcyclobutane-1-carboxylate (4h)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl (1s,3s)-1-formyl-3-((E)-4-methylstyryl)-3-phenylcyclobutane-1-carboxylate (4h)




HRMS spectrum of Methyl (1s,3s)-1-formyl-3-((E)-4-methylstyryl)-3-phenylcyclobutane-1-carboxylate (4h)


SJ-5 #379 RT: 2.06 AV: 1 NL: 1.85E8
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Methyl (1s,3s)-1-formyl-3-((E)-3-methylstyryl)-3-phenylcyclobutane-1-carboxylate (4i)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl (1s,3s)-1-formyl-3-((E)-3-methylstyryl)-3-phenylcyclobutane-1-carboxylate (4i)




HRMS spectrum of Methyl (1s,3s)-1-formyl-3-((E)-3-methylstyryl)-3-phenylcyclobutane-1-carboxylate (4i)


SJ-6 #384 RT: 2.08 AV: 1 NL: 3.60E8
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Methyl 4-((E)-2-((1s,3s)-3-formyl-3-(methoxycarbonyl)-1-phenylcyclobutyl)vinyl)benzoate (4j)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl 4-((E)-2-((1s,3s)-3-formyl-3-(methoxycarbonyl)-1-phenylcyclobutyl)vinyl)benzoate (4i)




HRMS spectrum of Methyl 4-((E)-2-((1s,3s)-3-formyl-3-(methoxycarbonyl)-1-phenylcyclobutyl)vinyl)benzoate (4i)


SJ-20 #357 RT: 1.91 AV: 1 NL: 1.82E8
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Methyl (E)-3-(4-cyanostyryl)-1-phenyl-2-oxabicyclo[2.1.1]hexane-4-carboxylate (3k)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl (E)-3-(4-cyanostyryl)-1-phenyl-2-oxabicyclo[2.1.1]hexane-4-carboxylate (**3k**)




HRMS spectrum of Methyl (E)-3-(4-cyanostyryl)-1-phenyl-2-oxabicyclo[2.1.1]hexane-4-carboxylate (3k)


SJ-18 #348 RT: 1.87 AV: 1 NL: 2.80E7
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Methyl (E)-3-(4-nitrostyryl)-1-phenyl-2-oxabicyclo[2.1.1]hexane-4-carboxylate (3I)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl (E)-3-(4-nitrostyryl)-1-phenyl-2-oxabicyclo[2.1.1]hexane-4-carboxylate (3l)




HRMS spectrum of Methyl (E)-3-(4-nitrostyryl)-1-phenyl-2-oxabicyclo[2.1.1]hexane-4-carboxylate (3I)


SJ-19 #363 RT: 1.96 AV: 1 NL: 1.74E7
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Methyl (1s,3s)-1-formyl-3-((E)-2-methoxystyryl)-3-phenylcyclobutane-1-carboxylate (4m)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl (1s,3s)-1-formyl-3-((E)-2-methoxystyryl)-3-phenylcyclobutane-1-carboxylate (4m)




HRMS spectrum of Methyl (1s,3s)-1-formyl-3-((E)-2-methoxystyryl)-3-phenylcyclobutane-1-carboxylate (4m)


SJ-7 #349 RT: 1.90 AV: 1 NL: 4.53E8
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Methyl (1s,3s)-1-formyl-3-((E)-2-(naphthalen-2-yl)vinyl)-3-phenylcyclobutane-1-carboxylate (4n)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl (1s,3s)-1-formyl-3-((E)-2-(naphthalen-2-yl)vinyl)-3-phenylcyclobutane-1-carboxylate (4n)




HRMS spectrum of Methyl (1s,3s)-1-formyl-3-((E)-2-(naphthalen-2-yl)vinyl)-3-phenylcyclobutane-1-carboxylate (4n)


SJR-2 #416 RT: 2.19 AV: 1 NL: 3.21E7
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Methyl (1s,3s)-1-formyl-3-phenyl-3-((E)-2-(thiophen-2-yl)vinyl)cyclobutane-1-carboxylate (4o)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl (1s,3s)-1-formyl-3-phenyl-3-((E)-2-(thiophen-2-yl)vinyl)cyclobutane-1-carboxylate (4o)




HRMS spectrum of Methyl (1s,3s)-1-formyl-3-phenyl-3-((E)-2-(thiophen-2-yl)vinyl)cyclobutane-1-carboxylate (4o)


SJR-10 #359 RT: 1.93 AV: 1 NL: 4.64E7
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Methyl (E)-3-(2-cyclohexylvinyl)-1-phenyl-2-oxabicyclo[2.1.1]hexane-4-carboxylate (3p)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl (E)-3-(2-cyclohexylvinyl)-1-phenyl-2-oxabicyclo[2.1.1]hexane-4-carboxylate (3p)




HRMS spectrum of Methyl (E)-3-(2-cyclohexylvinyl)-1-phenyl-2-oxabicyclo[2.1.1]hexane-4-carboxylate (3p)


SJ-17 #553 RT: 2.95 AV: 1 NL: 1.40E8
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Tert-butyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4q)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Tert-butyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (**4q**)




HRMS spectrum of Tert-butyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4q)


SJR-17 #864 RT: 4.66 AV: 1 NL: 3.20E5
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Benzyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4r)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Benzyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4r)




HRMS spectrum of Benzyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4r)


SJ-9 #449 RT: 2.44 AV: 1 NL: 1.96E7
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-N-methoxy-N-methyl-3-phenylcyclobutane-1-carboxamide (4s)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-N-methoxy-N-methyl-3-phenylcyclobutane-1-carboxamide (4s)




HRMS spectrum of (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-N-methoxy-N-methyl-3-phenylcyclobutane-1-carboxamide (4s)


SJ-11_20260813004434 #379 RT: 2.05 AV: 1 NL: 4.41E8
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of (E)-(3-(4-chlorostyryl)-1-phenyl-2-oxabicyclo[2.1.1]hexan-4-yl)(naphthalen-2-yl)methanone (3t)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of (E)-(3-(4-chlorostyryl)-1-phenyl-2-oxabicyclo[2.1.1]hexan-4-yl)(naphthalen-2-yl)methanone (3t)




HRMS spectrum of (E)-(3-(4-chlorostyryl)-1-phenyl-2-oxabicyclo[2.1.1]hexan-4-yl)(naphthalen-2-yl)methanone (3t)


SJ-21 #432 RT: 2.32 AV: 1 NL: 8.79E6
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Methyl (1s,3s)-3-((E)-4-chlorostyryl)-3-(4-fluorophenyl)-1-formylcyclobutane-1-carboxylate (4u)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl (1s,3s)-3-((E)-4-chlorostyryl)-3-(4-fluorophenyl)-1-formylcyclobutane-1-carboxylate (4u)




HRMS spectrum of Methyl (1s,3s)-3-((E)-4-chlorostyryl)-3-(4-fluorophenyl)-1-formylcyclobutane-1-carboxylate (4u)


SJR-19 #384-387 RT: 2.09-2.11 AV: 4 NL: 6.60E6
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Methyl (1s,3s)-3-(4-chlorophenyl)-3-((E)-4-chlorostyryl)-1-formylcyclobutane-1-carboxylate (4v)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl (1s,3s)-3-(4-chlorophenyl)-3-((E)-4-chlorostyryl)-1-formylcyclobutane-1-carboxylate (4v)




HRMS spectrum of Methyl (1s,3s)-3-(4-chlorophenyl)-3-((E)-4-chlorostyryl)-1-formylcyclobutane-1-carboxylate (4v)


SJR-20 #419 RT: 2.28 AV: 1 NL: 5.60E6
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Methyl (1s,3s)-3-(4-bromophenyl)-3-((E)-4-chlorostyryl)-1-formylcyclobutane-1-carboxylate (4w)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl (1s,3s)-3-(4-bromophenyl)-3-((E)-4-chlorostyryl)-1-formylcyclobutane-1-carboxylate (4w)




HRMS spectrum of Methyl (1s,3s)-3-(4-bromophenyl)-3-((E)-4-chlorostyryl)-1-formylcyclobutane-1-carboxylate (4w)


SJR-18 #437 RT: 2.37 AV: 1 NL: 4.00E6
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Methyl (1s,3s)-3-(4-(tert-butyl)phenyl)-3-((E)-4-chlorostyryl)-1-formylcyclobutane-1-carboxylate (4x)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl (1s,3s)-3-(4-(tert-butyl)phenyl)-3-((E)-4-chlorostyryl)-1-formylcyclobutane-1-carboxylate (4x)




HRMS spectrum of Methyl (1s,3s)-3-(4-(tert-butyl)phenyl)-3-((E)-4-chlorostyryl)-1-formylcyclobutane-1-carboxylate (4x)


SJR-12 #313 RT: 2.52 AV: 1 NL: 8.90E4
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Methyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-(4-(trifluoromethyl)phenyl)cyclobutane-1-carboxylate (4y)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-(4-(trifluoromethyl)phenyl)cyclobutane-1-carboxylate (4y)




HRMS spectrum of Methyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-(4-(trifluoromethyl)phenyl)cyclobutane-1-carboxylate (4y)


SJR-7 #404 RT: 2.14 AV: 1 NL: 2.90E6
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Methyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-(3-methoxyphenyl)cyclobutane-1-carboxylate (4z)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-(3-methoxyphenyl)cyclobutane-1-carboxylate (4z)




HRMS spectrum of Methyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-(3-methoxyphenyl)cyclobutane-1-carboxylate (4z)


SJ-12 #369 RT: 2.02 AV: 1 NL: 9.09E7
T: FTMS + p ESI Full ms [100.0000-1500.0000]



^1H NMR (400 MHz, CDCl_3) spectrum of Methyl(1s,3s)-3-([1,1'-biphenyl]-4-yl)-3-((E)-4-chlorostyryl)-1-formylcyclobutane-1-carboxylate (4aa)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl(1s,3s)-3-([1,1'-biphenyl]-4-yl)-3-((E)-4-chlorostyryl)-1-formylcyclobutane-1-carboxylate (4aa)




HRMS spectrum of Methyl(1s,3s)-3-([1,1'-biphenyl]-4-yl)-3-((E)-4-chlorostyryl)-1-formylcyclobutane-1-carboxylate (4aa)


SJ-15 #468 RT: 2.52 AV: 1 NL: 2.67E7
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Methyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-(naphthalen-1-yl)cyclobutane-1-carboxylate (4ab)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-(naphthalen-1-yl)cyclobutane-1-carboxylate (4ab)




HRMS spectrum of Methyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-(naphthalen-1-yl)cyclobutane-1-carboxylate (4ab)


SJ-13 #415 RT: 2.26 AV: 1 NL: 1.66E8
T: FTMS + p ESI Full ms [100.0000-1500.0000]




¹H NMR (400 MHz, CDCl₃) spectrum of (4R,8R,9R,10S,13S,17S)-10,13-dimethyl-17-((S)-6-methylheptan-2-yl)-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1H-cyclopenta[a]phenanthren-4-yl (1s,3S)-3-((E)-4-chloro-styryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4ac)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of (4R,8R,9R,10S,13S,17S)-10,13-dimethyl-17-((S)-6-methylheptan-2-yl)-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1H-cyclopenta[a]phenanthren-4-yl (1s,3S)-3-((E)-4-chloro-styryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4ac)



¹H NMR (400 MHz, CDCl₃) spectrum of (1S,2R,4R)-2-isopropyl-4-methylcyclohexyl (1s,3R)-3-((E)-4-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4ad)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of (1S,2R,4R)-2-isopropyl-4-methylcyclohexyl (1s,3R)-3-((E)-4-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4ad)




HRMS spectrum of (1S,2R,4R)-2-isopropyl-4-methylcyclohexyl (1s,3R)-3-((E)-4-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4ad)


SJR-9 #711 RT: 3.76 AV: 1 NL: 7.97E6
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of 5-(2,4-dimethylphenoxy)-2,2-dimethylpentyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4ae)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of 5-(2,4-dimethoxyphenyl)-2,2-dimethylpentyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-phenylcyclobutane-1-carboxylate (4ae)




HRMS spectrum of 5-(2,4-dimethylphenoxy)-2,2-dimethylpentyl (1s,3s)-3-((E)-4-chlorostyryl)-1-formyl-3-phen-ylcyclobutane-1-carboxylate (4ae)


SJR-11 #461 RT: 3.70 AV: 1 NL: 9.31E4
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of (E)-(3-(4-chlorostyryl)-3-phenylcyclobutane-1,1-diyl)dimethanol (5)




$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of (E)-(3-(4-chlorostyryl)-3-phenylcyclobutane-1,1-diyl)dimethanol (5)




HRMS spectrum of (E)-(3-(4-chlorostyryl)-3-phenylcyclobutane-1,1-diyl)dimethanol (5)


SJ-22 #499 RT: 2.70 AV: 1 NL: 1.73E8
T: FTMS + p ESI Full ms [100.0000-1500.0000]



¹H NMR (400 MHz, CDCl₃) spectrum of Methyl (1*r*,3*s*)-3-(4-(*tert*-butyl)phenyl)-3-(4-chlorophenethyl)-1-formylcyclobutane-1-carboxylate (6)



$^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) spectrum of Methyl (1*r*,3*s*)-3-(4-(*tert*-butyl)phenyl)-3-(4-chlorophenethyl)-1-formylcyclobutane-1-carboxylate (**6**)



HRMS spectrum of Methyl (1r,3s)-3-(4-(tert-butyl)phenyl)-3-(4-chlorophenethyl)-1-formylcyclobutane-1- carboxylate (6)

SJ-24 #373 RT: 2.04 AV: 1 NL: 5.39E8
T: FTMS + p ESI Full ms [100.0000-1500.0000]

