

Cooperative NHC/LA Catalysis for Stereoselective Strain-Release Difunctionalization of Bicyclo[1.1.0]butanes to 1,1,3,3-Tetrasubstituted Cyclobutanes

Santosh Mhaske

`sb.mhaske.ncl@csir.res.in`

CSIR-National Chemical Laboratory <https://orcid.org/0000-0002-5859-0838>

Sayali Jagtap

CSIR-National Chemical Laboratory

Gaurang Bhatt

CSIR-National Chemical Laboratory

Rajesh Gonnade

CSIR-National Chemical Laboratory

Article

Keywords:

Posted Date: September 22nd, 2026

DOI: <https://doi.org/10.21203/rs.3.rs-11093434/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Additional Declarations: Yes there is potential Competing Interest. A provisional patent application (Application number 202611102297, dated 24-08-2026) has been filed based on this chemistry.

Cooperative NHC/Lewis Acid Catalysis for Stereoselective Strain-Release Geminal Difunctionalization of Bicyclo[1.1.0]butanes to 1,1,3,3-Tetrasubstituted Cyclobutanes

Sayali G. Jagtap,^{a,c} Gaurang J. Bhatt,^a Rajesh G. Gonnade^{bc} and Santosh B. Mhaske^{*ac}


^aOrganic Chemistry Division, CSIR-National Chemical Laboratory (CSIR-NCL), Dr. Homi Bhabha Road, Pune-411008 (India).

^bPhysical and Materials Chemistry Division, CSIR-National Chemical Laboratory (CSIR-NCL), Dr. Homi Bhabha Road, Pune-411008 (India).

^cAcademy of Scientific and Innovative Research (AcSIR) Ghaziabad-201002 (India).

*E-mail: sb.mhaske.ncl@csir.res.in

The stereoselective synthesis of densely substituted cyclobutanes has emerged as an important frontier in contemporary molecular synthesis, owing to their value as three-dimensional building blocks and bioisosteric motifs. Here we report a cooperative N-heterocyclic carbene (NHC)/Lewis acid catalytic strategy for strain-release difunctionalization of bicyclo[1.1.0]butanes (BCBs) with α,β -unsaturated aldehydes, affording 1,1,3,3-tetrasubstituted cyclobutanes in up to 94% yield with excellent syn-dia stereoselectivity across a broad substrate scope. Most reactions reach completion within 15 min at room temperature. Mechanistic studies, including isolation of a 2-oxabicyclo[2.1.1]hexane intermediate and its subsequent conversion to the cyclobutane product under the catalytic conditions, establish its involvement in the reaction pathway. The data support Lewis acid-assisted C–O σ -bond cleavage followed by NHC-mediated ring opening and rearrangement. This transformation reveals an unconventional mode of NHC reactivity in strain-release chemistry and demonstrates how cooperative NHC/Lewis acid catalysis can redirect the reactivity of strained oxygenated bicyclic intermediates.



The increasing demand for structurally complex, sp^3 -rich molecular architectures has reshaped modern medicinal chemistry, as saturated three-dimensional (3D) frameworks can offer favorable physicochemical and pharmacokinetic properties compared with their planar counterparts.^[1-4] In the pursuit of ‘escaping from flatland’,^[5-9] cyclobutanes have emerged as valuable structural motifs owing to their rigid, conformationally constrained frameworks, which enable precise spatial orientation of substituents and impart distinct three-dimensional character to molecular architectures. Consequently, cyclobutane-containing scaffolds have found widespread applications in medicinal chemistry,^[10-15] and several pharmaceuticals incorporating 1,3-disubstituted cyclobutane motifs have advanced to clinical evaluation. These applications have stimulated the development of efficient and versatile strategies for the synthesis of increasingly complex and densely functionalized cyclobutanes (Fig. 1A).




Fig. 1 The difunctionalization of BCBs to access 1,1,3-trisubstituted and 1,1,3,3-tetrasubstituted cyclobutanes. A) Representative drug molecules containing polyfunctionalized cyclobutane skeleton. B) Strategies for the synthesis of highly functionalized cyclobutanes via strain release. C) Our protocol for the synthesis of 1,1,3,3-tetrasubstituted cyclobutanes via NHC/Lewis acid cooperative catalysis

BCBs have recently emerged as versatile strain-release synthons for the rapid construction of functionalized cyclobutanes.^[16-18] Their exceptionally weak and highly strained central C–C σ -bond exhibits pronounced π -bond-like character, rendering it susceptible to activation by nucleophilic, electrophilic, and radical species.^[19-28] Cleavage of this ‘spring-loaded’ bond releases substantial strain energy, enabling the efficient conversion of readily

accessible BCBs into structurally diverse cyclobutane frameworks. This distinctive reactivity has led to the development of diverse catalytic strain-release functionalization and polyfunctionalization reactions with high levels of regio- and stereocontrol across a range of catalytic manifolds.^[29-32] Despite these advances, the catalytic synthesis of highly substituted cyclobutanes from BCBs remains challenging, particularly when precise control over the relative stereochemistry generated during strain-release ring opening is required.^[21,22,27,28]

Several recent studies have highlighted the potential of BCBs for stereoselective access to highly substituted cyclobutanes.^[17] Aggarwal and co-workers established an electrophilic radical-induced ring opening of boronate-substituted BCBs (Fig. 1B, path 1), in which radical addition generates an α -boryl radical that undergoes stereospecific 1,2-boron migration to furnish highly functionalized cyclobutanes with excellent diastereoselectivity.^[24,33,34] Subsequently, Glorius and co-workers disclosed a photoredox-enabled thio-carbofunctionalization of BCBs using sulfur-containing bifunctional reagents (Fig. 1B, path 2). In this process, strain-release ring opening followed by C–S bond cleavage of a cyclic radical cation intermediate provides *syn*-functionalized cyclobutanes with high stereocontrol.^[35] Although these approaches demonstrate the versatility of BCB strain-release chemistry, they are largely centered on monosubstituted BCBs, limiting their direct application to more substituted cyclobutane architectures. Ackermann and co-workers reported an interesting Ru(II)-catalyzed C–C/C–H functionalization of BCBs, integrating strain-release ring opening, halogen-atom transfer, and site-selective C–H functionalization to access functionalized tri- and tetrasubstituted cyclobutanes (Fig. 1B, path 3).^[36] More recently, Zhao and co-workers reported an N-heterocyclic carbene (NHC)-catalyzed acylfluoroalkylation of BCBs through an NHC-derived ketyl radical, providing access to 1,1,3,3-tetrasubstituted cyclobutanes with high diastereoselectivity (Fig. 1B, path 4).^[37] This transformation utilizes fluoroalkyl radical precursor, such as a Togni's reagent, to achieve tetrasubstitution, with most of the substrates providing excellent diastereoselectivity. Thus, a general catalytic strategy that directly converts readily accessible BCBs into stereodefined 1,1,3,3-tetrasubstituted cyclobutanes through a simple two-component coupling remains highly desirable. In particular, achieving geminal difunctionalization of BCBs while simultaneously controlling the stereochemical outcome of strain release remains a significant synthetic challenge.


In recent years, cooperative catalysis with NHC has emerged as a powerful strategy for the development of diverse synthetic transformations.^[38] Here, we report a cooperative NHC/Lewis acid catalytic strategy for the stereoselective strain-release geminal difunctionalization of BCBs, enabling rapid access to 1,1,3,3-tetrasubstituted cyclobutanes

(Fig. 1C). The two-component protocol directly couples readily accessible BCBs with α,β -unsaturated aldehydes to furnish densely functionalized cyclobutanes in very good yield and with excellent diastereoselectivities. This cooperative catalytic strategy provides a direct and stereoselective approach to highly substituted cyclobutanes while expanding the reactivity landscape of NHC-mediated strain-release chemistry.

Results and Discussion

We began our investigation using 4-chlorocinnamaldehyde (**1d**) and methyl 3-phenylbicyclo[1.1.0]butane-1-carboxylate (**2a**) as reaction partners for the optimization of the reaction conditions (Table 1, selected entries, see SI).

Table 1 Optimization of the reaction conditions^a



Entry	Variation from Standard Condition ^a	Yield of 3d (%) ^b	Yield of 4d (%) ^b
1.	None	n.d.	85%
2.	20 mol% Sc(OTf) ₃ instead of 30 mol%	20%	57%
3.	NHC B instead of NHC A	7%	71%
4.	NHC C instead of NHC A	45%	33%
5.	1.5 equiv. of 2a instead of 2 equiv.	15%	40%
6.	Cs ₂ CO ₃ instead of K ₂ CO ₃	n.d.	24%
7.	DBU instead of K ₂ CO ₃	n.d.	18%
8.	Toluene instead of THF	n.d.	49%
9.	CHCl ₃ instead of THF	10%	48%
10.	CH ₃ CN instead of THF	89%	n.d.
11.	Without NHC	78%	5%
12.	Without Sc(OTf) ₃	n.d.	n.d.

^aReaction conditions: **1d** (0.16 mmol), **2a** (2.0 equiv.), NHC A (10 mol%), Sc(OTf)₃ (30 mol%), K₂CO₃ (40 mol%), THF (0.2 mL), Temp. = 0 °C to rt, t = 15 min. ^bYield of isolated products, n.d. = Not detected.

Gratifyingly, the reaction proceeded smoothly under the optimized cooperative catalytic conditions, employing 10 mol% of NHC A, 30 mol% Sc(OTf)₃, and 40 mol% K₂CO₃ in THF. Conducting the reaction from 0 °C to room temperature enabled rapid conversion within just

15 min, furnishing the desired 1,1,3,3-tetrasubstituted cyclobutane **4d** in 85% isolated yield (Table 1, entry 1). A reduction in the Sc(OTf)₃ loading to 20 mol% resulted in a significant decrease in the product yield to 57%, along with formation of **3d** in 20% yield (Table 1, entry 2). Several NHC catalysts featuring triazolium-, imidazolium-, and thiazolium-derived carbene frameworks were initially screened for this transformation (see SI). Among them, NHC catalysts based on the triazolium scaffold proved to be the most effective, affording the desired product in good yields. Notably, decreasing the steric bulk of the NHC catalyst resulted in a noticeable reduction in reaction efficiency and, consequently, lower expected product yields (Table 1, entries 3 and 4). Screening of various Lewis acids identified Sc(OTf)₃ as uniquely effective for this transformation (see the Supporting Information). When the amount of BCB was reduced from 2.0 to 1.5 equivalents, the yield of **4d** decreased to 40%, with **3d** formed in 15% yield (Table 1, entry 5). The nature of the base was found to have a pronounced impact on the reaction efficiency (see the Supporting Information). Substitution of K₂CO₃ with either Cs₂CO₃ or DBU led to a dramatic erosion in the yield of the desired product (Table 1, entries 6 and 7). Evaluation of various solvents identified THF as the optimal medium for this transformation (Table 1, entries 8 and 9). The use of ACN as the solvent interestingly resulted in the formation of **3d** exclusively in 89% yield, with no detectable formation of the desired product **4d** (Table 1, entry 10). When the reaction was performed in the absence of the NHC catalyst, **3d** was formed in 78% yield, whereas the desired product **4d** was obtained in only 5% yield (Table 1, entry 11). In contrast, in the absence of Sc(OTf)₃, neither **3d** nor **4d** was detected (Table 1, entry 12).

With the optimized reaction conditions in hand, we next investigated the generality of this bifunctionalization protocol using a diverse range of α,β -unsaturated aldehydes and BCBs (Fig. 2). We initially examined the substrate scope with respect to the α,β -unsaturated aldehyde component. The cooperative NHC-Lewis acid catalytic system exhibited broad applicability, efficiently converting a variety of α,β -unsaturated aldehydes into the corresponding 1,1,3,3-tetrasubstituted cyclobutane derivatives in good to excellent yields and with consistently high diastereoselectivities (**4a-4j**, **4m-4o**). Cinnamaldehyde worked well to provide the desired product **4a** in excellent yield and stereoselectivity. Substrates bearing halo substituents at the para, meta and ortho positions, as well as dihalo substituted derivative underwent smooth transformation to furnish the corresponding products **4b-4g** in moderate to very good yields (20-85%). The molecular structure of intermediate **3g**, corresponding to the proposed intermediate **3**, was unambiguously confirmed by single-crystal X-ray diffraction analysis (see SI). The relative stereochemistry of compound **4g** was established by single-crystal X-ray

diffraction analysis, confirming the cis arrangement of the styryl and aldehyde groups on the cyclobutane framework (see SI).




Fig. 2 Substrate scope of α,β -unsaturated aldehyde and BCBs. Reaction condition: **1** (0.16 mmol), **2** (2 equiv.), NHC A (10 mol%), $\text{Sc}(\text{OTf})_3$ (30 mol%), K_2CO_3 (40 mol%), THF (0.2 mL), T = 0 °C-rt, t = 15 min. Isolated product yields. Diastereomeric ratio (d.r.) values were determined by ^1H NMR. ^aReaction mixture heated to 50 °C for 10 h.

Para and meta methyl substituted enals were also readily accommodated, affording products **4h** and **4i** in 68% and 73% yields, respectively. Furthermore, the electron-withdrawing methyl ester substituted cinnamaldehyde was well tolerated, delivering the corresponding product **4j** in a good yield of 70%. However, the protocol didn't work well with cyano and nitro substituted cinnamaldehydes probably due to lesser nucleophilicity of the NHC-enolate. Only the corresponding intermediates **3k** and **3l** were observed in good yields. The influence of ortho-methoxy substitution was also examined, which furnished the corresponding product **4m** in 66% yield. In addition, 2-naphthyl-substituted and β -heteroaryl enals were compatible with the reaction conditions, providing the corresponding products **4n** and **4o** in moderate yields and with good diastereoselectivities. Aliphatic enal afforded only the corresponding intermediate **3p**, while the expected product **4p** was not observed.

Having established the scope of the α,β -unsaturated aldehyde component, we next investigated the effect of structural variation in the BCB substrates **2q-2ab**. We first explored variations in the ester functionality of the BCB substrate. Both *tert*-butyl and benzyl ester-substituted BCBs were readily accommodated, furnishing products **4q** and **4r** in good yields of 74% and 79%, respectively. The Weinreb amide-substituted BCB also participated in the reaction, affording the desired product **4s** in 49% yield, however it requires higher temperature and longer reaction time. In contrast, the keto-substituted BCB did not furnish the desired product **4t** and proceeded only to the corresponding intermediate **3t**. We subsequently examined the influence of substituents present on the aryl group of the BCB substrates. Para halo-substituted phenyl BCB derivatives underwent efficient transformation under the cooperative NHC-Lewis acid catalytic conditions, affording the corresponding cyclobutane products **4u-4w** in moderate to good yields (56-81%). Notably, the para *tert*-butyl-substituted phenyl BCB delivered the corresponding product **4x** in an excellent yield under the optimized reaction conditions. In contrast, the presence of a strongly electron-withdrawing CF₃ group at the para position resulted in the reduced yield of the corresponding product **4y**. The meta methoxy-substituted phenyl BCB was also well tolerated, providing the desired product **4z** in 82% yield. Furthermore, the biphenyl BCB exhibited excellent reactivity, furnishing the corresponding product **4aa** in 93% yield. The 1-naphthyl-substituted BCB **4ab** also afforded the desired product in a good yield of 74%. Encouraged by the excellent tolerance of diverse functional groups, we evaluated the applicability of this protocol to the substrates featuring structurally complex and biologically relevant molecules. Accordingly, BCB substrates derived from cholesterol, *L*-menthol, and the drug gemfibrozil were subjected to the optimized reaction conditions. The corresponding cyclobutane products **4ac-4ae** were obtained in 45-94% yield.

Among these substrates, the cholesterol-derived BCB displayed particularly high reactivity, delivering the corresponding 1,1,3,3-tetrasubstituted cyclobutane derivative **4ac** in an excellent yield of 94%. These results highlight the excellent functional group tolerance and potential utility of this methodology for the late-stage diversification of structurally complex and biologically relevant molecules.

To gain insight into the mechanism of this transformation, a series of control experiments were performed (Fig. 3A).




Fig. 3 A) Control Experiments: i) Reaction using saturated aldehyde ii) Reactivity of intermediate under standard reaction condition iv) Reactivity of intermediate under standard reaction conditions in the absence of NHC or Sc(OTf)₃ B) Plausible Mechanism

The importance of the α,β -unsaturated aldehyde functionality for the protocol was investigated by replacing enal **1a** with 3-phenylpropanal **1a'**. When **1a'** and BCB **2a** were subjected to the standard reaction conditions, **1a'** remained unreacted, and neither the corresponding intermediate **3a'** nor the desired cyclobutane product **4a'** was detected (Fig. 3A-i). In contrast, BCB **2a** was completely consumed, with formation of the corresponding cyclobutene derivative. These observations suggest that the α,β -unsaturated aldehyde functionality plays a crucial role in the reaction with the Sc(OTf)₃-activated BCB substrate. The corresponding 2-oxa-bicyclo[2.1.1]hexane **3a'** has previously been synthesized by the Huang group through a cobalt-promoted photocatalytic $[2\pi + 2\sigma]$ cycloaddition of BCBs with aldehydes under visible-light irradiation.^[39] To examine whether the intermediate **3d** could serve as a precursor to the final product. It was subjected to the standard reaction conditions. Complete consumption of **3d** was observed within 15 mins, affording the corresponding 1,1,3,3-tetrasubstituted cyclobutane **4d** in 93% isolated yield (Fig. 3A-ii). This outcome clearly supports **3d** as a competent intermediate in the formation of the final product. Furthermore, the observed conversion is consistent with the possibility that Sc(OTf)₃ coordinates with oxygen and facilitates C–O bond cleavage during the conversion of **3d** to the corresponding cyclobutane product.

Finally, two additional control experiments were performed to investigate the individual roles of Sc(OTf)₃ and the NHC catalyst in the conversion of intermediate **3d** into the product **4d**. When **3d** was subjected to the optimized reaction conditions in the absence of Sc(OTf)₃, no conversion to **4d** was observed (Fig. 3A-iii). Likewise, omission of the NHC catalyst also failed to promote the formation of **4d** (Fig. 3A-iii). Taken together, these results indicate that both Sc(OTf)₃ and the NHC catalyst are required for the conversion of intermediate **3d** into product **4d**. Overall, these observations are consistent with the involvement of cooperative NHC/Lewis acid catalysis in facilitating the transformation.

On the basis of the control experiments and relevant literature reports^[40–42], a plausible reaction pathway for this transformation is proposed (Fig. 3B). Initially, the BCB is activated through coordination with the Lewis acid Sc(OTf)₃, which facilitates cleavage of its highly strained central C–C bond^[42]. The resulting activated BCB intermediate then reacts with the enal, followed by intramolecular cyclization to furnish the bioisosteric^[43] 2-oxabicyclo[2.1.1]hexane intermediate **3d**. Meanwhile, the NHC catalyst is generated *in situ* through deprotonation of the triazolium salt in the presence of a base. The resulting NHC attacks the double bond of intermediate **3d**, generating intermediate **II**, which subsequently undergoes proton transfer to afford intermediate **III**, followed by the formation of intermediate

IV. Subsequently, coordination of the oxygen atom with $\text{Sc}(\text{OTf})_3$ and the presence of the NHC in intermediate **IV** facilitates the simultaneous cleavage of the C–C and C–O bonds, giving rise to intermediate **V**. The intermediate **V** upon proton transfer, affords intermediate **VI**. Finally, release of the NHC catalyst from intermediate **VI** furnishes the 1,1,3,3-tetrasubstituted cyclobutane product **4d**.

To demonstrate the scalability and practical applicability of the developed methodology, two independent scale-up experiments were conducted under the optimized reaction conditions using cinnamaldehyde **1d** as the α,β -unsaturated aldehyde component. The reaction of **1d** (3 mmol, 500 mg) with BCB **2x** furnished the corresponding bifunctionalized cyclobutane product **4x** in an excellent yield of 90%. The reaction was subsequently performed on a larger 6 mmol scale using BCB **2a**, affording the corresponding product **4d** in 71% yield (Fig. 4A). The stereoselectivity was not hampered even on bigger scale demonstrating the robustness of the protocol.




Fig. 4 A) Gram Scale synthesis. B) Product functionalization.

The synthetic versatility of product **4x** was further demonstrated through complementary downstream transformations (Fig. 4B). Catalytic hydrogenation of the styryl moiety in **4x** using H_2 and Pd/C smoothly afforded the corresponding saturated cyclobutane derivative **5** in 85% yield. Chemoselective reduction of the aldehyde functionality in **4d** with NaBH_4 furnished the corresponding primary alcohol **6** in an excellent yield of 96%. Collectively, these transformations demonstrate the synthetic versatility of the polyfunctionalized cyclobutane products and further highlight the utility of the present methodology as a versatile platform for accessing structurally diverse and synthetically valuable molecular scaffolds.

Conclusion

In summary, we have developed a cooperative NHC/Lewis acid catalytic strategy for the stereoselective strain-release difunctionalization of bicyclo[1.1.0]butanes with enals, providing rapid access to 1,1,3,3-tetrasubstituted cyclobutanes under mild conditions. The transformation accommodates electronically and structurally diverse enals and BCBs, delivering high efficiency, high regioselectivity and excellent syn-diastereoselectivity. The resulting cyclobutanes contain synthetically versatile geminal aldehyde and ester or amide functionalities at highly substituted stereogenic centres, providing opportunities for further molecular diversification. Beyond its synthetic utility, this transformation reveals an unconventional mode of NHC reactivity enabled by Lewis acid cooperation. The isolation and independent conversion of the 2-oxabicyclo[2.1.1]hexane intermediate under the catalytic conditions provide compelling evidence for its role as a mechanistic gateway to the substituted cyclobutane products. This reactivity highlights how Lewis acid cooperation can unlock C–O bond cleavage and redirect the fate of strained oxygenated intermediates through NHC-mediated rearrangement. More broadly, this work expands the reactivity landscape of NHC catalysis and establishes cooperative activation as a platform for accessing new NHC-mediated transformations.

Methods

General optimized procedure for the synthesis of polyfunctionalised cyclobutane derivatives 4: An oven-dried schlenk tube equipped with a magnetic stirring bar was charged with the NHC precatalyst-A (10 mol%) and K₂CO₃ (40 mol%). The reaction vessel was evacuated and backfilled with argon three times. Under an argon atmosphere, α,β -unsaturated aldehyde **1** (1 equiv.) and bicyclobutane derivative **2** (2 equiv.) were added followed by addition of THF. The reaction vessel was evacuated and backfilled with argon three times. The schlenk tube was placed in an ice bath and the reaction mixture was stirred for 5 minutes. After this, Sc(OTf)₃ (30 mol%) was added and the tube was tightly closed with a glass stopper. The resulting mixture was kept for stirring at rt for completion of the reaction. Upon completion of the reaction as observed by TLC, the reaction mixture was directly subjected to column chromatography using ethyl acetate:pet ether over 230-400 flash silica gel to afford the desired products **4** in good to excellent yields.

References

1. Shearer, J. et al. Rings in clinical trials and drugs: present and future. *J. Med. Chem.* **65**, 8699–8712 (2022).
2. Taylor, R. D., MacCoss, M. & Lawson, A. D. G. Rings in drugs. *J. Med. Chem.* **57**, 5845–5859 (2014).
3. Campos, K. R. et al. The importance of synthetic chemistry in the pharmaceutical industry. *Science* **363**, eaat0805 (2019).
4. Gomez-Angel, A. R. et al. Modular synthetic platform for the elaboration of fragments in three dimensions for fragment-based drug discovery. *J. Am. Chem. Soc.* **147**, 29292–29303 (2025).
5. Lovering, F., Bikker, J. & Humblet, C. Escape from flatland: increasing saturation as an approach to improving clinical success. *J. Med. Chem.* **52**, 6752–6756 (2009).
6. Mykhailiuk, P. K. Saturated bioisosteres of benzene: where to go next? *Org. Biomol. Chem.* **17**, 2839–2849 (2019).
7. Golfmann, M. & Walker, J. C. L. Bicyclobutanes as unusual building blocks for complexity generation in organic synthesis. *Commun. Chem.* **6**, 9 (2023).
8. Kelly, C. B. et al. Bicyclobutanes: from curiosities to versatile reagents and covalent warheads. *Chem. Sci.* **13**, 11721–11737 (2022).
9. Tsien, J. et al. Three-dimensional saturated C(sp³)-rich bioisosteres for benzene. *Nat. Rev. Chem.* **8**, 605–627 (2024).
10. Dembitsky, V. M. Bioactive cyclobutane-containing alkaloids. *J. Nat. Med.* **62**, 1–33 (2008).
11. Marson, C. M. New and unusual scaffolds in medicinal chemistry. *Chem. Soc. Rev.* **40**, 5514–5533 (2011).
12. Liu, Q. et al. Cyclobutane derivatives as novel nonpeptidic small-molecule agonists of glucagon-like peptide-1 receptor. *J. Med. Chem.* **55**, 250–267 (2012).
13. Bauer, M. R. et al. Put a ring on it: application of small aliphatic rings in medicinal chemistry. *RSC Med. Chem.* **12**, 448–471 (2021).

14. Van Der Kolk, M. R. et al. Cyclobutanes in small-molecule drug candidates. *ChemMedChem* **17**, e202200020 (2022).
15. Ertl, P., Altmann, E. & Wilcken, R. Ring systems in medicinal chemistry: a cheminformatics analysis of ring popularity in drug discovery over time. *Eur. J. Med. Chem.* **300**, 118178 (2025).
16. Zhan, X., He, H.-X., Peng, Q. & Feng, J.-J. Synthesis of cyclobutanes and cyclobutenes by strain-release-driven ring-opening of bicyclo[1.1.0]butanes. *Synthesis* **56**, 3829–3848 (2024).
17. Mukherjee, A. & Maity, S. Bicyclo[1.1.0]butane: a flexible template for strain-release functionalization in cyclobutane synthesis. *Synthesis* (2026).
18. Deswal, S., Das, R. C., & Biju, A. T. Recent advances in the strain-release driven reactions of bicyclobutanes: an emerging landscape in organic synthesis. *Chem. Soc. Rev.* **55**, 5572–5589 (2026).
19. Panish, R., Chintala, S. R., Boruta, D. T., Fang, Y., Taylor, M. T. & Fox, J. M. Enantioselective synthesis of cyclobutanes via sequential Rh-catalyzed bicyclobutanation/Cu-catalyzed homoconjugate addition. *J. Am. Chem. Soc.* **135**, 9283–9286 (2013).
20. Gianatassio, R. et al. Strain-release amination. *Science* **351**, 241–246 (2016).
21. Ernouf, G. et al. Photochemical strain-release-driven cyclobutylation of C(sp³)-centered radicals. *Angew. Chem. Int. Ed.* **59**, 2618–2622 (2020).
22. Ociepa, M., Wierzba, A. J., Turkowska, J. & Gryko, D. Polarity-reversal strategy for the functionalization of electrophilic strained molecules via light-driven cobalt catalysis. *J. Am. Chem. Soc.* **142**, 5355–5361 (2020).
23. Guo, L., Noble, A. & Aggarwal, V. K. α -Selective ring-opening reactions of bicyclo[1.1.0]butyl boronic ester with nucleophiles. *Angew. Chem. Int. Ed.* **60**, 212–216 (2021).
24. Bennett, S. H. et al. Difunctionalization of C–C σ -bonds enabled by the reaction of bicyclo[1.1.0]butyl boronate complexes with electrophiles: reaction development, scope, and stereochemical origins. *J. Am. Chem. Soc.* **142**, 16766–16775 (2020).

25. Lewis-Borrell, L. et al. Direct observation of reactive intermediates by time-resolved spectroscopy unravels the mechanism of a radical-induced 1,2-metalate rearrangement. *J. Am. Chem. Soc.* **143**, 17191–17199 (2021).
26. Xiao, Y. et al. Photochemical α -selective radical ring-opening reactions of 1,3-disubstituted acyl bicyclobutanes with alkyl halides: modular access to functionalized cyclobutenes. *Chem. Sci.* **14**, 13060–13066 (2023).
27. Pratt, C. J., Aycock, R. A., King, M. D. & Jui, N. T. Radical α -C–H cyclobutylation of aniline derivatives. *Synlett* **31**, 51–54 (2020).
28. Yu, X., Lübbesmeyer, M. & Studer, A. Oligosilanes as silyl radical precursors through oxidative Si–Si bond cleavage using redox catalysis. *Angew. Chem.* **133**, 685–689 (2021).
29. Panish, R. A., Chintala, S. R. & Fox, J. M. A mixed-ligand chiral rhodium(II) catalyst enables the enantioselective total synthesis of piperarborenine B. *Angew. Chem. Int. Ed.* **55**, 4983–4987 (2016).
30. Kerner, M. J. & Wipf, P. Semipinacol-type rearrangements of [3-(arylsulfonyl)bicyclo[1.1.0]butan-1-yl]alkanols. *Org. Lett.* **23**, 3615–3619 (2021).
31. Milligan, J. A., Busacca, C. A., Senanayake, C. H. & Wipf, P. Hydrophosphination of bicyclo[1.1.0]butane-1-carbonitriles. *Org. Lett.* **18**, 4300–4303 (2016).
32. Zhang, Z. & Gevorgyan, V. Palladium hydride-enabled hydroalkenylation of strained molecules. *J. Am. Chem. Soc.* **144**, 20875–20883 (2022).
33. Fawcett, A., Biberger, T. & Aggarwal, V. K. Carbopalladation of C–C σ -bonds enabled by strained boronate complexes. *Nat. Chem.* **11**, 117–122 (2018).
34. Silvi, M. & Aggarwal, V. K. Radical addition to strained σ -bonds enables the stereocontrolled synthesis of cyclobutyl boronic esters. *J. Am. Chem. Soc.* **141**, 9511–9515 (2019).
35. Wang, H. et al. syn-Selective difunctionalization of bicyclobutanes enabled by photoredox-mediated C–S σ -bond scission. *J. Am. Chem. Soc.* **145**, 23771–23780 (2023).

36. Chen, S. et al. Difunctionalization of bicyclo[1.1.0]butanes enabled by merging C–C cleavage and ruthenium-catalysed remote C–H activation. *Nat. Synth.* **4**, 655–663 (2025).
37. Xiao, C. et al. Stereoselective radical acylfluoroalkylation of bicyclobutanes via N-heterocyclic carbene catalysis. *Angew. Chem. Int. Ed.* **64**, e202416781 (2024).
38. Wang, M. H., & Scheidt, K. A. Cooperative Catalysis and Activation with N-Heterocyclic Carbenes. *Angew. Chem. Int. Ed.* **55**, 14912–14922 (2016).
39. Tang, S., et al. Photocatalytic synthesis of 2-oxabicyclo[2.1.1]hexanes: cobalt-enhanced efficiency. *Chem. Sci.* **16**, 11908–11917 (2025).
40. Zhang, M. et al. Lewis acid-catalyzed ring-opening reaction of bicyclobutanes for the highly selective synthesis of 1,1,3-trisubstituted cyclobutanes. *Org. Lett.* **27**, 9471–9476 (2025).
41. Wang, J. et al. Sc(OTf)₃-catalyzed diastereoselective hydroxyheteroarylation of C–C σ -bonds of bicyclo[1.1.0]butanes with azaheterocyclic N-oxides. *Org. Lett.* **27**, 3402–3408 (2025).
42. Roy, P., Deswal, S. & Biju, A. T. Lewis acid-assisted formal C=S insertion of thioindolinones to bicyclobutanes: diastereoselective access to trisubstituted cyclobutanethiols. *Org. Lett.* **27**, 10916–10922 (2025).
43. Denisenko, A. et al. 2-Oxabicyclo[2.1.1]hexanes as saturated bioisosteres of the ortho-substituted phenyl ring. *Nat. Chem.* **15**, 1155–1163 (2023).

Acknowledgment

S.B.M. acknowledges financial support from CSIR, India. S.G.J. thanks DST-Inspire, India and G.J.B. thanks CSIR, New Delhi.

Author contribution

S.B.M. conceived the project. S.G.J. and G.J.B. performed the synthetic experiments. R.G.G. carried out single crystal X-Ray diffraction analysis of compounds **3g** and **4g**. S.B.M. supervised the project.

Competing interest

Provisional patent (Application number 202611102297, dated 24-08-2026) has been filed based on this chemistry.

Additional information

Supplementary information The online version contains supplementary material available at...

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupportingInformation.pdf](#)