

Supplementary Information

Targeting Nicotinic Acetylcholine Receptor Subunits to Modulate


Brucella-Induced Inflammation in Murine Macrophages

Can Sun, Xiaoyang Liang, Ahmad Ali, Rakhshanda Rani, Huili Lu, Yuefeng Wu, Xingmei


Deng, Jia Guo, Hui Zhang, Yong Wei, Hongsu He, and Zhihua Sun

Original source Western-blot images corresponding to the cropped blot panels in Figures 1E, 2D, 4E, and 5A are provided below. Membrane edges and molecular-weight references are shown where available. The images are labelled to match the relevant main-figure panels.


Supplementary Figure S3. Original source blots corresponding to Figure 1E




NLRP3 (110 kDa)



Pro-caspase-1 (44 kDa)




β -actin (42 kDa)




Supplementary Figure S4. Original source blots corresponding to Figure 2D


$\beta 4$ nAChR (56 kDa)




$\beta 3$ nAChR (53 kDa)




$\alpha 7$ nAChR (57 kDa)




$\alpha 6$ nAChR (56 kDa)




$\alpha 5$ nAChR (53 kDa)




$\alpha 4$ nAChR (74 kDa)




Supplementary Figure S5. Original source blots corresponding to Figure 4E
NLRP3 (110 kDa)




α 4 nAChR/Chrna4 (74 kDa)




$\alpha 7$ nAChR/Chrna7 (56 kDa)




Pro-caspase-1 (44 kDa)




Cleaved caspase-1 (18 kDa)




β -actin (42 kDa)




Supplementary Figure S6. Original source blots corresponding to Figure 5A




p-ERK1/2 (42/44 kDa)




ERK1/2 (42/44 kDa)




p-NF- κ B p65 (65 kDa)




NF-κB p65 (65 kDa)




p-JAK2 (130 kDa)



JAK2 (130 kDa)



p-STAT3 (88 kDa)



STAT3 (88 kDa)



β -actin (42 kDa)

