

MiR-204-5p signalling modulates apoptosis and mitophagy-related changes after intracerebral hemorrhage in adult rats

Dongling Li

Affiliated Hospital of Jiujiang University

Xiaoqing Wen

Affiliated Hospital of Jiujiang University

wenqi Song

Affiliated Hospital of Jiujiang University

Yetong Ouyang

Jiangxi Provincial People's Hospital

Qinghua Luo

Second Affiliated Hospital of Nanchang University

Min Yin

Second Affiliated Hospital of Nanchang University

Yuqin Zhong

First Affiliated Hospital of Gannan Medical University

Zhengfang Qin

First People's Hospital of Fuzhou

Bing Bao

Affiliated Hospital of Jiujiang University

Xianming Cao

Affiliated Hospital of Jiujiang University

Moxin Wu

Jiujiang Clinical Precision Medicine Research Center

Zhiying Chen

zychen.jj@jju.edu.cn

Affiliated Hospital of Jiujiang University

Xiaoping Yin

Affiliated Hospital of Jiujiang University

Keywords: intracerebral hemorrhage, miR-204-5p, mitophagy-related changes, neuronal apoptosis, donepezil

Posted Date: October 28th, 2026

DOI: <https://doi.org/10.21203/rs.3.rs-11262375/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Additional Declarations: No competing interests reported.

MiR-204-5p signalling modulates apoptosis and mitophagy-related changes after intracerebral hemorrhage in adult rats

Dongling Li^{1,2+}, Xiaoqing Wen^{1,2+}, Wenqi Song^{1,2+}, Yetong Ouyang³, Qinghua Luo⁴, Min Yin⁴, Yuqin Zhong⁵, Zhengfang Qin⁶, Bing Bao^{1,2}, Xianming Cao^{1,2}, Moxin Wu^{2,7}, Zhiying Chen^{1,2#}, Xiaoping Yin^{1,2#}

¹ Department of Neurology, Affiliated Hospital of Jiujiang University, Jiujiang, Jiangxi, 332000, China

² Jiujiang Clinical Precision Medicine Research Center, Jiujiang, Jiangxi, 332000, China

³ Department of Neurology, Jiangxi Provincial People's Hospital, Nanchang, Jiangxi, 330000, China

⁴ Department of Neurology, The Second Affiliated Hospital of Nanchang University, Nanchang, Jiangxi, 330000, China

⁵ Department of Neurology, The First Affiliated Hospital of Gannan Medical College, Ganzhou, Jiangxi, 341000, China

⁶ Department of Neurology, First People's Hospital of Fuzhou, Fuzhou, Jiangxi, 344000, China

⁷ Department of Medical Laboratory, Affiliated Hospital of Jiujiang University, Jiujiang, Jiangxi 332000, China

+The first three authors contributed equally to this work.

#Correspondence authors: Xiaoping Yin and Zhiying Chen

Address correspondence to:

Xiaoping Yin, MD, PhD, Department of Neurology, Affiliated Hospital of Jiujiang University, Jiujiang, Jiangxi, 332000, China.

Email: xiaopingbuxiao@126.com

Zhiying Chen, MD, PhD, Department of Neurology, Affiliated Hospital of Jiujiang University, Jiujiang, Jiangxi, 332000, China.

Email: zychenjj@jjju.edu.cn

1 **Abstract**

2 **Background:** The survival of neurons around the hematoma is crucial for the recovery
3 of neurological function after spontaneous intracerebral hemorrhage (ICH). Neuronal
4 apoptosis and mitochondrial quality control disturbances are important factors for
5 secondary brain injury after ICH, but their molecular mechanisms have not been fully
6 elucidated. This study investigated the role of miR-204-5p in ICH and analyzed its
7 association with apoptotic and mitophagy-related signalling.

8 **Methods:** A collagenase-induced ICH model was established in adult male Sprague-
9 Dawley rats. miR-204-5p expression was assessed in perihematoma brain tissue and
10 in serum from patients with ICH. Neurological deficits, brain water content and blood-
11 brain barrier permeability were evaluated after intracerebroventricular administration
12 of a miR-204-5p agomir. Apoptosis-associated markers and mitophagy-related proteins
13 were assessed by western blotting, TUNEL staining, immunofluorescence and
14 transmission electron microscopy. Target prediction, luciferase reporter assays and
15 gene knockdown experiments were used to investigate TAOK1 and CAMK1. GATA3
16 promoter activity and the effects of donepezil pretreatment were also examined.

17 **Results:** miR-204-5p was reduced in perihematoma brain tissue and in serum from
18 patients with ICH, and lower serum levels were associated with higher NIHSS and
19 mRS scores in the available clinical samples. miR-204-5p overexpression improved
20 neurological deficits and reduced apoptosis-associated changes and mitophagy-related
21 molecular and ultrastructural changes after ICH. TAOK1 and CAMK1 were identified
22 as direct post-transcriptional targets of miR-204-5p by bioinformatic prediction,
23 expression analysis and dual-luciferase assays. TAOK1 knockdown reduced TUNEL-

positive signals, whereas CAMK1 knockdown attenuated mitochondrial ultrastructural abnormalities. GATA3 increased miR-204-5p promoter activity, and its expression was reduced after ICH. Donepezil pretreatment increased GATA3 expression and was associated with reduced apoptosis-associated and mitophagy-related changes.

Conclusions: Together, these findings suggest that the GATA3 – miR-204-5p – CAMK1/TAOK1 axis may contribute to neuronal injury after ICH. Mitophagic flux and the causal relationships among these components remain to be established. The clinical association and donepezil findings are preliminary and require further validation.

Keywords: intracerebral hemorrhage, miR-204-5p, mitophagy-related changes, neuronal apoptosis, donepezil

Introduction

Spontaneous intracerebral hemorrhage(ICH) accounts for 10 – 15% of strokes,with high morbidity,mortality,and disability[1·2]. With an ageing population, the lack of effective therapies to improve prognosis and functional recovery remains an important clinical challenge.

ICH involves primary mechanical compression and secondary injury[3]. Surgical hematoma removal has not consistently improved neurological outcomes, highlighting the importance of secondary injury,though its mechanisms remain complex.Survival of perihematoma neurons is critical for recovery[4].Numerous studies show that secondary injury induces substantial neuronal death around the hematoma[5-7],leading to severe deficits and poor prognosis,yet the specific mechanism is still unclear[8].

ICH induces excessive mitochondrial damage and neuronal apoptosis[9·10], both of which contribute to secondary brain injury. Mitophagy selectively removes damaged mitochondria and is an important component of mitochondrial quality control[11]; however, changes in mitophagy-related markers may reflect either increased mitochondrial turnover or impaired autophagic flux. Proper autophagy clears damaged organelles and proteins and can influence neuronal survival[12,13]. Previous studies have investigated the effects of modulating apoptosis and mitochondrial quality control in stroke and ICH[14]. Identifying molecular pathways that coordinately influence these processes may therefore provide additional insight into secondary brain injury.

MicroRNAs (miRNAs) are noncoding RNAs that regulate gene expression post-transcriptionally. Accumulating evidence links miRNA dysregulation to the pathological process of ICH[15,16]. Our group previously reported roles for miR-181c, miR-21-5p and miR-23a-3p in inflammatory or blood-brain barrier-related responses after ICH[15-17]. The present study extends this line of investigation to miR-204-5p, which showed a marked reduction in our previously generated ICH miRNA profile. Importantly, our data combine neuronal localization of miR-204-5p with two target-related branches involving TAOK1 and CAMK1, linking miR-204-5p to apoptosis-associated and mitochondrial quality-control phenotypes. We therefore investigated whether miR-204-5p contributes to secondary injury after ICH and explored upstream GATA3-associated regulation and the potential pharmacological relevance of donepezil.

Ethical approval

All animal experiments were reviewed and approved by the Ethics Committee of the Affiliated Hospital of Jiujiang University (approval number: 20200012). All animal-related experimental procedures were performed in accordance with the animal care guidelines of Jiujiang University.

Experimental animals

A total of 140 male Sprague-Dawley rats (18-week-old, weighing 500–600 g) were purchased from Beijing Weitong Lihua Laboratory Animal Technology Co., Ltd. These

animals were used as an adult rat model. Because only male adult rats were studied, the generalizability of the findings to females and to aged animals requires further investigation.

Materials and methods

Injecting agomir or virus and inducing a model of intracerebral hemorrhage

The rats were anaesthetized and fixed in a stereotaxic apparatus, the head skin was prepared and disinfected, and a median incision was made. The coordinates of the lateral ventricle (perpendicular to the bregma, 2 mm to the right, 1.2 mm to the back, and 4 mm deep) were used to connect the Hamilton microsyringe to the micropump. miR-204-5p agomir or scramble was injected at 0.5 μ l/min. One week later, a rat model of intracerebral hemorrhage was established using collagenase according to the method of Park et al. [18]. One day later, experiments such as behavioural experiments were performed. Similarly, CAMK1 knockdown lentivirus or TAOK1 knockdown lentivirus and corresponding control virus were injected in situ in the right basal ganglia coordinates (3 mm perpendicular to bregma, 0.2 mm anterior, 6 mm deep) for 3 weeks. After the intracerebral hemorrhage model was induced, the detection experiment was performed one day later.

Western blotting

Tissues or cells were lysed in RIPA buffer (Sigma - Aldrich), and protein concentrations were determined by BCA assay. Equal amounts of protein (20 μ g) were separated by

98 SDS-PAGE,transferred onto nitrocellulose membranes,and blocked with 3% BSA-
99 TBST for 1 h at room temperature.Membranes were then incubated overnight at 4° C
100 with primary antibodies against β -actin(Abcam,1:200),LC3II(Cell Signaling
101 Technology,1:1000),Parkin(CST,1:1000),PINK1(Abclonal,1:1000),Cleaved
102 caspase-3(CST,1:1000),total caspase-3(CST,1:1000), CAMK1(Abcam,1:5000),
103 TAOK1(CST,1:1000),GATA3(Abcam,1:1000),Cleaved PARP(CST,1:1000),or
104 Bak(CST,1:1000).After three TBST washes(10 min each),membranes were incubated
105 with HRP-conjugated secondary antibodies for 1 h at room temperature in the
106 dark,followed by another three TBST washes.Immunoreactive bands were visualized
107 using an Odyssey imaging system and quantified by densitometry.

108 **RT-qPCR**

109 According to the manufacturer's instructions,the miRNA Isolation
110 Kit(TIANGEN,Beijing,China).Lot:DP 501)extracts total RNA from rat brain
111 tissue,human serum specimens or HEK 293 T cells.Total RNA was reverse transcribed
112 for miRNA using the miRcute Enhanced miRNA First-Strand cDNA Synthesis
113 Kit(TIANGEN,Beijing,China,Lot KR 211).Total RNA was reverse transcribed for
114 mRNA using the QuantiNova™ Reverse transcription
115 Kit(QIAGEN,Germany,ID:205410).The SYBR Green I fluorescent dye
116 method(SYBR® PrimeScript™ RT-PCR Kit(Takara Biotechnology Co.)was used for
117 PCR detection.Amplification of miRNA/mRNA was performed on an Applied
118 Biosystems StepOne™ Real-Time PCR System(Applied Biosystems).The cycling

conditions were according to the manufacturer's instructions. U6 snRNA/GAPDH served as the internal control for RNA quantification. The relative concentration was calculated by the $2^{-\Delta\Delta CT}$ method.

Brain slice preparation

In short, the rat was perfused with normal saline, fixed with 4% paraformaldehyde and then treated with a gradient of 10%, 20%, 30%, and 30% sucrose solutions. Tissues were dehydrated by gradient sucrose solutions, embedded in OCT, and then placed in a microtome, and the corresponding parameters were set. Brain slices were stored in PBS-glycerol at 4 °C for subsequent experiments.

Immunofluorescence

The brain slices were rinsed 3 times in PBS at room temperature for 10 min each time; PBS solution of 0.5% Triton X-100 was added, and the membrane was broken by shaking slowly for 15 mins. LC 3II (Cell Signaling Technology, 1:2000) was incubated at 4 °C for 24-28 hours and then rinsed with PBS 3 times for 10 min each time. Thereafter, secondary antibody was added and incubated for 2 hours at room temperature in the dark, after which the sections were rinsed with PBS 3 times for 10 min each time. Next, DAPI was added to stain the nuclei for 10 min at room temperature in the dark, then rinsed 3 times with PBS, attached to a glass slide, air-dried naturally in the dark, and mounted with PBS-glycerol.

TUNEL

Brain slices were rinsed 3 times in PBS at room temperature for 10 min each, and 3% BSA was added to a shaker and slowly shaken for 10 mins. Then, the slices were rinsed 3 times with PBS for 5 min followed by 0.1% Triton X-100 containing 0.1% sodium citrate solution, shaken slowly for 8 min on a shaker in cold storage at 4 °C, rinsed with PBS 3 times for 5 min each time, and then attached to a glass slide. TUNEL detection solution was prepared according to the TUNEL kit (Beyotime, C1089) (5 µl TdT enzyme + 45 µl fluorescent labelling solution), TUNEL detection solution was added dropwise, the slides were placed in a wet box, incubated for 1 hour at 37 °C in the dark, and rinsed with PBS 3 times for 5 min each. Thereafter, DAPI was added to stain cell nuclei for 10 min at room temperature in the dark and rinsed 3 times with PBS every 5 min. Brain slices were naturally air-dried under dark conditions and mounted with PBS-glycerol.

Fluorescence in situ hybridization

RNase-free consumables and reagents were used. At room temperature, the brain slice was rinsed in PBS containing 0.1% RNase Remover, a PBS solution of 0.5% Triton X-100 was added, and the shaking table was slowly shaken for 15 min. Then, the sections were digested at 37°C for 1 min followed by fixation with 1% paraformaldehyde containing 0.1% DEPC for 15 min. The brain slices were placed into a wet box, prehybridization solution was added dropwise at 38°C for 4 hours, hybridization solution was added dropwise, and the sections were hybridized overnight at 42°C. After hybridization, the cells were washed twice with 2x SSC at 37°C for 5 min each and

then washed once with 0.5x SSC at 37°C for 15 min and twice with 0.2x SSC at 37 °C for 15 min. The blocking solution was added dropwise at 37 °C for 30 min, and target-protein primary antibodies (GFAP (Abcam 1:1000), Iba-1 (Abcam 1:1000), NeuN (Abcam 1:1000), TOM20 (Cell Signaling Technology, 1:400)) were added overnight at 4 °C. Biotinylated mouse anti-digoxigenin was added dropwise at 37 °C for 60 mins, then rinsed 4 times with PBS for 5 min each; 1% SABC-FITC-containing PBS solution was added dropwise for 30 min at 37 °C after rinsing with PBS. Next, target protein secondary antibody was added, incubated at 37 °C for 60 min, rinsed with PBS, and then DAPI was added to stain nuclei for 10 min at room temperature in the dark and rinsed with PBS, air-dried, and mounted with PBS-glycerol.

TEM observation

Rats were anaesthetized, and the volume of rice grains was taken from the fresh brain tissue around the foci, quickly placed under an electron microscope for fixation, and fixed at 4 °C for 4 hours. Then, the cells were rinsed with PBS 3 times for 15 mins each, followed by the addition of 1% osmic acid-containing PBS solution at room temperature, and fixed for 2 hours. The cells were rinsed 3 times with PBS for 15 min each. Then, 50%-70%-80%-90%-95%-100%-100% alcohol-100% acetone-100% acetone dehydration was added successively, each time for 15 min. Then, 1/2 acetone in 812 embedding medium was added for 2-4 hours, infiltrated in 1/3 acetone-containing 812 embedding medium overnight, in pure 812 embedding medium for 5-8 hours, and poured into pure 812 embedding medium. The samples were inserted into the

embedding plate, placed in a 37 °C oven overnight, polymerized at 60 °C for 48 hours and cut into 70 nm ultrathin sections in an ultramicrotome. Then, the sections were double-stained with uranium lead (2% uranyl acetate saturated alcohol solution, lead citrate, stained for 15 min each), dried at room temperature overnight and finally observed under a transmission electron microscope, and images were collected for analysis.

Cell transfection

The cells were cultured in a well plate to 70% density for transfection, and the medium was changed 2 h before transfection. Lipo3000 reagent was diluted with Opti-MEM and mixed, the plasmid was diluted with Opti-MEM, and P3000 reagent was added and mixed (transfected siRNA) or (miRNA without P3000 reagent). The diluted Lipo3000 was mixed with the plasmid diluent in equal proportions and incubated at room temperature for 10 min. Then, the incubated mixture was added to the cells and cultured for 48 hours before performing subsequent experiments.

Dual luciferase

For the luciferase reporter assay, psiCHECK2 constructs containing the wild-type (WT) or mutant (MUT) 3'UTRs of TAOK1 or CAMK1 were generated based on the predicted miR-204-5p binding sites and transfected into HEK293T cells together with miR-204-5p mimic or negative control. After 48 h, cells were harvested and luciferase activity was measured using the Dual Luciferase Reporter Assay System (Promega, WI, USA). Each transfection was performed in triplicate.

Corner experiment

Two plates forming a 30° corner with a small gap at the junction were used. Rats placed facing the corner naturally approached it; stimulated by whiskers, they turned to one side. Only complete turns were counted, and the number of right turns out of 10 trials was recorded, with intervals ≤ 2 min.

Determination of water content in brain tissue

Brain water content was measured by wet-dry weight method. Ipsilateral and contralateral peri-lesional tissues were collected after anesthesia, surface water blotted, and wet weight immediately weighed. Tissues were dried at 100° C for 24h to obtain dry weight. Water content = (wet weight - dry weight) / wet weight $\times$ 100%.

Blood-brain barrier permeability test

Evans blue assay was performed. After anesthesia, 2% Evans blue (4 ml/kg) was injected via tail vein. Two hours later, rats were perfused with saline to remove intravascular dye. Brain tissues were homogenized in 50% trichloroacetic acid, centrifuged, and supernatant diluted 1:3 in ethanol. Absorbance was measured at 620 nm.

Statistical methods

For statistical analysis, SPSS 22 software (IBM, Armonk, USA) was used, and the results are expressed as the mean $\pm$ standard error (mean $\pm$ SEM). $P < 0.05$ indicates a significant difference between the two groups of results. When analysing the data of the two groups, if the variance was homogeneous, the test was used to analyse the significant difference, and if the variance was unequal, Mann-Whitney

analysis was used. To analyse multiple groups of sample data, one-way ANOVA or two-way ANOVA was used for significant difference analysis if the variance was homogeneous, and the Kruskal–Wallis method was used to approximate the variance analysis if the variance was unequal.

Results

3.1 Reduction of miR-204-5p is associated with neurological deficits after ICH

We first examined the top 10 downregulated miRNAs identified in our previous ICH miRNA profiling study (Ouyang Y et al., 2019) (Fig. 1A). Four miRNAs—miR-501-3p, miR-409-3p, miR-204-5p and miR-150-5p—were significantly decreased in ICH rats (Fig. 1B). Among these candidates, miR-204-5p showed the largest reduction and was therefore selected for further study (Supplementary Table S1). We then examined miR-204-5p expression and neurological outcomes at different time points after ICH. Neurological impairment, assessed by mNSS and the corner test, was greatest at 24 h and gradually improved from days 1 to 7 (Fig. 1C–D), whereas miR-204-5p was most strongly reduced at 24 h and gradually recovered thereafter (Fig. 1E). Serum miR-204-5p in patients with spontaneous ICH showed a similar temporal pattern (Fig. 1F). In the available clinical samples, serum miR-204-5p was inversely associated with NIHSS and mRS scores at the assessed time point (Fig. 1G–H). These findings support an association between reduced miR-204-5p expression and ICH severity, but do not establish that circulating miR-204-5p originates from perihematoma neurons.




Fig.1

3.2 miR-204-5p overexpression attenuates apoptosis-associated and mitophagy-related changes after ICH

We used TargetScan, miRDB and miRWalk to predict potential targets of miR-204-5p (Supplementary Fig. S1A–D) and analysed the intersection of the predicted target genes.

250 GO enrichment analysis showed enrichment of biological processes related to cell death,
251 including apoptosis (GO:0043525, $P = 4.38 \times 10^{-4}$), whereas the enrichment for
252 autophagy did not reach conventional statistical significance (GO:0010506, $P = 0.058$)
253 (Supplementary Fig. S1E – G). (Supplementary Fig. S1E–G). Using a specific probe
254 for miR-204-5p, we co-labelled miR-204-5p with NeuN, GFAP and Iba-1 and found
255 that miR-204-5p was predominantly detected in neurons and was less abundant in
256 astrocytes and microglia (Fig. 1J). miR-204-5p signal was also observed in the
257 cytoplasm and mitochondrial compartment of neurons. We next examined apoptosis-
258 associated and mitophagy-related changes after miR-204-5p supplementation. ICH
259 increased LC3-II, Parkin, PINK1, cleaved PARP and Bak, together with increased
260 TUNEL and LC3-II signals (Fig. 2A–D). miR-204-5p agomir reduced these changes.
261 Transmission electron microscopy showed more autophagosome-like structures and
262 mitochondrial structural abnormalities after ICH, whereas miR-204-5p treatment was
263 associated with preservation of mitochondrial morphology (Fig. 2E). miR-204-5p
264 agomir also improved mNSS and corner-test performance and reduced brain water
265 content and Evans blue leakage (Fig. 2F–I). Collectively, these data indicate that miR-
266 204-5p overexpression is associated with improved neurological outcomes and reduced
267 apoptosis-associated and mitophagy-related changes. Because autophagic flux was not
268 directly measured, these findings should not be interpreted as definitive evidence of
269 increased or decreased mitophagic flux.




Fig.2

3.3 miR-204-5p directly targets CAMK1 and TAOK1

To identify candidate functional targets of miR-204-5p, we combined target prediction with pathway analysis and expression screening. Among the predicted candidates, 11 genes were present in the intersection of the selected target lists (Supplementary Fig.

276 S1I). PIP4K2B and TAOK1 were associated with apoptosis-related pathways, whereas
277 RPS6KA3 and CAMK1 were associated with mitochondrial quality-control pathways.
278 All four proteins were increased after ICH, while TAOK1 and CAMK1 mRNA levels
279 were comparable between ICH and sham groups (Fig. 3A–B), suggesting post-
280 transcriptional regulation. Conserved miR-204-5p binding sites were identified in the
281 3'UTRs of TAOK1 and CAMK1 (Fig. 3E,G). In HEK293T cells, miR-204-5p mimic
282 reduced luciferase activity from WT, but not MUT, TAOK1 and CAMK1 3'UTR
283 reporters (Fig. 3F,H). miR-204-5p agomir also reduced TAOK1 and CAMK1 protein
284 levels in perihematoma tissue (Fig. 3I–J). These results support direct post-
285 transcriptional regulation of TAOK1 and CAMK1 by miR-204-5p. However, rescue
286 experiments were not performed; therefore, TAOK1 and CAMK1 should be regarded
287 as direct targets rather than definitively established mediators of all miR-204-5p effects.




Fig.3

3.4 TAOK1 and CAMK1 knockdown phenocopies selected effects of miR-204-5p after ICH

To examine the functional contribution of TAOK1 and CAMK1, we injected AAV vectors carrying shRNAs targeting TAOK1 or CAMK1, or corresponding control vectors, into the basal ganglia before ICH induction. TAOK1 knockdown reduced

295 TUNEL-positive signals, whereas CAMK1 knockdown was associated with
296 preservation of mitochondrial ultrastructure and fewer abnormal autophagosome-like
297 structures (Fig. 4C–E). Neurological deficits were improved by TAOK1 or CAMK1
298 knockdown and were further reduced when both targets were inhibited (Fig. 4F–G).
299 These results show that TAOK1 and CAMK1 knockdown can phenocopy selected
300 protective effects associated with miR-204-5p overexpression. However, because
301 target re-expression experiments were not performed, the data do not establish epistatic
302 mediation or exclude contributions from additional miR-204-5p targets.




Fig.4

3.5 GATA3-associated regulation of miR-204-5p and exploratory effects of donepezil

To explore upstream regulation of miR-204-5p, we cloned 1-kb, 2-kb and 3-kb upstream regions into a pGL3 reporter (Supplementary Fig. S1J). The 2-kb construct showed the strongest promoter activity (Supplementary Fig. S1K). RegRNA and

LASAGNA prediction identified potential binding motifs for GATA1, GATA3, AP-1, Cart-1, NF-1 and PU.1; among the candidates tested, GATA3, but not the other tested factors, increased miR-204-5p promoter reporter activity (Supplementary Fig. S1L and Fig. 5A). Mutation of the tested GATA3-associated region reduced miR-204-5p promoter activity relative to the corresponding construct (Fig. 5B). GATA3 mRNA and protein levels were reduced after ICH (Fig. 5C–E). These findings support a regulatory association between GATA3 and miR-204-5p transcription, but direct chromatin binding was not assessed by ChIP. We next screened four compounds reported to influence GATA3-related signalling and found that only donepezil increased GATA3 expression in the tested cell system (Fig. 5F). Donepezil was administered intraperitoneally for 14 days before ICH induction. Pretreatment partially improved neurological scores, restored GATA3 mRNA expression and reduced apoptosis-associated and mitophagy-related molecular changes (Fig. 5G–L). Because donepezil was administered before ICH and no miR-204-5p loss-of-function experiment was performed, these data provide pharmacological support but do not establish that donepezil neuroprotection requires the GATA3–miR-204-5p pathway.




Fig.5

4Discussion

Our findings support a regulatory model in which miR-204-5p is reduced after ICH and its supplementation is associated with improved neurological outcomes and reduced apoptosis-associated and mitophagy-related changes. We identified TAOK1 and CAMK1 as direct post-transcriptional targets and found that their knockdown

phenocopied selected protective effects. We further identified GATA3 as a candidate upstream regulator of miR-204-5p promoter activity and found that donepezil pretreatment increased GATA3 expression. Importantly, the present study should be interpreted as a mechanistic framework rather than a fully established linear pathway because mitophagic flux, target-rescue/epistasis and pharmacological dependency were not directly tested.

4.1 Reduced miR-204-5p after ICH is associated with neurological impairment

Our group has previously reported miR-181c, miR-21-5p and miR-23a-3p as regulators of inflammatory or blood-brain barrier-related responses after ICH[15-17]. The present study addresses a different biological question by focusing on miR-204-5p and its relationship with apoptosis-associated and mitochondrial quality-control phenotypes. miR-204-5p was reduced in perihematoma brain tissue and in serum from patients with ICH, and lower serum levels were associated with higher NIHSS and mRS scores in the available samples. The neuronal localization of miR-204-5p and the effects of miR-204-5p supplementation provide a rationale for further investigation. Nevertheless, serum miR-204-5p cannot be assumed to reflect neuronal miR-204-5p directly because the present study did not examine cerebrospinal fluid or neuron-derived extracellular vesicles.

4.2 miR-204-5p is associated with apoptosis and mitochondrial quality-control changes after ICH

Excessive neuronal apoptosis contributes to secondary injury after ICH, while mitochondrial damage caused by oxidative stress, inflammation and excitotoxicity can further compromise neuronal survival[24,27]. Mitophagy is an important component of mitochondrial quality control, but its biological effect depends on the balance between mitochondrial clearance and degradation capacity. Our increases in LC3-II, Parkin and PINK1, together with electron-microscopic changes, indicate altered autophagy/mitochondrial quality-control signalling after ICH. However, these measurements alone cannot distinguish increased mitophagy from impaired autophagic flux. The reduction of these markers and improvement in mitochondrial morphology after miR-204-5p supplementation therefore support an association with improved mitochondrial homeostasis, but not a definitive conclusion about the direction of mitophagic flux. Future studies using flux reporters, lysosomal inhibition, mitochondrial-lysosomal colocalization and functional mitochondrial measurements will be required.

4.3 TAOK1 and CAMK1 are direct post-transcriptional targets of miR-204-5p

By combining target prediction, pathway analysis and expression screening, we identified TAOK1 and CAMK1 as candidate targets of miR-204-5p. Conserved 3'UTR binding sites, WT/MUT reporter assays and reduced endogenous protein levels after miR-204-5p agomir support direct post-transcriptional regulation. Knockdown experiments further showed that TAOK1 and CAMK1 contribute to distinct phenotypic components: TAOK1 knockdown was associated with reduced TUNEL signals,

whereas CAMK1 knockdown was associated with improved mitochondrial ultrastructure. These findings are consistent with previous reports linking TAOK1 to apoptosis and CAMK-related proteins to mitochondrial quality control[30,31]. Nevertheless, target re-expression using 3'UTR-less constructs was not performed. Thus, the present data support target regulation and phenotypic contribution but do not prove that TAOK1 and CAMK1 are the exclusive or obligatory mediators of miR-204-5p-mediated protection.

4.4 GATA3 and donepezil: a hypothesis-generating upstream regulatory component

GATA3 is a transcription factor involved in cell differentiation, apoptosis and immune regulation[32-34]. Our promoter-reporter experiments suggest that GATA3 can increase miR-204-5p promoter activity, and GATA3 expression was reduced after ICH. These findings support GATA3 as a candidate upstream regulator, but direct promoter occupancy was not demonstrated by ChIP and the effects of GATA3 loss- or gain-of-function on endogenous pri-miR-204 were not examined. We therefore regard the GATA3–miR-204-5p relationship as supportive but not definitive. Donepezil pretreatment increased GATA3 expression and was associated with improved neurological outcomes in our adult rat model. However, the treatment was initiated before ICH, the therapeutic window after ICH was not assessed, and the dependence of the effect on miR-204-5p was not tested. Molecular docking suggested an interaction with GATA3, but this interaction was not validated by mutational or biochemical

experiments. Accordingly, donepezil should be considered a hypothesis-generating pharmacological observation rather than evidence for a miR-204-5p-dependent therapeutic mechanism.

4.5 Study limitations

Several limitations should be considered. First, the present study used 18-week-old adult rats and therefore does not specifically represent aged ICH. Second, although changes in LC3-II, PINK1, Parkin and mitochondrial ultrastructure were observed, mitophagic flux was not directly assessed. Therefore, these findings should be interpreted as mitophagy-related changes rather than definitive evidence of altered mitophagy flux. Third, although TAOK1 and CAMK1 were identified as direct targets of miR-204-5p and their knockdown reproduced selected effects of miR-204-5p overexpression, rescue or epistasis experiments were not performed to establish a definitive causal relationship. Similarly, the regulatory relationship between GATA3 and miR-204-5p was not further validated by ChIP or endogenous transcriptional assays. Finally, the effects of donepezil were examined only in a pretreatment setting, and whether its effects depend on miR-204-5p remains to be determined. Further studies using aged animals, direct mitophagy-flux assays, rescue experiments and post-ICH treatment paradigms are needed to validate these findings.

Abbreviations

ICH:intracerebral hemorrhage,BBB:blood brain barrier,KEGG:Kyoto Encyclopedia of Genes and Genomes,CAMK1:calcium/calmodulin-dependent protein kinase 1,CAMKs:calcium/calmodulin-dependent protein kinases,TAOK1:Thousand and One Kinase 1

Figure Legends

Figure 1:Expression and subcellular location of miR-204-5p,and its relationship with neurological deficits.(A)Heat map showed the 10 downregulated candidate miRNAs after ICH with $|\log_2 \text{fold change}| > 2$ and $p < 0.05$ as the filter condition from the expression profile of miRNAs in perifocal brain tissue of rats between the ICH and sham group.(B)RT-qPCR was performed to validate the expression of the 10 miRNAs around the perifocal tissue,suggested that 4 miRNAs - miR-501-3p,miR-409-3p,miR-204-5p,miR-150-5p were found significantly decreased.Using the $2^{-\Delta\Delta CT}$ method with U6 expression levels as an internal reference.All results are presented as mean $\pm$ SEM of five independent experiments.(C)We established ICH rat models at different time points on day 1,day 3 and day 7,evaluated neurological damage like motor nerves,sensory function,reflex ability,abnormal activity with mNSS scores.(D)Assessed the degree of hemiplegia with right turn behavior experiment.(E)Detection of miR-204- 5p expression level in each group.(F)miR-204- 5p expression in the serum of patients at different time points after the onset

434 of ICH.(G-H)Correlation between the expression of miR-204-5p and ICH patient's
 435 NIHSS and mRS score.(I)Use
 436 TargetScan(<http://www.targetscan.org/>),miRDB(<http://mirdb.org/>)and
 437 miRWalk(<http://mirwalk.umm.uni-heidelberg.de/>)database to predicted the target
 438 genes of these four miRNAs,took target genes intersection and enriched them with GO
 439 function ,suggesting that only miR-204-5p target genes are mainly involved in the
 440 regulation of cell death biological processes including neuron
 441 apoptosis(GO:0043525,p=4.38E-04)and
 442 autophagy(GO:0010506,p=0.05800458).(J)The subcellular distribution of miR-204-
 443 5p was visualized by RNA fluorescence in situ
 444 hybridization(FISH).astrocytes,microglia,neurons and mitochondria were identified by
 445 GFAP,Iba-1 and NeuN,Tom20 antibody,respectively.miR-204-5p is primarily
 446 localized in the nucleus and mitochondria of neurons.Scale bar=20um.***p <
 447 0.001.****p<0.0001.

448

449 **Figure 2:Neuron apoptosis and mitophagy disorders increased 24h post-ICH,miR-**
 450 **204-5p overexpression rescued these change and improved neurological**
 451 **deficits.(A)**Western blotting detection of LC3II,Parkin,PINK1,Cleaved-PARP and Bak
 452 expression in sham-operated ,ICH group,Scramble+ICH group and miR-204-5p
 453 agomir(204Ago)+ICH group.(B)Quantitative analysis of western blotting
 454 results.(C)Apoptosis was visualized by the TUNEL Assay,and the TUNEL-

positive(red)cells were quantitatively analyzed.Scale
bar=20um.(D)Immunofluorescence detection and quantification of LC3II levels in
brain sections.Scale bar=20um.(E)Ultrastructure of neuronal mitochondria in perifocal
brain tissue was observed by transmission electron microscope(TEM).Mitochondrial
autophagosomes increased,cristae structures of some mitochondria swollen and
disappeared,mitochondrial structure was seriously damaged in ICH or Scramble+ICH
group,while that of 204Ago+ICH group was restored.Scale
bar=500nm .**p<0.01,****p<0.0001.The neurological damage of rats in each group
was evaluated by mNSS score(F)and corner behavioral test(G),n=10.(H)Dry-wet
weight method was used to detect the brain water content of the ipsilateral and
contralateral lesions of the rats in each group,n=5.(I)Evans blue assay to detect the
permeability of the perifocal blood-brain barrier,n=5.

Figure 3:miR-204-5p directly targets TAOK1 and CAMK1.(A)mRNA
and(B)protein expression levels of apoptosis-related genes(PIP4K2B,TAOK1)and
mitophagy-related genes(RPS6KA3,CAMK1)were detected.The mRNA expression
levels of TAOK1 and CAMK1 showed no significant change but protein expression
increased when ICH versus sham group,implying post-transcriptional
regulation.(C)Western blotting measure and(D)statistical analysis of TAOK1 and
CAMK1 protein expression.(E)Predictive binding sites in TAOK1 3'UTR on
mammalian miR-204-5p by bioinformatics analysis.(F)Constructed luciferase reporter

plasmid,psi-CHEK2 containing wild-type(WT-TAOK1)or mutant(MUT-TAOK1)3'UTR of TAOK1,transfected them to HEK293T cells with the mimic or negative control(NC)of miR-204-5p respectively for 48 hours,then performed luciferase reporter assay.Bar graph shows relative luciferase activities.(G)Putative binding sites in CAMK1 3'UTR on miR-204-5p among mammals.(H)Similar procedure to verify the direct bind relationship between CAMK1 and miR-204-5p.Relative luciferase activities were measured.(I)Protein expression of TAOK1 and CAMK1 around hematoma of ICH rats after injecting miR-204-5p agomir or negative scramble.(J)Quantification analysis of TAOK1 and CAMK1 protein levels.**p<0.01,***p<0.001,****p<0.0001.

Figure 4 :Knockdown of TAOK1 inhibits neuronal apoptosis,and knockdown of CAMK1 improves mitophagy disorder,protecting neurological function.(A)TAOK1 mRNA expression in rats 24h post-ICH after TAOK1 knockdown virus,corresponding control virus or TAOK1 knockdown virus + CAMK1 knockdown virus injected 14 days later.(B)CAMK1 mRNA expression 24h post-ICH with injection of CAMK1 knockdown virus,corresponding control virus or TAOK1 knockdown virus + CAMK1 knockdown virus for 14 days.(C)TUNEL staining apoptotic cells around the focus of rats in the TAOK1 knockdown virus + ICH group and control virus + ICH group,Scale bar=50um.(D)Counts the number of TUNEL positive(red)cells.(E)Ultrastructure of neuron mitochondria in the perihemetoma of

ICH-induced rats with injection of CAMK1 knockdown virus or control virus. Scale bar=500nm.(E)mNSS score and(F)Corner test to evaluate the neurological function damage of rats in each group.*p<0.05,**p<0.01,****p<0.0001.

Figure5 Donepezil promoted GATA-3 expression and drove miR-204-5p to work.(A)Predicted the transcription factors that might bind to the promoter region of miR-204-5p,constructed plasmids containing binding domain or its homeodomain deletion mutation(HDD)of the transcription factor,and transfected it to 293T cells with the plasmid containing miR-204-5p promoter for 48 hours, then use luciferase assay to detect fluorescence intensity.Only the group containing GATA-3 binding domain in relative light unit(RLU)pronouncedly enhanced compared with its mutant group.(B)Transfected the plasmid containing GATA-3 binding domain or its HDD into 293T cells,RT-qPCR detected miR-204-5p expression levels.(C)GATA-3 mRNA and(D)protein levels in sham group and ICH group.(E)Quantitative analysis of western blotting results.(F)GATA-3 mRNA expression after 4 drugs(Gemfibrozil,Roflumilast,Sinomenine and Donepezil)to 293t cells separately.Injected Donepezil or normal saline into rats once a day intraperitoneally for 14 days,then established ICH models,mNSS score(G)and Corner test(H)to evaluate the neurological deficits of rats in each group.(I)Detection expression of GATA-3 mRNA.(J)Western blotting detection of GATA3,Parkin,PINK1,Cleaved-PARP and Bak expression and(K)Quantitative analysis.(L)Apoptosis was visualized by the

518 TUNEL Assay, and the TUNEL-positive (red) cells were quantitatively analyzed. Scale
519 bar=50um. **p<0.01, ***p<0.001, ****p<0.0001

520 **Declarations**

521 **Ethics approval and informed consent**

522 This study was approved by the Ethics Committee of the Affiliated Hospital of Jiujiang
523 University. Written informed consent was obtained from all the participants.

524 **Consent for publication**

525 All authors and patients have given their consent for publication.

526 **Data availability**

527 Please contact the corresponding author for data requests.

528 **Funding**

529 This study was sponsored by the National Key R&D Program(2018FY100903 to
530 XPY),the National Natural Science Foundation of China(81960221 to XPY,82260454
531 to BB,82260249 to XPY),Jiangxi Provincial Health Commission Science and
532 Technology Plan project(202311506 to ZYC),Jiangxi Provincial Administration of
533 Traditional Chinese Medicine science and technology plan project(2022A322 to
534 ZYC),Youth Foundation of Natural Science Foundation of Jiangxi
535 Province(20224BAB216045 to ZYC).

536 **Competing interests**

537 The authors declare that there are no conflicts of interest regarding the publication of
538 this paper.

539 **Authors' contributions**

Xiao-ping Yin and Zhiying Chen conceived and designed the work. Dongling Li, Yetong Ouyang, Qinghua Luo and Min Yin conducted the experiments. Dongling Li, Xiaoqing Wen, Manqing Zhang and Yuqin Zhong analysed and interpreted the data. Bing Bao, Xianming Cao, Zhengfang Qin and Moxin Wu advised on the experimental design and contributed materials and animals. Xiaoping Yin and Zhiying Chen prepared the manuscript, while Song Wenqi handled the submission. All authors revised the manuscript and approved the final version.

Acknowledgements

We sincerely thank Prof. Lingqiang Zhu and the staff of the team from Tongji Medical College, Huazhong University of Science and Technology. We sincerely thank the Jiujiang Precision Clinical Medicine Research Center staff and the students Weixin Zhou, Zhi Tu, and Jinming Ma.

References:

1. Anderson, C.S.; Heeley, E.; Huang, Y.; Wang, J.; Stapf, C.; Delcourt, C.; Lindley, R.; Robinson, T.; Lavados, P.; Neal, B. et al. Rapid blood-pressure lowering in patients with acute intracerebral hemorrhage. *N. Engl. J. Med.* **2013**, *368*, 2355-2365, doi:10.1056/NEJMoa1214609.
2. van Asch, C.J.; Luitse, M.J.; Rinkel, G.J.; van der Tweel, I.; Algra, A.; Klijn, C.J. Incidence, case fatality, and functional outcome of intracerebral haemorrhage over time, according to age, sex, and ethnic origin: a systematic review and meta-analysis. *Lancet Neurol.* **2010**, *9*, 167-

- 176,doi:10.1016/S1474-4422(09)70340-0.
3. Ironside,N.;Chen,C.J.;Ding,D.;Mayer,S.A.;Connolly,E.J.Perihematomal Edema After Spontaneous Intracerebral Hemorrhage.*Stroke* **2019**,50,1626-1633,doi:10.1161/STROKEAHA.119.024965.
4. Keep,R.F.;Hua,Y.;Xi,G.Intracerebral haemorrhage:mechanisms of injury and therapeutic targets.*Lancet Neurol.***2012**,11,720-731,doi:10.1016/S1474-4422(12)70104-7.
5. Fang,Y.;Tian,Y.;Huang,Q.;Wan,Y.;Xu,L.;Wang,W.;Pan,D.;Zhu,S.;Xie,M.Deficiency of TREK-1 potassium channel exacerbates blood-brain barrier damage and neuroinflammation after intracerebral hemorrhage in mice.*J.Neuroinflamm.***2019**,16,96,doi:10.1186/s12974-019-1485-5.
6. Gong,C.;Boulis,N.;Qian,J.;Turner,D.E.;Hoff,J.T.;Keep,R.F.Intracerebral hemorrhage-induced neuronal death.*Neurosurgery* **2001**,48,875-882,882-883,doi:10.1097/00006123-200104000-00037.
7. Rahman,M.S.;Yang,J.;Luan,Y.;Qiu,Z.;Zhang,J.;Lu,H.;Chen,X.;Liu,Y.Attenuation of Acute Intracerebral Hemorrhage-Induced Microglial Activation and Neuronal Death Mediated by the Blockade of Metabotropic Glutamate Receptor 5 In Vivo.*Neurochem.Res.***2020**,45,1230-1243,doi:10.1007/s11064-020-03006-1.
8. Hua,Y.;Schallert,T.;Keep,R.F.;Wu,J.;Hoff,J.T.;Xi,G.Behavioral tests after intracerebral hemorrhage in the rat.*Stroke* **2002**,33,2478-2484,doi:10.1161/01.str.0000032302.91894.0f.
9. Wang,Z.;Zhou,F.;Dou,Y.;Tian,X.;Liu,C.;Li,H.;Shen,H.;Chen,G.Melatonin Alleviates Intracerebral Hemorrhage-Induced Secondary Brain Injury in Rats via Suppressing Apoptosis,Inflammation,Oxidative Stress,DNA Damage,and Mitochondria Injury.*Transl.Stroke Res.***2018**,9,74-91,doi:10.1007/s12975-017-0559-x.

- 582 10. Wu,X.;Cui,W.;Guo,W.;Liu,H.;Luo,J.;Zhao,L.;Guo,H.;Zheng,L.;Bai,H.;Feng,D.et al.Acrolein
583 Aggravates Secondary Brain Injury After Intracerebral Hemorrhage Through Drp1-Mediated
584 Mitochondrial Oxidative Damage in Mice.*Neurosci.Bull.***2020**,36,1158-1170,doi:10.1007/s12264-
585 020-00505-7.
- 586 11. Gustafsson,A.B.;Dorn,G.N.Evolving and Expanding the Roles of Mitophagy as a Homeostatic and
587 Pathogenic Process.*Physiol.Rev.***2019**,99,853-892,doi:10.1152/physrev.00005.2018.
- 588 12. Fricker,M.;Tolkovsky,A.M.;Borutaite,V.;Coleman,M.;Brown,G.C.Neuronal Cell
589 Death.*Physiol.Rev.***2018**,98,813-880,doi:10.1152/physrev.00011.2017.
- 590 13. Liu,P.;Yu,X.;Dai,X.;Zou,W.;Yu,X.;Niu,M.;Chen,Q.;Teng,W.;Kong,Y.;Guan,R.et al.Scalp
591 Acupuncture Attenuates Brain Damage After Intracerebral Hemorrhage Through Enhanced
592 Mitophagy and Reduced Apoptosis in Rats.*Front.Aging*
593 *Neurosci.***2021**,13,doi:10.3389/fnagi.2021.718631.
- 594 14. Chen,S.;Peng,J.;Sherchan,P.;Ma,Y.;Xiang,S.;Yan,F.;Zhao,H.;Jiang,Y.;Wang,N.;Zhang,J.H.et
595 al.TREM2 activation attenuates neuroinflammation and neuronal apoptosis via PI3K/Akt
596 pathway after intracerebral hemorrhage in mice.*J.Neuroinflamm.***2020**,17,168,doi:10.1186/s12974-
597 020-01853-x.
- 598 15. Ouyang,Y.;Li,D.;Wang,H.;Wan,Z.;Luo,Q.;Zhong,Y.;Yin,M.;Qing,Z.;Li,Z.;Bao,B.et al.MiR-21-
599 5p/dual-specificity phosphatase 8 signalling mediates the anti-inflammatory effect of haem
600 oxygenase-1 in aged intracerebral haemorrhage rats.*Aging Cell*
601 **2019**,18,e13022,doi:10.1111/accel.13022.
- 602 16. Yin,M.;Chen,Z.;Ouyang,Y.;Zhang,H.;Wan,Z.;Wang,H.;Wu,W.;Yin,X.Thrombin-induced,TNFR-

dependent miR-181c downregulation promotes MLL1 and NF-kappaB target gene expression in human microglia.*J.Neuroinflamm.***2017**,14,132,doi:10.1186/s12974-017-0887-5.

17. Hu,Y.L.;Wang,H.;Huang,Q.;Wang,G.;Zhang,H.B.MicroRNA-23a-3p promotes the perihematoma edema formation after intracerebral hemorrhage via ZO-1.*Eur.Rev.Med.Pharmacol.Sci.***2018**,22,2809-2816,doi:10.26355/eurev_201805_14980.

18. Park,J.;Kim,J.Y.;Choi,S.K.;Kim,J.Y.;Kim,J.H.;Jeon,W.B.;Lee,J.E.Thermo-sensitive assembly of the biomaterial REP reduces hematoma volume following collagenase-induced intracerebral hemorrhage in rats.*Nanomedicine* **2017**,13,1853-1862,doi:10.1016/j.nano.2017.04.001.

19. Cheng,Y.;Zhang,J.;Hou,W.;Wang,D.;Li,F.;Zhang,Y.;Yuan,F.Immunoregulatory effects of sinomenine on the T-bet/GATA-3 ratio and Th1/Th2 cytokine balance in the treatment of mesangial proliferative nephritis.*Int.Immunopharmacol.***2009**,9,894-899,doi:10.1016/j.intimp.2009.03.014.

20. Conti,E.;Tremolizzo,L.;Santarone,M.E.;Tironi,M.;Radice,I.;Zoia,C.P.;Aliprandi,A.;Salmaggi,A.;Dominici,R.;Casati,M.et al.Donepezil modulates the endogenous immune response:implications for Alzheimer's disease.*Hum.Psychopharmacol.-Clin.Exp.***2016**,31,296-303,doi:10.1002/hup.2538.

21. Dasgupta,S.;Roy,A.;Jana,M.;Hartley,D.M.;Pahan,K.Gemfibrozil ameliorates relapsing-remitting experimental autoimmune encephalomyelitis independent of peroxisome proliferator-activated receptor-alpha.*Mol.Pharmacol.***2007**,72,934-946,doi:10.1124/mol.106.033787.

22. Sulaiman,I.;Lim,J.C.;Soo,H.L.;Stanslas,J.Molecularly targeted therapies for asthma:Current

- development,challenges and potential clinical translation.*Pulm.Pharmacol.Ther.***2016**,40,52-68,doi:10.1016/j.pupt.2016.07.005.
23. Stein,C.;Hopfeld,J.;Lau,H.;Klein,J.Effects of Ginkgo biloba Extract EGb 761,Donepezil and their Combination on Central Cholinergic Function in Aged Rats.*J.Pharm.Pharm.Sci.***2015**,18,634-646,doi:10.18433/j3wc8v.
24. Qi,B.;Hu,L.;Zhu,L.;Shang,L.;Wang,X.;Liu,N.;Wen,N.;Hong,Y.;Fang,D.Metformin Attenuates Neurological Deficit after Intracerebral Hemorrhage by Inhibiting Apoptosis,Oxidative Stress and Neuroinflammation in Rats.*Neurochem.Res.***2017**,42,2912-2920,doi:10.1007/s11064-017-2322-9.
25. Shi,R.Y.;Zhu,S.H.;Li,V.;Gibson,S.B.;Xu,X.S.;Kong,J.M.BNIP3 interacting with LC3 triggers excessive mitophagy in delayed neuronal death in stroke.*CNS Neurosci.Ther.***2014**,20,1045-1055,doi:10.1111/cns.12325.
26. Tang,Y.C.;Tian,H.X.;Yi,T.;Chen,H.B.The critical roles of mitophagy in cerebral ischemia.*Protein Cell* **2016**,7,699-713,doi:10.1007/s13238-016-0307-0.
27. Kim-Han,J.S.;Kopp,S.J.;Dugan,L.L.;Diringer,M.N.Perihematomal mitochondrial dysfunction after intracerebral hemorrhage.*Stroke* **2006**,37,2457-2462,doi:10.1161/01.STR.0000240674.99945.4e.
28. Guan,R.;Zou,W.;Dai,X.;Yu,X.;Liu,H.;Chen,Q.;Teng,W.Mitophagy,a potential therapeutic target for stroke.*J.Biomed.Sci.***2018**,25,87,doi:10.1186/s12929-018-0487-4.
29. Zhang,X.;Yuan,Y.;Jiang,L.;Zhang,J.;Gao,J.;Shen,Z.;Zheng,Y.;Deng,T.;Yan,H.;Li,W.et al.Endoplasmic reticulum stress induced by tunicamycin and thapsigargin protects against transient ischemic brain injury:Involvement of PARK2-dependent mitophagy.*Autophagy*

2014,10,1801-1813,doi:10.4161/auto.32136.

30. Wu,M.F.;Wang,S.G.Human TAO kinase 1 induces apoptosis in SH-SY5Y cells.*Cell Biol.Int.***2008**,32,151-156,doi:10.1016/j.cellbi.2007.08.006.

31.

Zhang,X.;Yuan,D.;Sun,Q.;Xu,L.;Lee,E.;Lewis,A.J.;Zuckerbraun,B.S.;Rosengart,M.R.Calcium/calmodulin-dependent protein kinase regulates the PINK1/Parkin and DJ-1 pathways of mitophagy during sepsis.*Faseb J.***2017**,31,4382-4395,doi:10.1096/fj.201601096RRR.

32.

Jogdand,P.;Siddhuraj,P.;Mori,M.;Sanden,C.;Jonsson,J.;Walls,A.F.;Kearley,J.;Humbles,A.A.;Kolbeck,R.;Bjerner,L.et al.Eosinophils,basophils and type 2 immune microenvironments in COPD-affected lung tissue.*Eur.Resp.J.***2020**,55,doi:10.1183/13993003.00110-2019.

33. Ma,Y.;Li,J.;Wang,H.;Chiu,Y.;Kingsley,C.V.;Fry,D.;Delaney,S.N.;Wei,S.C.;Zhang,J.;Maitra,A.et al.Combination of PD-1 Inhibitor and OX40 Agonist Induces Tumor Rejection and Immune Memory in Mouse Models of Pancreatic Cancer.*Gastroenterology* **2020**,159,306-319,doi:10.1053/j.gastro.2020.03.018.

34. Takaku,M.;Grimm,S.A.;De Kumar,B.;Bennett,B.D.;Wade,P.A.Cancer-specific mutation of GATA3 disrupts the transcriptional regulatory network governed by Estrogen Receptor alpha,FOXA1 and GATA3.*Nucleic Acids Res.***2020**,48,4756-4768,doi:10.1093/nar/gkaa179.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryTableS1.docx](#)
- [SupplementaryFigureS2.docx](#)
- [SupplementaryFigureS1.docx](#)