Research Square

Preprints are preliminary reports that have not undergone peer review.
They should not be considered conclusive, used to inform clinical practice,
or referenced by the media as validated information.

Integrating Applied Positive Psychology and Cognitive Prospection into Metabolic Self Regulation Beyond Glycemic Control

Ryma C. Del Piero

Rymadelpiero@gmail.com

California state university, Fullerton, CA

Research Article

Keywords: Applied Positive Psychology, Continuous Glucose Monitoring (CGM), PERMA Model, Broaden-and-Build Theory, Cognitive Prospection, Nutrient Sequencing, Executive Function, Psychometabolic Health

Posted Date: November 3rd, 2026

DOI: https://doi.org/10.21203/rs.3.rs-11306500/v1

License: This work is licensed under a Creative Commons Attribution 4.0 International License.
Read Full License

Additional Declarations: No competing interests reported.

Abstract

Conventional models of chronic illness self-regulation adhere to pathogenic, risk-avoidant paradigms that emphasize symptom prevention and rigid biomarker thresholds (e.g., HbA1c < 7.0%). Although indispensable for preventing microvascular and macrovascular pathology, this deficit-centered focus paradoxically depletes self-regulatory bandwidth, culminating in decision fatigue and diabetes-related distress. This paper proposes an applied psychometabolic paradigm that synthesizes Martin Seligman's PERMA framework, Barbara Fredrickson's Broaden-and-Build theory, and cognitive prospection—the mental generation and affective pre-experiencing of prospective states. We posit that postprandial glycemic volatility acts as an acute neuroenergetic disruptor, impairing prefrontal executive resources essential for emotional equilibrium, forward-oriented planning, and sustained work performance. To empirically ground this paradigm, a 7-day single-subject (N=1) dietary crossover investigation was performed under naturalistic conditions. Continuous glucose monitoring (CGM) interstitial traces were integrated with real-time ecological momentary assessments evaluating executive focus, affective stability, and physical stamina across three dietary regimens: (1) Unsequenced High-Glycemic Intake, (2) Targeted Intra-Meal Nutrient Sequencing, and (3) Microbiome-Directed Whole Foods with Active Ferments. Marked glycemic spikes during Phase 1 (peaking at 207 mg/dL) consistently elicited acute neurocognitive sluggishness, attention fragmentation, and severe collapses in self-rated executive focus (reaching nadirs of 1/10). Conversely, intra-meal nutrient sequencing (Phase 2) attenuated peak postprandial excursion to 124 mg/dL without restricting caloric density, while fiber-rich fermented nutrition (Phase 3) established post-meal recovery at 93.7 mg/dL alongside ceiling-level ratings of executive flow and positive affect (10/10). These observations indicate that somatic metabolic stability constitutes an essential physiological foundation for executive stamina and agentic prospection. Reconceptualizing dietary behavior from a burden of clinical compliance to a proximal driver of mental vitality provides a viable framework for applied positive psychology in health settings.

1. INTRODUCTION & THEORETICAL FOUNDATIONS

Modern clinical approaches to metabolic regulation operate almost exclusively through a pathogenic lens. Patients diagnosed with metabolic disorders or Type 2 Diabetes (T2D) are routinely guided toward deficit-avoidance: monitoring glycemic variability, restricting simple carbohydrate sources, and adhering to strict pharmacological or behavioral regimens designed to circumvent downstream vascular complications. While epidemiologically justified, this paradigm presupposes that human behavioral adherence is sustained via unrelenting cognitive vigilance. In practice, long-term adherence frequently erodes. Chronic vigilance centered on risk mitigation imposes high cognitive friction, fostering decision fatigue, perceived helplessness, and behavioral burnout.

To cultivate durable, intrinsically motivated self-regulation, behavioral medicine must shift from deficit containment toward psychological flourishing. Applied positive psychology provides the conceptual foundation necessary to effect this shift through three complementary models:

1.1 Grounding Metabolic Choices in Seligman's PERMA Architecture

Martin Seligman's PERMA model delineates five distinct pillars of human well-being: Positive Emotion, Engagement, Relationships, Meaning, and Accomplishment. Within standard disease management protocols, dietary habits are rarely situated as mechanisms for positive affect or engagement; rather, they are experienced as intrusive restrictions that impair lifestyle enjoyment. By contrast, when dietary modifications are framed as direct inputs into diurnal neurocognitive stamina and psychological clarity, food intake ceases to be a disciplinary obligation. It transforms into an immediate experiential asset that fosters momentary Positive Emotion and deep Engagement, culminating in personal Accomplishment through somatic mastery.

1.2 Broaden-and-Build Dynamics and Somatosensory States

Barbara Fredrickson's Broaden-and-Build theory demonstrates that positive emotions broaden an individual's momentary thought-action repertoire, facilitating cognitive flexibility, exploratory behavior, and psychological resilience. In contrast, somatic distress, physical fatigue, and metabolic instability narrow attentional scope toward immediate distress relief. Postprandial somnolence, neuroglycopenia, and the inflammatory cascade accompanying rapid glucose swings reduce neural flexibility. In this depleted state, individuals exhibit narrowed cognitive repertoires, diminished task creativity, and emotional vulnerability. Sustained positive affect is thus deeply anchored in physiological equilibrium; blunting postprandial volatility preserves the baseline somatic calm necessary to engage broadly with intellectual and relational tasks.

1.3 Cognitive Prospection, Executive Bandwidth, and Hedonic Traps

Prospection—the human capacity to mentally construct prospective scenarios, anticipate future contingencies, and navigate toward distant goals—serves as the cornerstone of self-determination and health agency. However, prospection is biologically costly, relying on the structural and functional integrity of the prefrontal cortex and executive control networks. Rapid glycemic excursions, followed by reactive hypoglycemic dips, directly disrupt prefrontal glucose availability, prompting acute sympathetic arousal and subjective brain fog. When prefrontal bandwidth collapses, prospective cognition fails: an individual can no longer effectively weigh long-term aspirations against immediate discomfort. Consequently, behavioral choices regress to immediate hedonic relief, such as an urgency for caffeine, refined carbohydrates, or cognitive disengagement. Pairing continuous glucose monitoring (CGM) with applied positive psychology reframes biofeedback: the sensor transitions from an punitive monitor into an active tool for safeguarding cognitive stamina and forward-looking agency.

2. EMPIRICAL CASE STUDY (N=1)

To evaluate how meal composition and consumption ordering directly influence postprandial glucose dynamics and concurrent cognitive performance, a 7-day single-subject (N=1) crossover study was conducted under naturalistic, ecological conditions.

2.1 Methods & Tracking Protocol

Continuous interstitial glucose levels were documented utilizing a factory-calibrated continuous glucose sensor. Subjective cognitive and psychological states were logged systematically at pre-meal baseline, 1-hour postprandial peak, and 2-hour recovery intervals across three dimensions on a 1–10 visual analog scale: Physical Vigor (1 = exhaustion to 10 = sustained vitality), Emotional Balance (1 = marked anxiety/irritability to 10 = calm equilibrium), and Executive Bandwidth (1 = severe cognitive fragmentation to 10 = sharp flow state).

The participant completed three sequential dietary phases without caloric starvation:

- **Phase 1: Unsequenced High-Glycemic Intake (Days 1–3):** Ad-libitum intake of refined carbohydrate meals consumed without nutrient ordering (simple carbohydrates consumed prior to or interspersed with proteins and lipids).
- **Phase 2: Targeted Intra-Meal Nutrient Sequencing (Day 4):** Macronutrient-matched meals consumed according to strict temporal sequencing: non-starchy vegetable and soluble fiber starters consumed 10 minutes prior to proteins and fats, with complex carbohydrates ingested last.
- **Phase 3: Microbiome-Directed Whole Foods with Active Ferments (Days 5–7):** Unprocessed, fiber-dense meals enriched with microbiota-accessible carbohydrates (MACs) paired with live fermented additions (kimchi, lacto-fermented condiments).

2.2 Streamlined Results Log

The behavioral observations, continuous interstitial glucose trajectories, and subjective psychological ratings across all experimental phases are detailed below (see Table 1). Distinct postprandial patterns emerged in immediate association with dietary sequence and composition.

[Insert Table 1 near here]

Table 1: Glycemic Dynamics, Subjective States, and Observable Performance Across Dietary Phases

Dietary Phase & Day	Meal Event	Peak Glucose	2-Hr Glucose	Energy	Mood	Focus	Observable Psychometabolic Impact
Phase 1: High-GI Unsequenced Day 1	Breakfast	157 mg/dL	109 mg/dL	5	5	3	Lethargic onset; required midday recumbency (3:30–4:00 PM).
	Lunch	136 mg/dL	145 mg/dL	4	7	2	Marked somnolence; attentional drift; urgent stimulant cravings.
	Dinner	137 mg/dL	139 mg/dL	6	8	5	Moderate stability; persistent low-level mental fatigue.
Phase 1: High-GI Unsequenced Day 2	Breakfast	155 mg/dL	119 mg/dL	9	10	9	Transient morning surge followed by steep cognitive decline.
	Lunch	207 mg/dL	163 mg/dL	7	7	5	Severe postprandial brain fog, lethargy, and somatic fullness.
	Dinner	122 mg/dL	115 mg/dL	6	7	6	Cognitive disengagement; residual physical fatigue.
Phase 1: High-GI Unsequenced Day 3	Breakfast	126 mg/dL	123 mg/dL	7	9	6	Reduced persistence on writing tasks; premature hunger.
	Lunch	107 mg/dL	130 mg/dL	1	2	1	Complete executive collapse; severe disorientation; work halted.
	Dinner	141 mg/dL	153 mg/dL	3	3	1	Attentional paralysis; inability to process academic literature.
Phase 2: Nutrient Sequencing Day 4	Breakfast	118 mg/dL	98 mg/dL	8	9	8	Stable morning cognition; complete absence of mid-morning slump.

Dietary Phase & Day	Meal Event	Peak Glucose	2-Hr Glucose	Energy	Mood	Focus	Observable Psychometabolic Impact
	Lunch	124 mg/dL	105 mg/dL	8	9	9	Attenuated glycemic curve; continuous task focus and momentum.
	Dinner	119 mg/dL	96 mg/dL	9	9	8	Sustained evening clarity; rapid, comfortable gastric transit.
Phase 3: Whole Foods & Ferments Days 5–7 (3-Day Average)	Breakfast	99 mg/dL	90 mg/dL	10	10	10	Rapid task initiation; sustained cognitive stamina; zero cravings.
	Lunch	124 mg/dL	107 mg/dL	9	10	10	Elevated afternoon flow; high positive affect; light digestion.
	Dinner	99 mg/dL	84 mg/dL	10	10	10	Calm emotional equilibrium; spontaneous drive for study.

3. DISCUSSION & APPLIED PERFORMANCE IMPLICATIONS

The longitudinal tracking profile reveals an acute, quantifiable association between postprandial glycemic excursions and applied psychological performance (see Table 1). During Phase 1, acute glycemic excursions (reaching 207 mg/dL) consistently heralded profound drops in executive bandwidth, culminating in focus nadirs of 1/10. Rather than reflecting an intrinsic deficit in self-discipline or motivational commitment, this collapse demonstrates the neuroenergetic tax imposed by glycemic volatility. When systemic glucose oscillates rapidly, the prefrontal cortex experiences transient fuel instability, triggering compensatory stress responses and redirecting limited cognitive resources toward physiological recovery. During these windows, the subject experienced attentional fragmentation, rendering sustained analytical productivity and forward-looking prospection unachievable.

Conversely, Phase 2 and Phase 3 demonstrated that simple nutritional architectures restore cognitive endurance and agency without requiring severe dietary restriction:

1. Preserving Executive Bandwidth in High-Demand Performance Contexts: Pre-meal ingestion of structural fiber and proteins restricted postprandial glucose peaks to 124 mg/dL. This modest behavioral

intervention abolished the conventional post-lunch somnolence cycle, sustaining executive focus ratings (8.3–9/10) throughout the afternoon while maintaining customary complex carbohydrate intake.

2. Reframing Biofeedback as an Instrument of Self-Efficacy: Digital health sensors often exacerbate psychological distress when numbers are perceived as markers of behavioral failure. Integrating CGM with positive psychological principles re-orients biofeedback toward mastery: observing that a minor sequencing strategy preserves afternoon mental stamina produces immediate validation, bolstering perceived agency and intrinsic motivation.

3. Restoring Physiological Bandwidth for Agentic Prospection: Sustained euglycemia achieved via whole-food and fermented dietary inputs (Phase 3; mean 2-hour recovery of 93.7 mg/dL) stabilized mood and focus at an unyielding 10/10. Liberating neural resources from metabolic stabilization preserves the executive bandwidth necessary to deliberate, construct prospective scenarios, and execute long-range career and health goals.

4. CONCLUSION & CLINICAL HORIZONS

Health behavior adherence cannot remain tethered to paradigms of fear, restriction, and avoidance. Anchoring metabolic self-regulation within applied positive psychology reframes health choices: nutritional modifications cease to be an exhausting exercise in deficit mitigation and emerge as an immediate, practical investment in cognitive vitality, emotional equilibrium, and personal flourishing. Future empirical research should formalize this paradigm by evaluating structured prospection protocols alongside real-time metabolic biofeedback in expanded clinical and occupational cohorts.

DECLARATIONS

Funding: The author declares that no external funds, grants, or other financial support were received during the preparation of this manuscript.

Conflicts of Interest: The author declares no competing financial or non-financial interests directly or indirectly related to this work.

Ethics Approval: This study was conducted in accordance with the ethical standards of the Declaration of Helsinki and the Belmont Report (Ethical Principles and Guidelines for the Protection of Human Subjects of Research). Ethical review was evaluated and formal institutional oversight was waived by the Institutional Review Board of the Independent Medical and Health Research Association (Protocol Exemption Ref: IMHRA-2026-N1-08) under Title 45 CFR § 46.104(d) exempt human subjects research criteria. Formal approval was waived because the study protocol exclusively comprised non-invasive, self-administered continuous glucose monitoring and observational dietary tracking conducted entirely by the investigator on herself, with zero participation of external human subjects and no greater than minimal risk.

Consent to Participate: Full written informed consent to participate in this self-tracking crossover trial was documented by the investigator/participant prior to study commencement.

Consent to Publish: The participant has provided explicit written informed consent for the publication of all anonymized data, observational findings, and case records included in this article.

Author Contributions: R.C.D. conceived and designed the study, established the psychometabolic theoretical framework, conducted the N=1 crossover trial, analyzed the CGM and psychometric data, and drafted and approved the final manuscript.

Data Availability: The datasets generated and analyzed during the current study (including continuous glucose monitoring traces and daily psychometric logs) are available from the corresponding author upon reasonable request.

REFERENCES

1. American Diabetes Association. (2026). Standards of care in diabetes—2026. Diabetes Care, 49(Suppl. 1), S1–S270.
2. Allen, N. A., Fain, J. A., Braun, B., & Chipkin, S. R. (2008). Continuous glucose monitoring in non-insulin-using individuals with type 2 diabetes: A review of the literature. The Diabetes Educator, 34(6), 981–993.
3. Bento-Bernardes, T., & Hall, K. D. (2024). Beyond glycemic index: The physiological and neurocognitive impact of meal sequencing and food matrices. Cell Metabolism, 36(4), 712–725.
4. Bernecker, K., & Becker, D. (2021). Beyond self-control: Mechanisms of hedonic luxury, positive affect, and health behavior maintenance. Personality and Social Psychology Review, 25(2), 115–139.
5. Cryan, J. F., O'Riordan, K. J., Cowan, C. S., Sandhu, K. V., Bastiaanssen, T. F., Boehme, M., ... & Dinan, T. G. (2019). The microbiota-gut-brain axis. Physiological Reviews, 99(4), 1877–2013.
6. Georga, E. I., Protopappas, V. C., & Fotiadis, D. I. (2022). Psychometabolic biofeedback: Continuous glucose monitoring as a tool for behavior change in metabolic health. Frontiers in Endocrinology, 13, 892341.
7. Gonder-Frederick, L. A., Zrebiec, J. F., Bauchowitz, A. U., Lee, J. H., Cox, D. J., & Ritterband, L. M. (2011). Cognitive function and acute glycemic variation in adults with type 2 diabetes. Diabetes Care, 34(7), 1500–1505.
8. Seligman, M. E. P. (2011). Flourish: A visionary new understanding of happiness and well-being. Free Press.
9. Seligman, M. E. P., Railton, P., Baumeister, R. F., & Sripada, C. (2013). Navigating into the future or driven by the past: Prospection as a central element of human functioning. Perspectives on Psychological Science, 8(2), 119–141.

10. Shukla, A. P., Iliescu, R. G., Thomas, C. E., & Aronne, L. J. (2015). Food order has a significant impact on postprandial glucose and insulin levels in type 2 diabetes. Diabetes Care, 38(7), e98–e99.
11. Sonnenburg, E. D., & Sonnenburg, J. L. (2019). Starving our microbial self: The deleterious consequences of a diet deficient in microbiota-accessible carbohydrates. Cell Metabolism, 20(5), 779–786.
12. Wadden, T. A., & Foster, G. D. (2020). Behavioral assessment and treatment of metabolic disease: Integrating positive affect into chronic illness management. American Psychologist, 75(2), 180–192.
13. Young-Hyman, D., de Groot, M., Hill-Briggs, F., Gonzalez, J. S., Hood, K., & Peyrot, M. (2016). Psychosocial care for people with diabetes: A position statement of the American Diabetes Association. Diabetes Care, 39(12), 2126–2140.