

Evidence that the perinatal maternal microbiome mediates metabolite signaling programs that drive sex-specific microglial metabolic alterations and Alzheimer's disease susceptibility

Piyali Saha

University of Chicago

Nicholas Slimmon

The University of Texas Southwestern Medical Center

Ananya Nadgauda

University of Chicago

Priyam Patel

Northwestern University

Diya Mishra

University of Chicago

Pedro Blauth Poli

University of Chicago

Ana Inukai

University of Chicago

Ahtziri Fernandez

University of Chicago

Ian Weigle

University of Chicago

Xiaoqiong Zhang

University of Chicago

Sangram S. Sisodia

ssisodia@bsd.uchicago.edu

University of Chicago

Research Article

Keywords:

Posted Date: October 23rd, 2026

DOI: <https://doi.org/10.21203/rs.3.rs-11307065/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Additional Declarations: No competing interests reported.

Abstract

Maternal microbiome-dependent programming of neuroimmune and metabolic pathways during early development may critically shape sex-specific susceptibility to Alzheimer's disease (AD) later in life. We have documented that in mouse models of A β deposition, broad-spectrum antibiotic (ABX)-mediated perturbations of the gut microbiome postnatally (P14-P21) selectively reduces A β pathology only in male, but not in female animals. Here, we shift the developmental window upstream to test whether perinatal maternal microbiome perturbations shapes the pathological trajectories of offspring. Dams were treated with the broad-spectrum ABX from preconception through gestation, without direct exposure of the offspring. Strikingly, maternal microbiome disruption selectively reduced cortical A β burden in female offspring, with no effect in males, effectively reversing the sex bias observed following postnatal ABX manipulations. To explore the mechanisms underlying this effect, we now report that female offspring exhibited coordinated shifts in microbial composition and amino acid metabolism, that are coupled to cortical transcriptional programs enriched for GPCR signaling and Rho GTPase pathways. As many microbial metabolites and amino acid-derived ligands signal through GPCRs that are expressed by microglia, these transcriptional changes suggest enhanced GPCR-mediated nutrient sensing in microglia of female animals. These systems level changes in microglia drive a metabolically engaged amyloid clearing phenotype, providing a mechanistic link between maternal microbiome disruption and enhanced A β clearance in females. Together, these findings demonstrate that maternal microbial inheritance and developmental timing critically shape sex specific AD susceptibility and position temporal microbiome modulation as a potential precision strategy for disease prevention.

Introduction

Alzheimer's disease (AD) is characterized by the deposition of amyloid β (A β) peptides in extracellular plaques and the accumulation of hyperphosphorylated tau as intracellular neurofibrillary tangles within neurons. These pathological lesions lead to synaptic dysfunction, progressive neurodegeneration, neuronal loss, and ultimately cognitive decline and dementia (Price and Sisodia, 1998). Despite decades of research, there are few effective disease-modifying treatments that can prevent or halt disease progression, and current anti-A β therapies that are expensive provide only modest, temporary benefits and have significant side effects (Parums, 2024). Recent research has uncovered a pivotal role of the gut microbiome in modulating brain physiology through the gut-brain axis (Cryan et al., 2019). Dysbiosis has been linked to neurodevelopmental and neurodegenerative disorders, including AD, through immune, metabolic, and endocrine pathways (Bicknell et al., 2023). We have shown that postnatal ABX-induced microbiome dysbiosis reduces A β pathology and microglial activation in mouse models of A β amyloidosis that occurs specifically in male, but not female animals (Dodiya et al., 2019; Minter et al., 2016). To these efforts, we have demonstrated that estradiol-mediated elevation of amyloid pathology at later ages is, at least in part, responsible for lack of an effect of ABX on females (Saha et al., 2024). These findings underscore the temporal specificity of gut microbiome perturbation as a confounding factor in sex-dependent microbial-neural interactions. In this study, we therefore sought to investigate

how perinatal maternal microbiome perturbations during early development influences microbiome composition, cortical transcriptomes and amyloid pathology in the offspring.

The maternal microbiome plays a foundational role in shaping fetal immune and metabolic programming (Koren et al., 2024). Microbial metabolites and immune mediators cross the placental barrier, influencing neurodevelopmental pathways and establishing life-long trajectories of brain function (Di Gesu et al., 2021; Edwards et al., 2017; Vuong et al., 2020). Perturbations in maternal microbiota during pregnancy have been implicated in disorders with pronounced sex bias, including autism spectrum disorders (ASD) and metabolic syndromes (Biagioli et al., 2025; Salia et al., 2025), suggesting that maternal microbial inheritance may confer differential risk across sexes.

In this study, we hypothesized that disruption of the maternal gut microbiome could alter the microbiome of offspring and by so doing, reprogram the metabolic landscape of microglia and other CNS cell types that could lead to alterations in amyloid pathology. Using the APPPS1-21 transgenic mouse model of A β amyloidosis, we treated dams with a broad-spectrum ABX cocktail from preconception through gestation to induce maternal microbiome dysbiosis without directly altering the postnatal microbiota of the offspring. This design allowed us to isolate prenatal microbial effects on fetal brain and immune development. We refer to this approach as maternal microbiome perturbation because the intervention specifically targets the gut microbiota of the dams during the critical window of conception and pregnancy, thereby influencing fetal development largely through maternal immune, hormonal, and metabolic pathways rather than directly altering the postnatal microbiome of the offspring. In this model, dams were separated on the day of copulation, ensuring that the altered maternal microbiome exerted the predominant influence on fetal development and subsequent disease trajectory.

We implemented an integrative multi-omics approach combining 16S rRNA and shotgun metagenomic sequencing, metabolomics, and cortical transcriptomics, complemented by immunohistochemical analyses of A β pathology and microglial phenotypes. Our results reveal that maternal microbiome perturbation induces sex-specific microbial and metabolic remodeling leading to alterations in amino acid metabolism and activation of GPCR-linked nutrient-sensing pathways in female offspring. Our findings establish a mechanistic framework in which maternal microbial perturbation programs female-specific neuroprotection through metabolic reorganization and cortical GPCR activation. This work underscores the developmental timing and sex specificity of microbiome-brain interactions and highlights maternal microbial inheritance as a key determinant of AD vulnerability.

Results

16s rRNA sequencing reveals maternal gut microbiome perturbation leads to changes in microbial diversity in offspring

A schematic representation of the ABX treatment and experimental timeline used to evaluate the impact of maternal microbiome perturbations is shown in Fig. 1A. Analysis of cecum weight relative to body

weight shows a significant increase in male (MM(ABX)_Male) and female (MM(ABX)_Female) pups born to ABX exposed dams compared with male (MM(Ctrl)_Male) or female (MM(Ctrl)_Female) offspring from water-treated dams (Fig. 1S, left panel). Morphological examination further reveals that the ceca from all ABX-treated pups are enlarged and distended as a result of localized inflammation and fluid retention (Fig. 1B, right panel). Taxonomic profiling of cecal bacteria at the phylum level demonstrates a pronounced shift in microbial composition across treatment groups (Fig. 1B, a). Interestingly, the microbial communities of male and female pups born to ABX-treated dams exhibit clear sex-specific patterns, with male and female offspring forming distinct clusters. Alpha diversity analysis using the Chao1, Inverse Simpson, and Shannon indices reveals changes in microbial richness and evenness in all offspring from ABX-treated mice compared with offspring from vehicle-treated dams (Fig. 1B, b). Offspring from ABX-treated dams also show a reduction in microbial diversity compared with controls, although this reduction is less pronounced than that observed in cecal contents of dams that were treated with ABX. Statistical comparisons indicate a significant difference for the Chao1 index ($p < 0.05$), whereas the Inverse Simpson and Shannon indices do not reach statistical significance. The Chao1 index estimates species richness by approximating the number of distinct taxa present, the Inverse Simpson index emphasizes community dominance and evenness, and the Shannon index integrates both richness and evenness. Together, these metrics suggest that maternal ABX exposure alters microbial diversity and community balance, with sex-dependent differences in the magnitude and pattern of these changes.

Principal coordinates analysis (PCoA) of beta diversity, based on both Unweighted UniFrac and Bray–Curtis dissimilarity metrics, further reveals distinct clustering patterns (Fig. 1B, c). PERMANOVA analysis confirms that the overall microbiome composition differs significantly between offspring from ABX-treated and vehicle-treated dams. Notably, male and female offspring segregate into separate clusters, indicating sex-dependent microbial restructuring following maternal microbiome perturbation. Collectively, these findings demonstrate that maternal ABX exposure differentially shapes the gut microbial landscape of male and female progeny rather than exerting an equivalent effect across sexes.

MaAsLin2 analysis revealed pronounced sex-dependent microbial responses to maternal ABX exposure (Fig. 1B,e). (MM(ABX)_Female) animals exhibited a broader and stronger enrichment of several microbial taxa compared with (MM(ABX)_Male) animals, indicating enhanced colonization and restructuring of the gut microbiome. At the family level, *Bacteroidaceae*, *Akkermansiaceae*, *Enterococcaceae*, *Porphyromonadaceae*, *Rikenellaceae*, and *Sutterellaceae* showed stronger positive associations in (MM(ABX)_Female) pups compared with (MM(ABX)_Male) pups. Similarly, several genera displayed markedly higher enrichment in (MM(ABX)_Female) pups, including *Blautia*, *Phocaeicola*, *Akkermansia*, *Enterococcus*, *Bacteroides*, *Prevotella*, *Parasutterella*, *Dubosiella*, and *Faecalicoccus*, suggesting an expansion of Bacteroidota- and Verrucomicrobia-associated lineages. In contrast, (MM(ABX)_Male) pups exhibited fewer positively associated taxa and retained a greater proportion of genera typically enriched in (MM(Ctrl)_Male) pups, including *Lactobacillus*, *Bifidobacterium*, *Muribaculum*, *Ruminococcus*, and *Clostridium XIVa*, indicating reduced microbial recolonization following maternal microbiome perturbation. Collectively, these findings demonstrate that maternal ABX exposure induces a

more pronounced restructuring and enrichment of specific microbial lineages in female offspring, whereas male offspring display comparatively limited microbiome reconstitution.

Collectively, these findings indicate that female offspring exhibit a stronger microbial response to maternal ABX-induced perturbation, characterized by broader enrichment of multiple taxa and greater restructuring of the gut microbial community. In contrast, male offspring display a comparatively limited taxonomic shift and retain a larger proportion of taxa associated with offspring from vehicle-treated dams, consistent with reduced recolonization and a more constrained microbiome response following maternal ABX exposure.

Gut Microbiota Perturbation of Dams Reduces A β Burden in Female Offspring

Immunofluorescence analysis of cortical A β plaque deposition using A β -specific 3D6 antibody (Fig. 2A) revealed sex-specific effects of gut microbiota perturbations. While A β burden remained comparable between (MM(Ctrl)_Male) and (MM(ABX)_Male) animals at the time of sacrifice, a statistically significant reduction in A β plaques was observed in (MM(ABX)_Female) relative to their control counterparts from vehicle-treated dams ($p < 0.05$; Fig. 2B). These findings suggest that maternal microbiome alterations selectively modulate A β accumulation in a sex-dependent manner, with a pronounced effect in females. Western blot analysis of the transgene-encoded full-length amyloid precursor protein 751 isoform (APP-FL) in cortical extracts (Fig. 2C) demonstrated no significant differences in APP levels between pups from vehicle- and ABX-treated dams of either sex. Densitometric quantification across both male and female groups revealed that the observed reduction in A β burden in (MM(ABX)_Female) pups relative to (MM(Ctrl)_Female) pups is not attributable to altered steady-state levels of APP. We further examined the steady-state levels of BACE1 and PS1, two key enzymatic components of the amyloidogenic APP-processing pathway. Western blot analysis showed no significant differences in levels of BACE1 or PS1 (PS1-NT is the amino-terminal derivative generated by autocatalytic processing of full-length PS1) between control and maternal ABX groups in either male or female offspring (Fig. 2D–E). Together, these findings indicate that the female-specific reduction in A β pathology following maternal ABX treatment occurs without detectable changes in APP abundance or in the protein levels of these major components of the amyloidogenic APP-processing pathway.

Metagenomic profiling reveals sex-specific alterations in gut microbiota of offspring

The taxonomic composition of the gut microbiota in offspring varied markedly across sex and ABX treatment (Fig. 3A). In (MM(Ctrl)_Male) pups, the microbiota was dominated by *Bacteroides acidifaciens*, *Bacteroides caecimuris*, and *Candidatus Arthromitus* sp. SFB_mouse. In (MM(ABX)_Male), we observed a drastic reduction of these taxa, accompanied by increases in *Blautia pseudococcoides*, *Burkholderiaceae bacterium*, and *Phocaeicola sartorii*. In comparison, (MM(Ctrl)_Female) exhibited a more diverse

microbiota, characterized by higher relative abundances of *Bacteroidales bacterium*, *Lactobacillus johnsonii*, *Muribaculaceae bacterium isolate 104 HZI*, and *Lactobacillus reuteri*. Moreover, in (MM(ABX)_Female) pups, we observed a marked loss of these latter taxa, with concomitant increases in *Xylanibacter rodentium*, *Parabacteroides distasonis*, and *Phocaeicola sartorii*, indicating a pronounced ABX-driven restructuring of the gut community. Several strains exhibited baseline sex differences in (MM(Ctrl)_Male) and (MM(Ctrl)_Female) pups (Fig. 3C), including higher *Bacteroidales bacterium* and *Adlercreutzia sp. DSM 108611* in (MM(Ctrl)_Female) pups and *Adlercreutzia sp. DSM 109821* in (MM(Ctrl)_Male) pups. Both *Adlercreutzia* species were nearly depleted in (MM(ABX)_Female) pups. Notably, *L. johnsonii* and *L. reuteri*, species that are linked to methionine-associated metabolic pathways critical for one-carbon metabolism and methyl-donor balance, were significantly more abundant in (MM(Ctrl)_Female) pups than in (MM(ABX)_Female) pups ($p < 0.005$), and both species were either low or undetectable in males, underscoring their female-specific depletion following maternal microbiome perturbation. A subset of taxa, including *Akkermansia muciniphila*, *Dubosiella newyorkensis*, *Blautia pseudococcoides*, *Burkholderiaceae bacterium*, *Parabacteroides distasonis*, and *Phocaeicola sartorii*, increased significantly in both sexes after ABX exposure ($p < 0.05$ to $p < 0.0001$), suggesting sex-independent recolonization. Conversely, *Bifidobacterium pseudolongum*, *Bacterium_1XD21_13*, and *Clostridia bacterium* declined significantly in both sexes ($p < 0.05$ to $p < 0.0001$). ABX-induced maternal microbial perturbation also produced more extensive compositional changes in (MM(ABX)_Female) pups than in (MM(ABX)_Male) pups (Fig. 3A). Specifically, *Muribaculum intestinale*, *Clostridium innocuum*, *Clostridium cocleatum*, *Desulfovibrionaceae bacterium*, *Rodentibacter heilii*, and *Bacteroides muris* were markedly reduced in (MM(ABX)_Female) pups ($p < 0.05$ to $p < 0.001$), while *Xylanibacter rodentium* and *Staphylococcus xylosus* showed female-specific increases ($p < 0.05$ to $p < 0.001$). Male-specific differences included elevated *Bacteroides acidifaciens* and *Bacteroides caecimuris* in (MM(Ctrl)_Male) ($p < 0.0001$) and reduced *Turicimonas muris* and *Candidatus Arthromitus* in (MM(ABX)_Male) ($p < 0.05$ to $p < 0.0001$).

Overall, the magnitude and diversity of microbial alterations were greater in females, highlighting their heightened susceptibility of female pups to maternal ABX exposure. Some non-significant trends, including reduced abundance of *Bacterium_1XD42_1*, *Alistipes sp. DSM 112343*, *Alistipes indistinctus*, *Coriobacteriaceae bacterium*, and *Muribaculaceae* isolates, further support sex- and ABX-dependent remodeling of the gut microbial ecosystem.

To further evaluate similarities in microbial restructuring across groups, we performed linear regression-based correlation analyses comparing species-level abundances (Fig. 3B). A very weak but statistically significant positive correlation was observed between MM(ABX)_Female and MM(ABX)_Male microbiomes ($R^2 = 0.1208$, $p = 0.0208$), indicating limited overlap in taxa that recolonize following maternal ABX exposure. Together, these analyses indicate that maternal ABX exposure drives a unique restructuring of the offspring gut microbiota, with females exhibiting greater taxonomic divergence and more extensive microbial remodeling.

Metabolomic profiling reveals sex-specific changes in metabolite composition

ABX-mediated maternal microbiome perturbation resulted in significant metabolic alterations, particularly in amino acid (Fig. 4A and 4S), short-chain fatty acid (SCFA) (Fig. 4S), and bile acid (Fig. 4S) metabolism. While many metabolite levels changed in male or female pups from ABX-treated dams compared to pups from vehicle-treated dams, the most significant alterations were observed in the amino acid panel (Fig. 4A), where there appears to be a sex-dependent effect on microbial fermentation.

Amino acid profiling revealed that (MM(Ctrl)_Female) animals showed higher overall amino acid abundance under control conditions, which markedly decreased in (MM(ABX)_Female) pups, while (MM(ABX)_Male) animals displayed comparatively minor changes compared with (MM(Ctrl)_Male) animals (Fig. 4S). For instance, alanine was significantly reduced in (MM(ABX)_Female) animals compared with (MM(Ctrl)_Female) animals ($p < 0.01$) but remained unchanged between (MM(Ctrl)_Male) and (MM(ABX)_Male) animals. Interestingly, (MM(Ctrl)_Female) animals exhibited a significantly higher alanine concentration than (MM(Ctrl)_Male) animals ($p < 0.05$). Similarly, essential amino acids, including leucine, lysine, methionine, and phenylalanine, as well as branched-chain amino acids (BCAAs) including leucine and isoleucine, followed the same pattern, with a significant decrease in (MM(Ctrl)_Female) but no change in (MM(ABX)_Male) animals. Other amino acids, including pantothenic acid, tryptophol, and niacin, also showed a similar trend.

Cysteine and tyrosine showed a significantly higher increase in (MM(Ctrl)_Female) animals compared with (MM(Ctrl)_Male) animals ($p < 0.01$). Aspartate, PreQ1, and tryptamine exhibited a significant decrease in (MM(ABX)_Female) animals ($p < 0.05$) compared with (MM(Ctrl)_Female) animals, with no significant change in males from vehicle or ABX-treated dams. Additionally, dopamine and quinine levels were significantly increased in (MM(ABX)_Female) animals ($p < 0.01$), with no detectable differences between males from vehicle or ABX-treated dams.

Functional metagenomic–metabolomic integration (Fig. 4B) revealed strong sex-specific correlations between microbial pathways and amino acid metabolites. In (MM(Ctrl)_Female) animals, *Lactobacillus johnsonii* and *Lactobacillus reuteri* were enriched and strongly associated with methionine-related routes, including the S-adenosyl-L-methionine (SAM) salvage I (PWY-6151; $p = 0.00053$) and the L-lysine, L-threonine, and L-methionine biosynthesis II (PWY-724; $p = 0.0021$) pathways. These associations were reduced in (MM(ABX)_Female) animals, indicating disruption of methionine metabolism, methyl-donor balance, and one-carbon cycling. Together, these findings indicate that ABX-mediated maternal microbiome perturbations induces sex-specific microbial and metabolic reorganization, wherein female offspring exhibit amino acid restriction through reduced microbial methionine metabolism and host one-carbon regulation.

Transcriptomic analysis reveals sex-specific transcriptional responses in offspring from ABX-exposed dams

We extracted RNA from the dorsal cerebral cortex of male and female APPPS1-21 transgenic mouse offspring born to dams treated with either vehicle or ABX, and performed RNA sequencing to evaluate cortical transcriptional responses to maternal microbiome perturbations. Principal component analysis (PCA) revealed clear separation of samples by sex along the first two principal components (PC1: 33% variance; PC2: 28% variance), with male and female samples forming distinct clusters (Fig. 5A,a). Within each sex, maternal ABX exposure further shifted the transcriptional profile, with RNA reads from (MM(ABX)_Female) samples forming a distinct cluster relative to RNA reads from (MM(Ctrl)_Female) samples. Consistent with this pattern, a chord diagram illustrating transcriptomic relationships among groups revealed stronger divergence between (MM(Ctrl)_Female) and (MM(ABX)_female) samples compared with (MM(Ctrl)_Male) versus (MM(ABX)_Male) samples, indicating a larger transcriptional shift in the female cortex following maternal microbiome perturbation (Fig. 5A,b).

Differential expression analysis demonstrated a markedly stronger transcriptional response in females compared with males. (MM(ABX)_Female) exhibited 372 differentially expressed genes (DEGs) compared with (MM(Ctrl)_Female), whereas (MM(ABX)_Male) showed only 81 DEGs compared with (MM(Ctrl)_Male) (FDR-adjusted $p < 0.05$) (Fig. 5A,c). Only six genes overlapped between the two comparisons, indicating that the transcriptional response to maternal microbiome perturbation is largely sex-specific.

Hierarchical clustering analysis further illustrated these differences (Fig. 5A,d). RNA from (MM(ABX)-Female) formed a distinct transcriptional cluster that separated clearly from RNA from (MM(Ctrl)_Female), reflecting coordinated gene expression changes across multiple gene modules. In contrast, RNA from (MM(ABX)_Male) showed comparatively smaller transcriptional shifts and clustered closer to transcripts from (MM(Ctrl)_Male), indicating a more limited cortical response to maternal ABX exposure.

Pathway enrichment analysis highlighted GPCR downstream signaling and the Rho GTPase cycle as the most prominent pathways altered in female offspring following maternal ABX exposure (Fig. 5B). These pathways are central regulators of intracellular signaling and cytoskeletal remodeling. Additional enriched categories included regulation of neuron projection development, plasma membrane–bounded cell projection morphogenesis, and neurotrophin signaling, indicating coordinated alterations in neuronal structure and connectivity. Multiple synaptic processes were also affected, including modulation of chemical synaptic transmission, presynaptic modulation of synaptic transmission, regulation of vesicle fusion, and regulation of neurotransmitter levels. Behavioral and systems-level processes, such as behavior and locomotory behavior, were also represented among the enriched categories. In contrast, pathway enrichment in males was substantially more limited, consistent with the smaller number of DEGs observed in the male cortex.

Pathway enrichment analysis comparing female ABX-treated offspring to male ABX-treated offspring revealed a distinct set of sex-biased biological processes (Fig. 5B,c). The most significantly enriched pathway was histone demethylation (GO:0016577), indicating sex-specific epigenetic regulation in response to maternal microbiome disruption. Additional enrichment of positive regulation of translation (GO:0045727) suggests differential control of protein synthesis between sexes under ABX conditions. Immune-related processes, including antigen processing and presentation (GO:0019882), were also overrepresented, pointing to heightened or altered immune signaling in females relative to males. GPCR-related signaling pathways, including class A rhodopsin-like GPCRs (WP189), were enriched, consistent with sex-dependent modulation of receptor-mediated signaling. Developmental and transport-related pathways, such as embryonic organ development (GO:0048568), organic anion transport (GO:0015711), and cellular response to abiotic stimulus (GO:0071214), further distinguished female from male ABX-treated cortices. Together, these findings indicate that maternal ABX exposure induces sex-specific epigenetic, translational, immune, and GPCR-mediated pathway alterations, with females exhibiting a broader and more coordinated transcriptional response than males under identical microbiome perturbation conditions.

Microglial Remodeling Around Amyloid Plaques in Response to Maternal Microbiome Perturbation

To assess how maternal microbiome perturbation influences A β -associated neuroinflammation, Iba1 immunofluorescence staining, three-dimensional (3D) reconstruction, and quantitative morphological analyses were performed in cortical sections from female offspring (Fig. 6A). Confocal z-stacks were reconstructed in 3D using Imaris to visualize individual Iba1⁺ microglia and their spatial relationship with A β plaques (Fig. 6B). In (MM(Ctrl)_Female) mice, large, dense A β plaques were surrounded by compact clusters of Iba1⁺ microglia exhibiting enlarged somata, shorter processes, and reduced branching, consistent with a less ramified plaque-associated morphology. In contrast, (MM(ABX)_Female) mice displayed smaller A β plaques surrounded by microglia with smaller somata, longer processes, and greater process branching, reflecting a more highly ramified morphology. These differences were also evident in the 3D reconstructions of microglia surrounding A β plaques.

Quantification revealed a significant reduction in the number of Iba1⁺ microglia in close proximity to A β plaques in (MM(ABX)_Female) compared with (MM(Ctrl)_Female) mice. Quantitative analysis of individual Iba1⁺ microglia further confirmed these morphological differences, demonstrating a significant decrease in soma area (Fig. 6a) and significant increases in total process length (Fig. 6b), branch points (Fig. 6c), junctions (Fig. 6c), and terminal endpoints (Fig. 6d) in (MM(ABX)_Female) compared with (MM(Ctrl)_Female) mice. Collectively, these findings demonstrate that the reduction in A β pathology following maternal microbiome perturbation is accompanied by reduced microglial accumulation around plaques and a shift toward a more highly branched and ramified microglial morphology.

To further characterize microglial phenotypes within the plaque microenvironment, we examined Clec7a, a marker associated with the microglial neurodegenerative/disease-associated (MGnD/DAM) state, and P2RY12, a marker associated with homeostatic microglia (Butovsky and Weiner, 2018; Paolicelli et al., 2022). Compared with (MM(Ctrl)_Female) mice, (MM(ABX)_Female) mice exhibited a significant reduction in Clec7a⁺ microglia and a concomitant increase in P2RY12⁺ microglia surrounding A β plaques (Fig. 6C). These molecular changes, together with the decreased soma area and increased process length, branch points, junctions, and terminal endpoints observed in the morphological analyses, support a shift away from a disease-associated microglial phenotype toward a more homeostatic and highly ramified microglial state following maternal microbiome perturbation. Importantly, this altered microglial phenotype was associated with the reduced A β burden observed in (MM(ABX)_Female) mice.

Discussion

In a series of preceding efforts, we demonstrated that postnatal ABX-induced microbiome depletion attenuates A β deposition and microglial activation in male, but not female, mouse models of A β amyloidosis (Dodiya et al., 2019; Minter et al., 2017; Saha et al., 2024). In view of the established role of the maternal microbiome in shaping fetal immune and metabolic programs (Koren et al., 2024), we have examined the impact of ABX-mediated maternal microbiome perturbations on the microbiome composition of male and female offspring and the effects on metabolic processes that might be associated with alterations in A β deposition. Our studies offer several important insights into the role of the maternal microbiome in regulation of cellular and physiological processes that mediate AD-type pathology and neuroinflammation in a mouse model. First, we demonstrate that maternal ABX exposure induces a more pronounced restructuring and enrichment of specific microbial lineages in female offspring, whereas male offspring display comparatively limited microbiome reconstitution. Second, we show that while A β burden remained comparable between male animals from vehicle-treated or ABX-treated dams, we observe a statistically significant reduction in A β plaques in female offspring from ABX-treated dams compared with their counterparts from vehicle-treated dams. These findings suggest that maternal microbiome alterations selectively modulate A β accumulation in a sex-dependent manner, with a pronounced effect in females. Third, metabolomic profiling of cecal contents from offspring of ABX-treated dams revealed sex-specific changes in metabolite composition, particularly in amino acids, short-chain fatty acids (SCFA), and bile acids (Fig. 4S) compared with the levels in ceca from offspring of vehicle-treated dams. Functional metagenomic–metabolomic integration revealed strong sex-specific correlations between selected microbes and amino acid metabolites. In female offspring from vehicle-treated dams, *Lactobacillus johnsonii* and *Lactobacillus reuteri* were enriched and strongly associated with methionine-related metabolism, including the S-adenosyl-L-methionine (SAM) salvage I (PWY-6151; $p = 0.00053$) and the L-lysine, L-threonine, and L-methionine biosynthesis II (PWY-724; $p = 0.0021$) pathways. These associations were reduced in female offspring from ABX-treated dams indicating disruption of methionine metabolism, methyl-donor balance, and one-carbon cycling, pathways that regulate epigenetic remodeling and redox homeostasis (Xing and Tu, 2025; Yan et al., 2025). As epigenetic changes are attributed to methylation of lysine residues on histones, and SAM is a methyl

donor, pathway enrichment analysis comparing female and male offspring from ABX-treated dams showed a significant enrichment of histone demethylation (GO:0016577) in females. In this regard, methionine restriction (MR), previously been shown to enhance metabolic flexibility and longevity (Kosakamoto et al., 2023; Latimer et al., 2018; Yan et al., 2025; Zhang et al., 2024), also leads to reduced A β accumulation in APP/PS1 mice (Xi et al., 2023).

Fourth, our transcriptomic studies of cortical RNAs in male and female offspring from vehicle- or ABX-treated dams and now observe a markedly stronger transcriptional response in females compared with males. Pathway enrichment analysis highlighted GPCR downstream signaling and the Rho GTPase cycle as the most prominent pathways altered in female offspring following maternal ABX exposure. While these pathways are central regulators of intracellular signaling and cytoskeletal remodeling, the impact on specific cortical cell types is uncertain. Finally, *Ms4a3*, a gene associated with peripheral myeloid lineages, was downregulated in the cortex of female offspring from ABX-treated dams, that is coincident with reduced amyloid burden. Because *Ms4a3* expression is most commonly associated with chronic inflammatory states, vascular dysfunction, and heightened peripheral immune signaling (Mattiola et al., 2021), its reduction in female offspring from ABX-treated dams might reflect attenuation of myeloid transcriptional signatures associated with amyloid reduction. In this altered inflammatory context, morphological analyses using Iba1 staining revealed that microglia surrounding A β plaques in female offspring born to microbiome-perturbed dams displayed elongated, highly ramified dendritic processes, consistent with a homeostatic state. Consistent with these effects, reduced amyloid burden in offspring from ABX-treated dams was associated with a decreased abundance of Clec7a⁺ degenerative microglia (MGnD/DAM) and a concomitant increase in P2ry12⁺ homeostatic microglia within the immediate plaque microenvironment. Collectively, these findings support a model in which maternal microbiome perturbation induces immune transcriptional reprogramming that biases microglia toward a metabolite-responsive, GPCR-driven activation state that favors morphological plasticity and efficient A β clearance while limiting degenerative microglial phenotypes. This GPCR-linked microglial activation state aligns with the paradigm that microbial metabolites crossing the blood–brain barrier directly modulate microglial and clearance capacity through ligand–receptor signaling (Erny et al., 2015; Young and Samuel, 2025). We propose that maternal microbiome perturbation primes microglia in females toward a GPCR-driven, metabolite-responsive phenotype that facilitates A β clearance and maintains neuroimmune equilibrium—analogueous to metabolic reprogramming observed under MR conditions (Kosakamoto et al., 2023; Latimer et al., 2018; Yan et al., 2025; Zhang et al., 2024).

It should be noted GO terms in our analyses (Fig. 5B-a) such as presynaptic modulation of neurotransmission (GO:0099171), inhibitory postsynaptic regulation (GO:0098828), neurotransmitter homeostasis (GO:0001505), vesicle fusion (GO:0031338), neurotrophin signaling (GO:0038179), and neuronal projection development (GO:0010975) indicate that the enriched pathways are predominantly associated with neuronal physiology and synaptic signaling. This is not unexpected or surprising as developmental processes and organization in the offspring are still ongoing during the period of maternal microbiome dysbiosis and after birth. Indeed, the volcano plots (Fig. 5A-e) clearly show an elevation in NT3 (neurotrophin 3), a neurotrophic factor secreted by astrocytes that is known to regulate

neuronal survival, differentiation, and synaptic function in both the peripheral and central nervous systems (BARBACID, 1995; Ernfors et al., 1994) and synaptotagmin (Syt17), a multifunctional regulator of intracellular protein trafficking in excitatory hippocampal neurons (Ruhl et al., 2019). We also observe a significant elevation in Kallikrein-related peptidase 12 (KLK12), a member of the serine protease family that is expressed by astrocytes in the human brain. A homologue of KLK12, termed KLK7, is a known A β degrading enzyme (Kidana et al., 2018) and in view of the sequence and structural similarity of the KLKs, it is not inconceivable that KLK12 also shares A β -degradative activity, that would explain the reduction in A β burden in female offspring of ABX-treated dams. Whether KLK12 is a *bona fide* A β degrading enzyme is uncertain but knockout and overexpression studies *in vitro* and *in vivo* would be critical to support, or refute this proposal. We also examined whether the reduction in A β pathology could be associated with alterations in the amyloidogenic APP-processing pathway. However, protein levels of APP-FL, BACE1, and PS1-NT were unchanged following maternal ABX treatment in both male and female offspring. These findings suggest that the female-specific reduction in A β pathology is unlikely to result from altered APP abundance or processing through the BACE1/PS1-dependent pathway and instead support the possibility that mechanisms affecting A β clearance or degradation may contribute to the observed phenotype.

The concept that maternal microbial inheritance imprints long-term neuroimmune trajectories is supported by the seminal work of Garel, Thion and colleagues (Thion et al., 2018), which demonstrated that the maternal microbiota regulates fetal microglial maturation at transcriptomic and chromatin-accessibility levels. Their study revealed sex- and time-specific windows of vulnerability to maternal microbiome absence—males being more affected in utero, while females exhibit greater alterations in adulthood. Multiple studies now demonstrate that female offspring are consistently more vulnerable to maternal perturbations involving the gut microbiome and diet. Maternal diet–microbiome disturbances drive epigenetic and developmental alterations that are more pronounced in females, highlighting their heightened sensitivity to early microbial and nutritional cues (Keleher et al., 2018; Masuyama et al., 2015). Recent work further shows that disruptions to the maternal microbiome produce sex-dependent alterations in brain development of offspring, with female offspring exhibiting more pronounced structural changes in the prefrontal cortex and distinct shifts in microbial colonization, underscoring a clear female-biased vulnerability (Gohmann et al., 2025). In our previous studies, direct ABX treatment of offspring during P14–P21 produced sex-dependent effects on AD pathology (Dodiya et al., 2018, 2021; Saha et al., 2024), whereas the present study uses maternal ABX treatment and therefore represents a substantially different developmental exposure. Consistent with this concept, Lynch et al. (2023) demonstrated that treatment with the same broad-spectrum ABX cocktail during closely spaced postnatal windows (P2–P9, P12–P18, or P21–P27) produced distinct and even opposing sex-dependent immune responses. P2–P9 treatment increased LY6C–/CX3CR1 + monocytes in adult males but not females, whereas P21–P27 treatment increased this population in females but not males; early P2–P9 treatment also increased microglial soma size selectively in females (Lynch et al., 2023). These findings highlight how relatively small shifts in the developmental timing of microbiome perturbation can produce markedly different sex-dependent neuroimmune outcomes. This is consistent with our finding that

female offspring from ABX-treated dams exhibit a distinct postnatal response to maternal microbiome perturbations, marked by microglial metabolic reprogramming and enhanced A β clearance.

Although this study provides important insights into how maternal microbiome perturbation shapes sex-dependent neuroimmune and metabolic pathways in AD, there are several limitations. Our broad-spectrum ABX-mediated maternal microbiome perturbation paradigm does not identify the specific taxa or metabolites responsible for the observed effects on offspring. Future studies using targeted microbial manipulations, gnotobiotic models, or metabolite supplementation will be required to define the microbial species and metabolic pathways that drive these neuroimmune changes. In addition, although our multi-omics analyses suggest a link between microbial metabolic alterations, GPCR signaling, and microglial remodeling, the causal receptor–ligand interactions remain to be experimentally validated. Functional studies using GPCR-specific perturbations, microglia-specific receptor knockouts, or metabolite restoration approaches will be necessary to directly test this signaling axis and determine how maternal microbiome perturbations influence neuroimmune trajectories across development and aging. An additional limitation is that the offspring were examined at 3 months of age. In the APPPS1-21 model, cortical A β plaque deposition begins at approximately 6–7 weeks of age and is already robust by 3 months, allowing us to examine the effects of maternal microbiome perturbation at a stage of established amyloid pathology. However, aging is accompanied by substantial changes in microglial phenotype and function, including the emergence of senescence-associated and dysfunctional states. Therefore, the microglial response to maternal microbiome perturbation and its relationship to A β pathology may differ in aged animals. Examining later ages will be important for determining whether the female-specific reduction in A β pathology and associated microglial phenotype observed at 3 months persist, diminish, or change with aging.

A potential caveat in this study is that while pups were still nursing before weaning, they may have consumed small amounts of ABX-containing water from the maternal drinking bottle, resulting in direct exposure estimated at approximately 1/50th of the gavage dose. Nevertheless, given the substantially lower potential direct exposure, the effects observed in this paradigm are expected to be predominantly driven by maternal microbiome perturbations. Future studies examining different developmental windows and exposing pups directly to this low ABX dose without altering the maternal microbiome will help further distinguish maternal from direct postnatal effects.

Collectively, our findings position the maternal microbiome as a lifelong determinant of the neuroimmune milieu, capable of biasing sex specific trajectories of neurodegenerative risk. By integrating microbiome, metabolomic, and transcriptomic data, our study supports a developmental model in which maternal microbiome perturbation programs female specific neuroprotection through metabolic and GPCR linked reorganization. In female offspring from ABX-treated dams, maternal dysbiosis induces amino acid restriction and one carbon imbalance that engage GPCR mediated nutrient sensing pathways, promoting microglial remodeling and enhanced A β clearance (Fig. 7), whereas males exhibit relative metabolic stability that limits transcriptional adaptation and fails to alter amyloid burden. Modulating the maternal microbiome or restoring key microbial metabolites either early or later in life

may therefore represent a promising avenue for precision medicine aimed at leveraging sex informed metabolic and immune mechanisms to influence neurodegenerative trajectories.

Material and Methods

Animal handling and housing

APPPS1-21 mice, acquired from M. Jucker at the University of Tübingen, Germany, were raised on a C57BL6Cj background (Radde et al., 2006). In this model, cortical A β plaque deposition begins at approximately 6–7 weeks of age and becomes robust by 3 months of age. The mice were kept in sterile micro-isolator cages and provided with standard chow ad libitum. All procedures involving animal experimentation were conducted following approved Animal Care and Use Protocols by the Institutional Animal Care and Use Committee at the University of Chicago. We hereby confirm that this study has been reported in accordance with the ARRIVE (Animal Research: Reporting of In Vivo Experiments) guidelines, as outlined at <https://arriveguidelines.org>.

Antibiotics treatment

Maternal microbiome depletion was performed following the protocol described by Vuong et al. 2018 (Vuong et al., 2020), consisting of an initial short term ABX gavage (ST ABX) followed by long term ABX exposure (1/50th of gavage dose) in drinking water (LT ABX). Beginning on day 28 (D28), both males and females assigned to the ABX cohort received the standard ABX cocktail by oral gavage for microbiome depletion. The cocktail included 4 mg/ml kanamycin (Sigma Aldrich K4000 5g), 0.35 mg/ml gentamicin (Sigma Aldrich G1914 250mg), 8,500 U/ml colistin (Sigma Aldrich C4461 1g), 2.15 mg/ml metronidazole (Sigma Aldrich M1547 25g), and 0.45 mg/ml vancomycin (Sigma Aldrich V2002 1g), prepared fresh in sterile water (Minter et al., 2016).

At 6 weeks of age, ABX treated males were paired with ABX treated females for mating. To prevent any paternal microbial influence during pregnancy, males were removed immediately upon confirmation of copulation (vaginal sperm detection or vaginal plug) and were not used for further experimental procedures. Following mating, dams transitioned to the LT ABX phase, in which the same ABX cocktail was provided in drinking water at 1/50 of the standard gavage dose throughout pregnancy and lactation. This ensured continuous maternal microbiome depletion from preconception through offspring postnatal day 21 (PND21). Control animals were housed identically but received autoclaved drinking water without ABX. Following weaning at PND21, offspring were group housed according to sex and experimental condition.

Pups remained with their ABX treated dams until PND21, receiving developmental exposure to maternal microbiome disruption without any direct ABX administration. After weaning, all offspring were placed on autoclaved drinking water (no ABX) and maintained under standard conditions until 3 months of age.

Necropsy and tissue harvesting

The procedure for harvesting various organs from mice at the time of sacrifice was conducted according to the approved and published Animal Care and Use Protocols of the University of Chicago. On the day of euthanasia, mice were given an intraperitoneal injection containing a mixture of ketamine and xylazine. After confirming deep anesthesia, blood samples were drawn through cardiac puncture using a 25-gauge needle, then stored in EDTA-coated collection tubes (BD Vacutainer; 365974) and placed on ice. Immediately following blood collection, the descending aorta was clamped, and the mice were perfused with physiological cold saline (pH 7.4) for 3 minutes. The brains were then removed from the skulls and divided into two hemispheres: one was postfixed with 4% paraformaldehyde for histological examination, and the other was immediately frozen for transcriptome analysis. The cecum was collected and weighed to assess the efficacy of the ABX treatments. Upon completion of these procedures, plasma was separated by centrifuging at 2,000 rpm for 10 minutes at 4°C using a Beckman Coulter centrifuge and stored at – 80°C for future use.

Cerebral cortex RNA extraction

Total RNA was extracted from the dorsal cerebral cortex using Trizol reagent, as per the protocol by Dodiya et al., and cleaned with the RNeasy Micro kit (Qiagen). Quality assessment was done with an Agilent Bioanalyzer. RNA-seq library preparations and sequencing were performed using an Illumina HiSeq4000 at The University of Chicago Genomics Core Facility. The data were collected in FASTQ format for bioinformatic analysis.

RNA-seq bioinformatics analysis

The quality of DNA reads, provided in FASTQ format, was assessed using FastQC. Adapters were trimmed, and reads that were either of poor quality or aligned to rRNA sequences were filtered using TrimGalore http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/. The cleaned reads were then aligned to the mouse genome (mm10) using STAR. Read counts for each gene were computed using HTSeq-Counts, in conjunction with a gene annotation file for mm10 obtained from Ensembl (<http://useast.ensembl.org/index.html>). A comprehensive quality control report was assembled using MultiQC. Differential expression was analyzed using DESeq2, with a cut-off for identifying significant differentially expressed genes (DEGs) set at an FDR-adjusted P value of < 0.05. Gene Ontology (GO) analysis and identification of DEGs within specific pathways were conducted using Metascape, and the findings were validated using the DAVID and Gorilla online services.

Fecal DNA extraction

Bacterial DNA was extracted from cecal and fecal contents following the method previously described and in accordance with standardized Earth Microbiome Project protocols (www.earthmicrobiome.org/emp-standard-protocols/). In brief, tissues were dissolved in an extraction buffer (50 mg tissue/ml buffer, 50 mM Tris (pH 7.4), 100 mM EDTA (pH 8.0), 400 mM NaCl, 0.5% w/v SDS) containing 0.4 mg/ml of proteinase K. Upon the addition of glass beads with a diameter of 0.1 mm (500 µl/ml buffer, BioSpec Products), microbial cells were lysed using a Mini-Beadbeater-8K Cell Disrupter (BioSpec Products), followed by overnight incubation in a water bath at 55°C. Total DNA was

subsequently extracted using a Phenol:Chloroform:IAA mixture (25:24:1 v/v, pH 8.0, Ambion), following the manufacturer's guidelines. The DNA yield was quantified, and its quality was determined using a combination of Nanodrop Lite (Thermo Fisher) and Qubit® fluorometer (Invitrogen) assessments.

16S rRNA gene Illumina® MiSeq sequencing

The 16S rRNA gene copy number was quantified from DNA isolated from fecal and cecal contents using quantitative PCR (QPCR). A concentration of 2–5 ng/μl DNA was combined with iQ-SYBR Green PCR Supermix (Bio-Rad), 518F (5'-TCC-TAC-GGG-AGG-CAG-CAG-T-3') and 338R (5'-GGA-CTA-CCA-GGG-TAT-CTA-ATC-CTG-TT-3') primers (2.5 μM), and Q-PCR was carried out on a LightCycler® 480 system (Roche). The reaction conditions were as follows: an initial denaturation at 95°C for 5 minutes, followed by 35 cycles of 95°C for 10 seconds, 64°C for 45 seconds, and 72°C for 45 seconds, and a final extension at 40°C for 30 seconds. The 16S rRNA gene copy number was determined by referencing Cp values to a standard curve created from the pCR4-TOPO plasmid containing the 16S rRNA gene amplicon. The copy number was then expressed relative to the specific DNA concentration added per reaction, as determined by a prior assessment using a Qubit® fluorometer (Invitrogen). All reactions were conducted in triplicate and included appropriate negative controls.

Fecal microbiota analysis

We utilized the Quantitative Insights Into Microbial Ecology (QIIME) pipeline and vsearch 8.1 to conduct standard demultiplexing and quality filtering. Unique ASVs were identified utilizing the Deblur method, and taxonomy was assigned through the Greengenes Database (May 2013 release; <http://greengenes.lbl.gov>). Statistical analyses were carried out in R, relying primarily on the 'vegan' and 'phyloseq' libraries.

Western Blot Analysis

Brain lysates were prepared from APPPS1-21 mice hemibrains using a lysis buffer composed of 50 mM Tris (pH 7.4), 150 mM NaCl, 5 mM EDTA, 0.5% NP-40, 0.5% sodium deoxycholate, 1x protease inhibitor cocktail, 1 mM PMSF, and 1x phosphatase inhibitor cocktail. Post-homogenization and sonication on ice, samples were centrifuged at 10,000 g for 20 minutes at 4°C. The supernatant was then collected as the total protein lysate. Protein concentrations were quantified using BCA analysis. For gel electrophoresis, samples were denatured at 95°C in reducing buffer prior to running on 12% glycine SDS-PAGE or 10–16.5% tricine SDS-PAGE gels, as described in Minter et al., 2016. This was followed by a wet transfer to a 0.2 μm nitrocellulose membrane. The membranes were blocked in 5% non-fat milk in PBS-T for 1 hour at room temperature. Full-length human APP (APP-FL) was detected using the C1/6.1 antibody (BioLegend; 1:1000). BACE1 (Abcam, #108394, 1:3000) and PS1-NT (Biolegend, #823404, 1:1000) were detected using their respective primary antibodies. β-Actin (Santa Cruz Biotechnology, #56459, 1:10000) and GAPDH (Cell Signaling Technology, #D4C6R, 1:10000) were used as loading controls and detected using the corresponding primary antibodies. Following washing, membranes were incubated with the appropriate HRP-conjugated secondary antibodies at a 1:10,000 dilution for 1 h at room temperature. Chemiluminescent signals were detected using Western Lightning® Plus ECL substrate (PerkinElmer).

and visualized using the ChemiDoc MP Imaging System (Bio-Rad). Densitometry analysis was performed using ImageJ.

Immunocytochemistry

To assess microglia in the context of A β deposits, immunofluorescence staining was performed, following a previously published protocol. Specifically, a complete series of 40- μ m-thick brain sections were stained for A β (3D6, 1:10,000) [9], while a half series was used for microglia staining with primary antibodies Iba1 (Invitrogen, 1:1000), P2RY12 (Invitrogen, 1:1000) and Clec7a (Invitrogen, 1:100). The procedure began with washing free-floating, level-matched 40- μ m sections in dilution media, with a duration of 60 minutes (10 min per wash). Sections were then immersed in serum blocking solution for an hour at ambient temperature, succeeded by overnight incubation with primary antibodies at 4°C. The subsequent day, the sections underwent a 60-minute wash in dilution media (10 min/wash), followed by a one-hour incubation with secondary antibodies at room temperature. Donkey anti-mouse 488 (Invitrogen; 1:500 and donkey anti-rabbit 647 (Invitrogen; 1:500) were employed as secondary antibodies for A β and microglia staining, correspondingly. Afterward, sections were washed, mounted on glass slides, and coverslipped using Fluoromount aqueous mounting medium (Sigma-Aldrich; F4680). The images of 3D6+ A β plaques were obtained with a 3D Histech Panoramic MIDI whole-slide scanner and a Zeiss AxioCam MRM CCD camera, handled by personnel from The University of Chicago Integrated Light Microscopy Facility. Microglia Iba1 images were taken utilizing a Leica SP5 3D STED laser-scanning confocal microscope at 40 $\times$ 1.5 oil magnification and Clec7a/P2ry12 images at 63 $\times$ oil magnification. For visualization of microglial morphology in three dimensions, confocal z-stacks were reconstructed using Imaris (Oxford Instruments). Three-dimensional reconstruction was used to visualize individual Iba1 + microglia and their spatial relationship to A β plaques. Quantitative analysis of individual microglial morphology was performed using Fiji/ImageJ. Individual Iba1 + microglia were analyzed for morphological parameters including soma area, total process length, number of branch points/junctions, and process endpoints. The same image-processing and quantification criteria were applied across experimental groups.

Abeta burden analysis

Evaluation of A β burden was conducted in accordance with a previously defined protocol. Briefly, six equidistant sections at 480 μ m intervals, ranging from beyond the olfactory bulb to the mid-hippocampal level, were employed for the analysis of 3D6 + amyloid plaques. The Fiji ImageJ software (NIH; ImageJ 1.51n) was used by two independent and blinded observers to determine the A β fractional area, also known as the amyloid burden. Individual images were processed using the 3D Histech Panoramic viewer software (3DHistech Kft), and subsequently normalized. Automated thresholding, based on histogram entropy, was applied to identify the amyloid plaques. After converting the images into an 8-bit format, a specific threshold number was applied, and this was followed by the use of the “fill holes” and “watershed” algorithms for binary conversion. Finally, the total area encompassed by plaques (i.e., A β burden) were computed using the “analyze particles” function. The A β burden, represented as the area

fraction, was calculated by dividing the total area occupied by plaques by the total area of the cerebral cortex, which was the region of interest.

Statistical analysis

Statistical analyses were carried out using the GraphPad Prism software, version 7.0e. For comparing the α - and β -diversity indices in the 16S rRNA microbiota analysis across multiple groups, the Wilcoxin analysis was utilized. This allowed for the careful comparison of distributions and medians among different groups, and corrections were made for multiple comparisons to ensure the integrity of the results. For other studies, one-way ANOVA (Analysis of Variance) was employed, unless otherwise specified in the methodology. The one-way ANOVA enabled the examination of differences among means for several groups, and it was followed by post hoc comparisons to identify which specific groups were different from each other. In all the analyses, a statistical P value below 0.05 was considered indicative of significant differences between the groups under comparison. Further details and specifics regarding the statistical methods and findings, including any additional tests or adjustments that were made, can be found in the figure legends accompanying the results.

Declarations

Ethics approval and consent to participate

All animal experiments were performed in accordance with institutional guidelines and were approved by the Institutional Animal Care and Use Committee (IACUC) at the University of Chicago.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by Cure Alzheimer's Fund, The Luminescence Foundation, the Open Philanthropy Project and Good Ventures Foundation, Creative Minds Care Foundation, and the BrightFocus Foundation.

Author Contribution

Author contributions P. Saha performed the ABX experiments; analyzed all omics datasets (including gut microbiota and RNA-seq); analyzed immunohistochemistry and western blot data; and performed SP5 laser confocal imaging., and prepared the final figures and manuscript; N. Slimmon assisted with immunohistochemistry using 3D6 antibody; P. Patel performed the bioinformatics analyses and assisted in generating RNA-seq figures; D. Mishra conducted the western blot experiments; A. Nangudga performed microglial staining; I. Weigle assisted with mouse sacrifice; P. P. Blauth carried out brain sectioning; A. Inukai and A. Fernandez performed RNA isolation; X. Zhang maintained and genotyped the mice; S. S. Sisodia developed the microbiota-related hypothesis, co-designed the experimental study, and reviewed and critiqued the manuscript.

Acknowledgement

Acknowledgement We thank the Slide Scanner Image Processing service at the University of Chicago Integrated Light Microscopy Core Facility. We thank Dr. Pieter W. Faber for his assistance with the Bioanalyzer and next-generation sequencing services at the University of Chicago Genomics Core Facility. We also thank Prof. Pamer for providing access to the Duchossois Family Institute facilities at the University of Chicago, including the full-service pipelines in metabolomics (DFI Microbiome Metabolomics Facility) and genomics (Microbiome Metagenomics Facility). These studies were supported by the Cure Alzheimer's Fund (to S.S. Sisodia), the Open Philanthropy Project and Good Ventures Foundation (to S.S. Sisodia), the Luminescence Foundation (to S.S. Sisodia), Creative Minds Care Foundation (to S.S. Sisodia) and the Safadi Program for Excellence in Clinical and Translational Neuroscience Postdoctoral Fellowship (to P. Saha).

Availability of data and materials

The datasets generated and/or analyzed during the current study are available from the corresponding author upon reasonable request. Sequencing data will be deposited in an appropriate public repository prior to publication.

References

1. BARBACID M. Structural and Functional Properties of the TRK Family of Neurotrophin Receptors. *Ann N Y Acad Sci.* 1995;766:442–58.
2. Biagioli V, Matera M, Ramenghi LA, Falsaperla R, Striano P. 2025. Microbiome and Pregnancy Dysbiosis: A Narrative Review on Offspring Health. *Nutrients* 17.
3. Bicknell B, Liebert A, Borody T, Herkes G, McLachlan C, Kiat H. Neurodegenerative and Neurodevelopmental Diseases and the Gut-Brain Axis: The Potential of Therapeutic Targeting of the Microbiome. *Int J Mol Sci.* 2023;24:9577.

4. Butovsky O, Weiner HL. Microglial signatures and their role in health and disease. *Nat Rev Neurosci.* 2018;19:622–35.
5. Cryan JF, O'Riordan KJ, Cowan CSM, Sandhu KV, Bastiaanssen TFS, Boehme M, Codagnone MG, Cusotto S, Fulling C, Golubeva AV, Guzzetta KE, Jaggar M, Long-Smith CM, Lyte JM, Martin JA, Molinero-Perez A, Moloney G, Morelli E, Morillas E, O'Connor R, Cruz-Pereira JS, Peterson VL, Rea K, Ritz NL, Sherwin E, Spichak S, Teichman EM, van de Wouw M, Ventura-Silva AP, Wallace-Fitzsimons SE, Hyland N, Clarke G, Dinan TG. The Microbiota-Gut-Brain Axis. *Physiol Rev.* 2019;99:1877–2013.
6. Di Gesu CM, Matz LM, Buffington SA. Diet-induced dysbiosis of the maternal gut microbiome in early life programming of neurodevelopmental disorders. *Neurosci Res.* 2021;168:3–19.
7. Dodiya HB, Kuntz T, Shaik SM, Baufeld C, Leibowitz J, Zhang X, Gottel N, Zhang X, Butovsky O, Gilbert JA, Sisodia SS. Sex-specific effects of microbiome perturbations on cerebral Abeta amyloidosis and microglia phenotypes. *J Exp Med.* 2019;216:1542–60.
8. Edwards SM, Cunningham SA, Dunlop AL, Corwin EJ. The Maternal Gut Microbiome During Pregnancy. *MCN Am J Matern Child Nurs.* 2017;42:310–7.
9. Ernfors P, Lee K-F, Jaenisch R. Mice lacking brain-derived neurotrophic factor develop with sensory deficits. *Nature.* 1994;368:147–50.
10. Erny D, Hrabé de Angelis AL, Jaitin D, Wieghofer P, Staszewski O, David E, Keren-Shaul H, Muhlaker T, Jakobshagen K, Buch T, Schwierzeck V, Utermohlen O, Chun E, Garrett WS, McCoy KD, Diefenbach A, Staeheli P, Stecher B, Amit I, Prinz M. Host microbiota constantly control maturation and function of microglia in the CNS. *Nat Neurosci.* 2015;18:965–77.
11. Gohmann L, Cusick JA, Demas GE, Wellman CL. Maternal Stress and Maternal Microbiome Manipulations Remodel Offspring Medial Prefrontal Cortex in a Sex-Dependent Manner. *Dev Neurobiol.* 2025;85:e22971.
12. Keleher MR, Zaidi R, Shah S, Oakley ME, Pavlatos C, Idrissi SE, Xing X, Li D, Wang T, Cheverud JM. Maternal high-fat diet associated with altered gene expression, DNA methylation, and obesity risk in mouse offspring. *PLoS ONE.* 2018;13:e0192606.
13. Kidana K, Tatebe T, Ito K, Hara N, Kakita A, Saito T, Takatori S, Ouchi Y, Ikeuchi T, Makino M, Saido TC, Akishita M, Iwatsubo T, Hori Y, Tomita T. 2018. Loss of kallikrein-related peptidase 7 exacerbates amyloid pathology in Alzheimer's disease model mice. *EMBO Mol Med* 10.
14. Koren O, Konnikova L, Brodin P, Mysorekar IU, Collado MC. The maternal gut microbiome in pregnancy: implications for the developing immune system. *Nat Rev Gastroenterol Hepatol.* 2024;21:35–45.
15. Kosakamoto H, Obata F, Kuraishi J, Aikawa H, Okada R, Johnstone JN, Onuma T, Piper MDW, Miura M. Early-adult methionine restriction reduces methionine sulfoxide and extends lifespan in *Drosophila*. *Nat Commun.* 2023;14:7832.
16. Latimer MN, Freij KW, Cleveland BM, Biga PR. Physiological and Molecular Mechanisms of Methionine Restriction. *Front Endocrinol (Lausanne).* 2018;9:217.

17. Lynch CMK, Cowan CSM, Bastiaanssen TFS, Moloney GM, Theune N, van de Wouw M, Florensa Zanuy E, Ventura-Silva AP, Codagnone MG, Villalobos-Manríquez F, Segalla M, Koc F, Stanton C, Ross P, Dinan TG, Clarke G, Cryan JF. Critical windows of early-life microbiota disruption on behaviour, neuroimmune function, and neurodevelopment. *Brain Behav Immun*. 2023;108:309–27.
18. Masuyama H, Mitsui T, Nobumoto E, Hiramatsu Y. The Effects of High-Fat Diet Exposure In Utero on the Obesogenic and Diabetogenic Traits Through Epigenetic Changes in Adiponectin and Leptin Gene Expression for Multiple Generations in Female Mice. *Endocrinology*. 2015;156:2482–91.
19. Mattioli I, Mantovani A, Locati M. The tetraspan MS4A family in homeostasis, immunity, and disease. *Trends Immunol*. 2021;42:764–81.
20. Minter MR, Hinterleitner R, Meisel M, Zhang C, Leone V, Zhang X, Oyler-Castrillo P, Zhang X, Musch MW, Shen X. Antibiotic-induced perturbations in microbial diversity during post-natal development alters amyloid pathology in an aged APPSWE/PS1ΔE9 murine model of Alzheimer's disease. *Sci Rep*. 2017;7:10411.
21. Minter MR, Zhang C, Leone V, Ringus DL, Zhang X, Oyler-Castrillo P, Musch MW, Liao F, Ward JF, Holtzman DM. Antibiotic-induced perturbations in gut microbial diversity influences neuro-inflammation and amyloidosis in a murine model of Alzheimer's disease. *Sci Rep*. 2016;6:30028.
22. Paolicelli RC, Sierra A, Stevens B, Tremblay M-E, Aguzzi A, Ajami B, Amit I, Audinat E, Bechmann I, Bennett M, Bennett F, Bessis A, Biber K, Bilbo S, Blurton-Jones M, Boddeke E, Brites D, Brône B, Brown GC, Butovsky O, Carson MJ, Castellano B, Colonna M, Cowley SA, Cunningham C, Davalos D, De Jager PL, de Strooper B, Denes A, Eggen BJL, Eyo U, Galea E, Garel S, Ginhoux F, Glass CK, Gokce O, Gomez-Nicola D, González B, Gordon S, Graeber MB, Greenhalgh AD, Gressens P, Greter M, Gutmann DH, Haass C, Heneka MT, Heppner FL, Hong S, Hume DA, Jung S, Kettenmann H, Kipnis J, Koyama R, Lemke G, Lynch M, Majewska A, Malcangio M, Malm T, Mancuso R, Masuda T, Matteoli M, McColl BW, Miron VE, Molofsky AV, Monje M, Mracsko E, Nadjar A, Neher JJ, Neniskyte U, Neumann H, Noda M, Peng B, Peri F, Perry VH, Popovich PG, Pridans C, Priller J, Prinz M, Ragozzino D, Ransohoff RM, Salter MW, Schaefer A, Schafer DP, Schwartz M, Simons M, Smith CJ, Streit WJ, Tay TL, Tsai L-H, Verkhratsky A, von Bernhardi R, Wake H, Wittamer V. Microglia states and nomenclature: A field at its crossroads. *Neuron*. 2022;110:3458–83. S.A. Wolf, L.-J. Wu, and T. Wyss-Coray.
23. Parums DV. A Review of the Current Status of Disease-Modifying Therapies and Prevention of Alzheimer's Disease. *Med Sci Monit*. 2024;30:e945091.
24. Price DL, Sisodia SS. Mutant genes in familial Alzheimer's disease and transgenic models. *Annu Rev Neurosci*. 1998;21:479–505.
25. Radde R, Bolmont T, Kaeser SA, Coomaraswamy J, Lindau D, Stoltze L, Calhoun ME, Jäggi F, Wolburg H, Gengler S, Haass C, Ghetti B, Czech C, Hölscher C, Mathews PM, Jucker M. Abeta42-driven cerebral amyloidosis in transgenic mice reveals early and robust pathology. *EMBO Rep*. 2006;7:940–6.

26. Ruhl DA, Bomba-Warczak E, Watson ET, Bradberry MM, Peterson TA, Basu T, Frelka A, Evans CS, Briguglio JS, Basta T, Stowell MHB, Savas JN, Roopra A, Pearce RA, Piper RC, Chapman ER. Synaptotagmin 17 controls neurite outgrowth and synaptic physiology via distinct cellular pathways. *Nat Commun.* 2019;10:3532.
27. Saha P, Weigle IQ, Slimmon N, Poli PB, Patel P, Zhang X, Cao Y, Michalkiewicz J, Gomm A, Zhang C, Tanzi RE, Dylla N, Al-Hendy A, Sisodia SS. Early modulation of the gut microbiome by female sex hormones alters amyloid pathology and microglial function. *Sci Rep.* 2024;14:1827.
28. Salia S, Burke FF, Hinks ME, Randell AM, Matheson MA, Walling SG, Swift-Gallant A. Gut microbiota transfer from the preclinical maternal immune activation model of autism is sufficient to induce sex-specific alterations in immune response and behavioural outcomes. *Brain Behav Immun.* 2025;123:813–23.
29. Thion MS, Low D, Silvin A, Chen J, Grisel P, Schulte-Schrepping J, Blecher R, Ulas T, Squarzoni P, Hoeffel G, Culpier F, Siopi E, David FS, Scholz C, Shihui F, Lum J, Amoyo AA, Larbi A, Poidinger M, Buttgereit A, Lledo PM, Greter M, Chan JKY, Amit I, Beyer M, Schultze JL, Schlitzer A, Pettersson S, Ginhoux F, Garel S. Microbiome Influences Prenatal and Adult Microglia in a Sex-Specific Manner. *Cell.* 2018;172:500–16. e516.
30. Vuong HE, Pronovost GN, Williams DW, Coley EJL, Siegler EL, Qiu A, Kazantsev M, Wilson CJ, Rendon T, Hsiao EY. The maternal microbiome modulates fetal neurodevelopment in mice. *Nature.* 2020;586:281–6.
31. Xi Y, Zhang Y, Zhou Y, Liu Q, Chen X, Liu X, Grune T, Shi L, Hou M, Liu Z. Effects of methionine intake on cognitive function in mild cognitive impairment patients and APP/PS1 Alzheimer's Disease model mice: Role of the cystathionine-beta-synthase/H(2)S pathway. *Redox Biol.* 2023;59:102595.
32. Xing Z, Tu BP. Mechanisms and rationales of SAM homeostasis. *Trends Biochem Sci.* 2025;50:242–54.
33. Yan C, He B, Wang C, Li W, Tao S, Chen J, Wang Y, Yang L, Wu Y, Wu Z. 2025. Methionine in embryonic development: metabolism, redox homeostasis, epigenetic modification and signaling pathway. *Crit Rev Food Sci Nutr* 1–24.
34. Young CE, Samuel MA. Microglia sensing of peripheral signals that bridge the brain and body. *Mol Neurodegener.* 2025;20:114.
35. Zhang Y, Wang Y, Li Y, Pang J, Höhn A, Dong W, Gao R, Liu Y, Wang D, She Y, Guo R, Liu Z. Methionine restriction alleviates diabetes-associated cognitive impairment via activation of FGF21. *Redox Biol.* 2024;77:103390.

Figures




Figure 1

Maternal ABX exposure induces persistent and sex-dependent gut microbiome remodeling. A, Experimental design. Female breeders received short-term antibiotic (ABX) treatment prior to mating followed by continued exposure during gestation. Offspring were weaned at P21 and analyzed at 3 months of age. Fecal samples from dams and offspring were collected for 16s rRNA sequencing. B, Microbiome profiling. a, Relative taxonomic abundance at the indicated taxonomic levels in maternal and

offspring microbiota following control or ABX exposure. b, Alpha diversity indices (Chao1 richness, Shannon index, and Inverse Simpson index). c, Unweighted UniFrac principal coordinates analysis (PCoA) showing phylogenetic β -diversity. d, Bray-Curtis PCoA showing compositional β -diversity. e, Differentially abundant taxa identified using *MaAsLin2*. Statistical significance for alpha diversity was determined using two-sided Wilcoxon rank-sum tests with false discovery rate (FDR) correction. β -diversity differences were assessed using two-sided PERMANOVA (999 permutations) with FDR correction. Differential abundance was determined using a two-sided linear model with Benjamini–Hochberg FDR correction (FDR < 0.05). Data represent individual biological replicates (n = 6 pups per group). *P < 0.05; **P < 0.01; NS, not significant.





Figure 2

Maternal antibiotic exposure alters cerebral cortical amyloid burden in a sex-dependent manner. A, Representative images of 3D6-immunolabeled amyloid plaques in the cerebral cortex of male and female offspring derived from control (MM(Ctrl)) and antibiotic-exposed (MM(ABX)) dams. B,

Quantification of cerebral cortical amyloid- β burden across groups. Statistical analysis was performed using one-way ANOVA followed by multiple-comparison correction. C–E, Western blot analysis and densitometric quantification of (C) full-length amyloid precursor protein (APP-FL), (D) β -site amyloid precursor protein cleaving enzyme 1 (BACE1), and (E) presenilin-1 N-terminal fragment (PS1-NT) in the cerebral cortex of male and female offspring. APP-FL and PS1-NT protein levels were normalized to β -actin, and BACE1 protein levels were normalized to GAPDH. Western blot comparisons were performed using two-sided Student's *t*-tests. Data are presented as mean $\pm$ s.e.m.; each point represents an independent biological replicate (immunohistochemistry: *n* = 6–9 per group; western blot: *n* = 3 per group). Scale bar, 50 μ m. *P* < 0.05; NS, not significant.

A Metagenomics analysis

a Piechart showing species distribution



b Linear regression–based correlation analysis





Figure 3

Maternal antibiotic exposure alters species-level microbial composition and intergroup correlations. a, Shotgun metagenomics analysis showing species-level distribution across MM(Ctrl)_Male, MM(ABX)_Male, MM(Ctrl)_Female and MM(ABX)_Female groups (n = 6 mice per group). Pie charts represent relative abundance of dominant bacterial species identified by metagenomic sequencing. b, Linear regression–based correlation analysis comparing relative species abundance between MM(ABX)_Female and the indicated groups. Correlation strength was assessed using two-sided linear regression analysis; the coefficient of determination (R^2) and corresponding P values are shown. Each point represents one bacterial species (n = 44 total species tested; species with missing or zero values in either group were excluded from regression). Statistical significance was defined as $P < 0.05$.

A Metabolomics analysis



B Functional metagenomics analysis




Figure 4

Maternal antibiotic exposure alters metabolite profiles and microbial functional pathways in a sex-dependent manner. a, Targeted metabolomics analysis showing relative abundance of the indicated metabolites in Male_Vehicle, Male_ABX, Female_Vehicle and Female_ABX groups (n = 6 mice per group; total n = 24). Bars represent mean relative abundance across biological replicates. b, Functional metagenomics analysis showing pathway abundance associated with *Lactobacillus johnsonii*(PWY-6151:

S-adenosyl-L-methionine salvage I) and *Limosilactobacillus reuteri* (PWY-724: superpathway of L-lysine, L-threonine and L-methionine biosynthesis. Relative abundance of *Lactobacillus johnsonii* (c) and *Lactobacillus reuteri* (d) across groups (n = 6 mice per group; total n = 24). Statistical significance was determined using two-sided Kruskal–Wallis tests followed by multiple-comparison correction. Data are presented as mean ± s.e.m., and each point represents one independent biological replicate. *P < 0.05; **P < 0.01.




Figure 5

Maternal antibiotic exposure induces sex-specific transcriptional remodeling in the cerebral cortex. A, Cortical RNA transcriptomic analysis of male and female offspring. Principal component analysis (PCA) of variance-stabilized normalized transcript counts demonstrates separation among MM(Ctrl)_Male, MM(ABX)_Male, MM(Ctrl)_Female, and MM(ABX)_Female groups; PC1 and PC2 explain 33% and 28% of the total variance, respectively. Differential gene expression was identified using DESeq2 with a two-sided Wald test for both up- and downregulated transcripts, followed by Benjamini–Hochberg false discovery rate (FDR) correction ($FDR < 0.05$). The overlap of significantly differentially expressed genes between Female_Ctrl versus Female_ABX and Male_Ctrl versus Male_ABX comparisons is shown. A heatmap of differentially expressed genes with hierarchical clustering was generated using Euclidean distance and complete linkage on scaled expression values. B, Pathway enrichment analysis of differentially expressed genes was performed using Gene Ontology (GO) and Reactome databases with a two-sided hypergeometric test against a whole-genome background, followed by multiple-hypothesis testing with FDR correction ($FDR < 0.05$). Enriched pathways are ranked by $-\log_{10}(P)$. Data represent $n = 6$ mice per group. Statistical significance was defined as $FDR < 0.05$.




Figure 6

Maternal antibiotic exposure alters microglial plaque association and phenotype in the female cortex. A, Representative confocal images of DAPI (blue), 3D6-immunolabeled amyloid plaques (green), and Iba1⁺ microglia (red) in the cerebral cortex of MM(Ctrl)_Female and MM(ABX)_Female mice. Merged images show Iba1⁺ microglia in close proximity to 3D6⁺ amyloid plaques. Quantification of Iba1⁺ microglial somata in close proximity to 3D6⁺ plaques is shown at right. B, Representative Imaris 3D reconstructions

of plaque-localized microglia in the cerebral cortex of MM(Ctrl)_Female and MM(ABX)_Female mice. Quantification of microglial morphological parameters, including (a) cell body area, (b) total dendritic length, (c) number of dendritic branches, (d) number of junctions, and (e) number of terminal points, is shown at right. C, Representative confocal images of 3D6-immunolabeled amyloid plaques (green), Clec7a⁺ microglia (yellow), and P2ry12⁺ microglia (red) in the cerebral cortex of MM(Ctrl)_Female and MM(ABX)_Female mice. Merged images show Clec7a⁺ and P2ry12⁺ microglia in close proximity to 3D6⁺ amyloid plaques. Quantification of P2ry12⁺ and Clec7a⁺ microglia in close proximity to 3D6⁺ plaques is shown at right. Statistical comparisons were performed using two-sided unpaired tests. Each point represents an independent biological replicate. Data are presented as mean $\pm$ s.e.m. Scale bars, 20 μ m (A, C) and 5 μ m (B). **P < 0.01; ****P < 0.0001.




Figure 7

Proposed model linking maternal microbiome disruption to sex-specific cortical remodeling and amyloid-associated microglial responses. Maternal antibiotic exposure alters gut microbial composition, leading to changes in microbial-derived metabolites and functional pathways. These circulating metabolites influence host signaling networks, including growth factor and neurotransmitter pathways, and modulate blood–brain communication. In female offspring, altered metabolic signaling reshapes

cortical transcriptional programs and microglial phenotype, influencing plaque-associated microglial responses and amyloid burden. This model integrates metagenomic, metabolomic and transcriptomic findings to propose a gut–metabolite–brain axis mediating sex-dependent effects on neuroinflammatory and synaptic regulation pathways.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Figure2S.pdf](#)
- [Figure4S.pdf](#)
- [Figure1S.pdf](#)