

Chronic Illness Burden Regular Healthcare Providers and Telehealth Use among African American and Latino Public Housing Residents

Mohsen Bazargan

Charles R. Drew University of Medicine and Science

Ehsan Yaghmaei

Charles R. Drew University of Medicine and Science

Attallah Dillard

Charles R. Drew University of Medicine and Science

Shervin Assari

shervinassari@cdrewu.edu

Charles R. Drew University of Medicine and Science

Research Article

Keywords: telehealth, racial and ethnic minorities, public housing, intersectionality, health disparities, mediation analysis, health needs, chronic illness

Posted Date: October 29th, 2026

DOI: <https://doi.org/10.21203/rs.3.rs-11537261/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Additional Declarations: The authors declare no competing interests.

Abstract

Background

Access to an established healthcare provider may help explain differences in telehealth use among residents of public housing.

Objective

As a continuation of Cobb et al. (2025), this study used the same survey dataset to examine whether having a regular healthcare provider may account for the association between chronic illness burden and telehealth use.

Methods

This cross-sectional analysis included 252 African American and Latino adults living in four public housing developments in South Los Angeles. A structural equation model examined associations among chronic illness burden, regular provider status, and telehealth use, accounting for age, gender, education, and language.

Results

Chronic illness burden was positively associated with having a regular provider ($B = 0.158$, $p = 0.015$), and regular provider status was positively associated with telehealth use ($B = 0.145$, $p = 0.016$). The adjusted direct association between chronic illness burden and telehealth use was not statistically significant ($B = 0.094$, $p = 0.141$).

Conclusion

The observed associations are consistent with a possible pathway through access to a regular provider. They do not establish mediation or causal direction. Improving access to ongoing care may be relevant to efforts to support telehealth use in this population.

1. Background

Telehealth use among public housing residents needs to be understood within the conditions in which care is sought. Housing disadvantage and racial inequities may shape access to both local healthcare and the resources needed for remote visits [1–6]. An intersectional approach considers how these conditions occur together, rather than treating income, racial and ethnic identity, and housing as separate

influences [7, 8]. For African American and Latino residents with limited financial resources, the availability of a clinician may be as relevant to telehealth use as the availability of technology.

A greater need for care does not necessarily translate into greater use of services. People with low incomes may encounter costs, transportation problems, and limited access to providers even when they have ongoing health concerns [9–13]. Residential segregation and housing policies may further shape where services are located and which communities can reach them [14–18]. Research on healthcare needs and service access in underserved settings supports examining the connection between illness burden and access to a regular source of care [19, 20].

Language and access to health information may also influence how residents seek care. Studies of Latino help-seeking and of health information use among underserved populations suggest that needs, preferences, and available resources differ across groups [23–28]. These differences make it useful to examine telehealth alongside established patterns of healthcare access, rather than assuming that a remote visit is equally accessible to all residents.

Cobb et al. [32] previously analyzed telehealth use in this same sample of African American and Latino public housing residents. That study examined perceived benefits and self-efficacy as potential mediators and did not find evidence supporting those pathways. The present analysis continues that work by shifting attention to an aspect of healthcare access: whether residents have a regular healthcare provider. Telehealth may reduce travel demands [29], but arranging a remote visit may still depend on having a clinical connection through which care can be obtained.

Using the same survey dataset as Cobb et al. [32], we examined whether chronic illness burden was associated with regular provider status and whether regular provider status was associated with telehealth use. The proposed pathway differs from the attitude-based pathways assessed previously. This is a secondary analysis of an existing sample, not an independent replication or a new round of data collection.

2. Methods

2.1 Study Design

This cross-sectional secondary analysis continues the research reported by Cobb et al. [32]. Both reports use the same face-to-face survey of public housing residents. The earlier publication focused on telehealth attitudes; the present analysis examines a proposed pathway from chronic illness burden through regular provider status to telehealth use.

2.2 Participants

The analytic sample comprised 252 adults who identified as African American or Latino and lived in four public housing developments in South Los Angeles. Interviews were conducted from August 2023 through January 2024. The four developments were located in Service Planning Area 6 and contained

2,306 housing units in total. They were selected from the 14 developments managed by the Housing Authority of the City of Los Angeles. Additional details about the setting and recruitment are available in Cobb et al. [32].

2.3 Sampling

Recruitment was nonrandom. Eligible residents were at least 18 years old, lived in Service Planning Area 6, identified as African American or Latino, and could complete an interview in English or Spanish. The current report uses the participants included in the earlier analysis [32].

2.4 Ethics

Participants gave written informed consent before the interviews. Charles R. Drew University of Medicine and Science approved the study protocol through its Institutional Review Board. The survey covered social and demographic characteristics, health conditions, and experiences with healthcare access and use [32].

2.5 Measurement

Measures were drawn from the survey described by Cobb et al. [32]. The present analysis retained the variables relevant to illness burden, regular provider access, and telehealth use.

2.5.1 Socioeconomic and Demographic Variables

Age: Age in years was reported by participants; observed ages ranged from 18 to 77.

Gender: The survey recorded self-reported gender as female (1) or male (2).

Education: Respondents selected their highest completed level of schooling, from eighth grade or less through a bachelor's degree or higher. Table 1 combines the upper education categories, as in the earlier report [32].

Primary language: Interview data distinguished English (1) from Spanish (2) as the participant's primary language.

Chronic illness burden: A count summarized participants' self-reported diagnosed chronic conditions [32].

Regular healthcare provider: Respondents indicated whether they had a doctor or other provider for primary care. Responses were coded no (0) and yes (1). This indicator represents access to a regular provider; it does not directly measure the duration or quality of continuity of care.

2.5.2 Telehealth Utilization

Telehealth use: The survey asked about healthcare visits conducted remotely using a telephone, smartphone, computer, or tablet [32].

Telehealth frequency: Respondents reported the number of telehealth visits in the preceding 12 months. Table 1 presents the distribution of those visits.

2.6 Statistical analysis

Descriptive statistics and the relevant subset of bivariate correlations summarize the shared dataset previously reported by Cobb et al. [32]. The present structural equation model specifies chronic illness burden as a predictor of regular provider status and telehealth use, with regular provider status also predicting telehealth use. Age, gender, education, and primary language are included in both equations. Table 3 reports coefficients (B), standard errors, 95% confidence intervals, and p values.

3. Results

Table 1 describes the same 252 participants reported by Cobb et al. [32]. Their mean age was 45 years (SD = 15), and 188 (75%) were women. Educational attainment was eighth grade or less for 99 participants (39%), some high school without a diploma for 56 (22%), and high school completion or a GED for 66 (26%). The remaining 31 participants (12%) were in the combined category of some college, an associate degree, or a bachelor’s degree.

A regular provider was reported by 207 participants (82%), and 187 (74%) identified Spanish as their primary language. The chronic illness count was zero for 122 participants (48%), one for 47 (19%), two for 39 (15%), and three or more for 44 (18%).

During the preceding year, 133 participants (53%) reported no telehealth visits. The counts for one, two, three, and four or more visits were 50, 37, 19, and 13, respectively. These descriptive results overlap with the earlier publication because both analyses use the same sample [32].

Table 1
Characteristics of the shared study sample (N = 252).

Adapted from Cobb et al. [32]; retained values are unchanged, with measures outside the present analysis omitted.

Age (Yrs)	Mean	SD
	45	15
Number of Chronic Illnesses (0–8)	1	2
	N	%
Gender		
Male	64	25
Female	188	75
Education		
8th Grade or Less	99	39
Some High School, No diploma	56	22
Graduated from High/ GED equivalent	66	26
Associate Degree or Some College/Bachelor's Degree	31	12
Language		
English	65	26
Spanish	187	74
Primary Care Provider		
No	45	18
Yes	207	82
Chronic Illnesses		
0	122	48
1	47	19
2	39	15
3 or more	44	18
Used Telehealth in the Past 12 Months		
0	133	53
1	50	20

Age (Yrs)	Mean	SD
	45	15
2	37	15
3	19	7
4 or more	13	5

Table 2 presents the subset of previously reported correlations relevant to this analysis [32]. Telehealth use was positively correlated with chronic illness burden and regular provider status (both $r = 0.20$, $p < 0.01$, at the precision shown in the table). Negative correlations with male gender ($r = -0.14$) and Spanish language ($r = -0.15$) were significant at $p < 0.05$. Age and education were not significantly correlated with telehealth use. These coefficients describe the shared dataset and are not new estimates from an independent sample.

Table 2
Correlations among variables retained in the present analysis.

Adapted from Cobb et al. [32]; all retained coefficients and significance markers are unchanged.

1. Age	1	2	3	4	5	6	7
	1.00						
2. Gender	-0.08	1.00					
3. Education	-0.49**	0.08	1.00				
4. Language	0.32**	-0.14*	-0.51**	1.00			
5. Number of Chronic Illnesses	0.34**	-0.05	-0.13*	0.05	1.00		
6. Primary Care Provider	0.09	-0.11	-0.01	-0.04	0.18**	1.00	
7. Telehealth Care Utilization	0.04	-0.14*	0.03	-0.15*	0.20**	0.20**	1.00
Note: ** Correlation is significant at the 0.01 level (two-tailed), and * correlation is significant at the 0.05 level (two-tailed).							

Table 3 reports the model examined in this continuation study. Unlike the earlier model of telehealth attitudes [32], it includes regular provider status as an intermediate variable between illness burden and telehealth use. The reported coefficients are described below.

Predictors of Telehealth Utilization

The analysis examined factors associated with telehealth utilization among low-income racial and ethnic minority residents of public housing in Los Angeles. Having a regular healthcare provider was positively

associated with telehealth use ($B = 0.145$, $SE = 0.060$, $95\% \text{ CI} = 0.027, 0.264$, $p = 0.016$), indicating that individuals with a designated provider were more likely to use telehealth services.

Gender was significantly associated with telehealth use ($B = -0.153$, $SE = 0.060$, $95\% \text{ CI} = -0.270, -0.035$, $p = 0.011$), suggesting that females were more likely to engage with telehealth services compared to males. Additionally, Spanish as the primary language was negatively associated with telehealth utilization ($B = -0.230$, $SE = 0.069$, $95\% \text{ CI} = -0.365, -0.095$, $p = 0.001$), implying that Spanish-speaking participants were less likely to use telehealth compared to English-speaking individuals.

Neither age ($B = 0.049$, $SE = 0.073$, $95\% \text{ CI} = -0.093, 0.192$, $p = 0.497$) nor education level ($B = -0.049$, $SE = 0.076$, $95\% \text{ CI} = -0.197, 0.099$, $p = 0.517$) showed a significant relationship with telehealth use. Similarly, the number of chronic illnesses was not significantly associated with telehealth utilization ($B = 0.094$, $SE = 0.064$, $95\% \text{ CI} = -0.031, 0.220$, $p = 0.141$).

Predictors of Having a Regular Healthcare Provider

The second model assessed predictors of having a regular healthcare provider, which was hypothesized to mediate the relationship between chronic illness and telehealth use.

Chronic illness burden was positively associated with having a regular healthcare provider ($B = 0.158$, $SE = 0.065$, $95\% \text{ CI} = 0.031, 0.285$, $p = 0.015$), suggesting that individuals with a higher number of chronic illnesses were more likely to have an established healthcare provider.

Other factors, including age ($B = 0.053$, $SE = 0.075$, $95\% \text{ CI} = -0.094, 0.200$, $p = 0.477$), gender ($B = -0.108$, $SE = 0.062$, $95\% \text{ CI} = -0.229, 0.013$, $p = 0.081$), education level ($B = 0.011$, $SE = 0.078$, $95\% \text{ CI} = -0.142, 0.164$, $p = 0.890$), and primary language ($B = -0.072$, $SE = 0.072$, $95\% \text{ CI} = -0.214, 0.069$, $p = 0.315$), were not significant predictors of having a regular healthcare provider.

Table 3
Structural equation model examining regular provider status and telehealth use.

Telehealth use	B	SE	95% CI	p
Regular Healthcare Provider	0.145	0.060	0.027 0.264	0.016
Age	0.049	0.073	-0.093 0.192	0.497
Gender	-0.153	0.060	-0.270 -0.035	0.011
Education	-0.049	0.076	-0.197 0.099	0.517
Spanish Language	-0.230	0.069	-0.365 -0.095	0.001
Chronic Illness (n)	0.094	0.064	-0.031 0.220	0.141
Intercept	1.871	0.505	0.881 2.861	< 0.001
Regular Healthcare Provider				
Age	0.053	0.075	-0.094 0.200	0.477
Gender	-0.108	0.062	-0.229 0.013	0.081
Education	0.011	0.078	-0.142 0.164	0.890
Spanish Language	-0.072	0.072	-0.214 0.069	0.315
Chronic Illness (n)	0.158	0.065	0.031 0.285	0.015
Intercept	2.448	0.510	1.449 3.447	< 0.001
Mean				
Age	2.949	0.146	2.663 3.234	< 0.001
Gender	2.881	0.143	2.601 3.161	< 0.001
Education	1.987	0.109	1.774 2.199	< 0.001
Spanish Language	3.982	0.188	3.613 4.351	< 0.001
Chronic Illness (n)	0.744	0.071	0.605 0.884	< 0.001

4. Discussion

Access to a regular provider was associated with telehealth use in this sample, while chronic illness burden was associated with having a regular provider. The adjusted direct association between illness burden and telehealth use was not statistically significant. This pattern is compatible with the proposed pathway, but the available results do not establish that provider status mediates the association or fully explains it.

This report extends Cobb et al. [32] by examining a different question within the same dataset. The shared participants, descriptive statistics, and correlations should therefore be understood as the basis for a continuation analysis. The potential contribution lies in examining regular provider access as an intermediate variable, rather than in presenting a new sample or independent confirmation of earlier findings.

The earlier study did not support perceived benefits or self-efficacy as mediators of telehealth use [32]. The current findings suggest that access to an established clinician deserves further study alongside these attitudes. A provider relationship may offer a route into remote care, although the present survey cannot determine whether that relationship preceded telehealth use or resulted partly from greater healthcare contact.

Education was not significantly associated with either regular provider status or telehealth use in the current model. Marginalization-Related Diminished Returns theory offers one possible context for interpreting weak associations between education and health-related resources [33, 34]. However, this analysis does not directly test that theory: it includes neither a White comparison group nor an estimate of differences in educational returns across racial groups. The null findings may also reflect limited variation, measurement, or statistical precision.

Women and English-speaking participants reported greater telehealth use in the adjusted model. Prior work has described gender differences in healthcare use [35–39] and language-related barriers to services [40–43]. In this setting, language support and accessible communication may be relevant to remote care. The associations do not identify the specific barriers encountered by participants, and explanations based on language access should be examined directly in future work.

A regular provider may help patients learn about remote appointments, arrange follow-up, and maintain contact with services. Earlier studies connect regular sources of care with access and patient-reported care experiences [47–50]. These possible explanations remain hypotheses here. Having a regular provider is a limited indicator of access and should not be treated as a direct measure of trust, relationship quality, or continuity over time.

Several limitations affect interpretation. The cross-sectional design does not establish the order of illness burden, provider access, and telehealth use, and greater use of care could itself increase the likelihood of reporting a regular provider. Nonrandom recruitment from four developments limits generalizability beyond the surveyed residents. Self-reported measures may be affected by recall and reporting differences. Insurance coverage, digital access, disability, provider availability, and other unmeasured factors could contribute to the observed associations. In addition, the reported paths alone do not demonstrate mediation; an estimated indirect effect with uncertainty is required. Because this is the same dataset analyzed by Cobb et al. [32], agreement between the two reports is not evidence of independent replication.

Future studies could follow residents over time to assess whether gaining a regular provider is followed by changes in telehealth use. Interviews could clarify how appointments are arranged and which language, technology, and service barriers residents encounter. Analyses that measure continuity directly, account for digital and insurance access, and estimate indirect effects would help assess the pathway proposed here. Broader recruitment would also help determine whether the pattern applies to other public housing communities.

5. Conclusions

In this continuation of Cobb et al. [32], chronic illness burden was associated with having a regular healthcare provider, and regular provider status was associated with telehealth use. These findings support further examination of provider access as a possible link between health needs and remote care. They do not establish a causal pathway or full mediation. Efforts to improve telehealth access may benefit from considering ongoing access to providers and language support alongside technology and patient attitudes.

Declarations

Author Contributions: Methodology, A.D., M.B. and S.A.; Software, E.Y.; Conceptualization, S.A. and M.B.; Formal analysis, A.D. and E.Y.; Data curation, A.D. and S.A.; Writing—original draft, S.A.; Writing—review and editing, E.Y., M.B., A.D. and S.A.; Funding acquisition, M.B. All authors have read and agreed to the published version of the manuscript.

Funding: Mohsen Bazargan (Co-PI), Ehsan Yaghmaei (Statistician), and Shervin Assari (Co-I) receive support from Clinical Research Education and Career Development (CRECD) program at Charles R. Drew University of Medicine and Science through the NIMHD/NIH Award number R25 MD007610.

Institutional Review Board Statement: The study protocol received approval from the Institutional Review Board (IRB) at Charles R. Drew University of Medicine and Science (IRB Approval Number: 1746099, approval date: 22 April 2021).

Informed Consent Statement: Informed consent was obtained from all subjects involved in this study.

Data Availability Statement: The original contributions presented in this study are included in this article. Further inquiries can be directed to the corresponding authors.

Acknowledgments: The researchers wish to acknowledge the participants as well as the Housing Authority of the City of Los Angeles (HACLA) who made this study possible.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Aratani, Y. Public housing revisited: Racial differences, housing assistance, and socioeconomic attainment among low-income families. *Social Science Research* 2010, 39, 1108-1125.
2. Fauth RC, Leventhal T, Brooks-Gunn J (2004) Short-term effects of moving from public housing in poor to middle-class neighborhoods on low-income, minority adults' outcomes. *Soc Sci Med* 59:2271–2284
3. Carter WH, Schill MH, Wachter SM (1998) Polarisation, public housing and racial minorities in US cities. *Urban Stud* 35:1889–1911
4. Murdie RA (1994) Blacks in near-ghettos? black visible minority population in metropolitan Toronto housing authority public housing units. *Hous Stud* 9:435–457
5. DeLuca S, Garboden PM, Rosenblatt P (2013) Segregating shelter: How housing policies shape the residential locations of low-income minority families. *ANNALS Am Acad Political Social Sci* 647:268–299
6. Jahoda M, West PS (1951) Race relations in public housing. *J Soc Issues* 7
7. Bowleg L (2012) The problem with the phrase women and minorities: intersectionality—an important theoretical framework for public health. *Am J Public Health* 102:1267–1273
8. Atewologun D (2018) Intersectionality theory and practice. 1–20
9. Dutton DB (1978) Explaining the low use of health services by the poor: costs, attitudes, or delivery systems? *Am Sociol Rev* 348–368
10. Price JH, Desmond SM, Snyder FF, Kimmel SR (1988) Health Care and Poor Patients. *J Fam Pract* 27:615–621
11. Kirby JB (2008) Poor people, poor places and access to health care in the United States. *Soc Forces* 87:325–355
12. Preker AS, Carrin G, Dror D, Jakab M, Hsiao W, Arhin-Tenkorang D (2004) Rich-poor differences in health care financing. *Health Financing for Poor People—Resource Mobilization and Risk Sharing*, pp 3–52
13. Carder P, Luhr G, Kohon J (2016) Differential Health and Social Needs of Older Adults Waitlisted for Public Housing or Housing Choice Vouchers. *J Aging Soc Policy* 28:246–260. 10.1080/08959420.2016.1156507
14. Fagan J, Davies G, Carlis A (2012) Race and selective enforcement in public housing. *J Empir legal Stud* 9:697–728
15. Goetz EG (2011) Where have all the towers gone? The dismantling of public housing in US cities. *J Urban Affairs* 33:267–287
16. Holloway SR, McNulty TL (2003) Contingent urban geographies of violent crime: Racial segregation and the impact of public housing in Atlanta. *Urban Geogr* 24:187–211
17. Haley DF, Linton S, Luo R, Hunter-Jones J, Adimora AA, Wingood GM, Bonney L, Ross Z, Cooper HL (2017) Public Housing Relocations and Relationships of Changes in Neighborhood Disadvantage

- and Transportation Access to Unmet Need for Medical Care. *J Health Care Poor Underserved* 28:315–328. 10.1353/hpu.2017.0026
18. Yamamoto T, Hanazato M, Hikichi H, Kondo K, Osaka K, Kawachi I, Aida J (2023) Change in Geographic Accessibility to Dental Clinics Affects Access to Care. *J Dent Res* 102:719–726. 10.1177/00220345231167771
 19. Hodgkinson S, Godoy L, Beers LS, Lewin A (2017) Improving Mental Health Access for Low-Income Children and Families in the Primary Care Setting. *Pediatrics* 139. 10.1542/peds.2015-1175
 20. Lascher S, Tasir-Rodriguez W, Moon G, Irizzary M, Baney M, Kellogg FR (2013) Government public housing health needs assessment: focus on race, ethnicity, and the older adult: background, methods, and demographics. *Care Manag J* 14:118–124. 10.1891/1521-0987.14.2.118
 21. Foster B, Ledgerwood DM, Struble CA, Fodor MC, Jordan P, Krishnan AC, Turner B, Pearson C, Twiner MJ, Levy PD (2022) Public Housing Resident Perspectives on Smoking, Barriers for Smoking Cessation, and Changes in Smoking Mandates. *Inquiry* 59:469580221092814. 10.1177/00469580221092814
 22. Wang X, Seyler BC, Chen T, Jian W, Fu H, Di B, Yip W, Pan J (2024) Disparity in healthcare seeking behaviors between impoverished and non-impoverished populations with implications for healthcare resource optimization. *Humanit Social Sci Commun* 11:1208. 10.1057/s41599-024-03712-z
 23. Gearing RE, Washburn M, Brewer KB, Yu M, Bjugstad A, Torres LR (2023) Predictors of Help-Seeking for Mental Health Treatment Among Latinos. *J Racial Ethn Health Disparities* 10:1869–1877. 10.1007/s40615-022-01370-y
 24. Siddiqui NY, Ammarell N, Wu JM, Sandoval JS, Bosworth HB (2016) Urinary Incontinence and Health-Seeking Behavior Among White, Black, and Latina Women. *Female Pelvic Med Reconstr Surg* 22:340–345. 10.1097/spv.0000000000000286
 25. Powe BD (2015) Health Information Seeking Among Rural African Americans, Caucasians, and Hispanics: It Is Built. Did They Come? *Nurs Clin North Am* 50:531–543. 10.1016/j.cnur.2015.05.007
 26. Lee YJ, Boden-Albala B, Larson E, Wilcox A, Bakken S (2014) Online health information seeking behaviors of Hispanics in New York City: a community-based cross-sectional study. *J Med Internet Res* 16:e176. 10.2196/jmir.3499
 27. Bjarnadottir RI, Millery M, Fleck E, Bakken S (2016) Correlates of online health information-seeking behaviors in a low-income Hispanic community. *Inf Health Soc Care* 41:341–349. 10.3109/17538157.2015.1064429
 28. Sherman LD, Goidel K, Bergeron CD, Smith ML (2021) Web-Based Health Information Seeking Among African American and Hispanic Men Living With Chronic Conditions: Cross-sectional Survey Study. *J Med Internet Res* 23:e26180. 10.2196/26180
 29. Jezewski E, Miller A, Eusebio M, Potter J (2022) Targeted telehealth education increases interest in using telehealth among a diverse group of low-income older adults. *Int J Environ Res Public Health* 19:13349

30. Luo J, Ahmad SF, Alyaemeni A, Ou Y, Irshad M, Alyafi-Alzahri R, Alsanie G, Unnisa ST (2024) Role of perceived ease of use, usefulness, and financial strength on the adoption of health information systems: the moderating role of hospital size. *Humanit Social Sci Commun* 11:516. 10.1057/s41599-024-02976-9
31. Broffman L, Harrison S, Zhao M, Goldman A, Patnaik I, Zhou M (2023) The Relationship Between Broadband Speeds, Device Type, Demographic Characteristics, and Care-Seeking Via Telehealth. *Telemed J E Health* 29:425–431. 10.1089/tmj.2022.0058
32. Cobb S, Dillard A, Yaghmaei E, Bazargan M, Assari S (2025) Telehealth Perceived Benefits and Self-Efficacy Do Not Mediate the Effects of Demographic, Health, and Social Determinants on Telehealth Use of Low-Income African American and Latino Residents of Public Housing in Los Angeles. *Healthcare* 13:286. <https://doi.org/10.3390/healthcare13030286>
33. Assari S (2018) Health Disparities due to Diminished Return among Black Americans: Public Policy Solutions. *Social Issues Policy Rev* 12:112–145. 10.1111/sipr.12042
34. Assari S (2018) Unequal Gain of Equal Resources across Racial Groups. *Int J Health Policy Manag* 7:1–9. 10.15171/ijhpm.2017.90
35. Liu L, Wang C, Yang S, Lo K, Yang X (2020) Gender Differences in Healthcare-Seeking Behaviors in China's Ageing Population. An Empirical Study of Shanghai
36. Kana S, Arjyal A, Wallace L, Nwameme A, Mbachu C, Islam F, Huque S, Salma U, Shrestha G, Elsey H (2023) Influence of changing gender and social norms on health seeking behaviour and health systems in cities
37. Onah MN, Horton S (2018) Male-female differences in households' resource allocation and decision to seek healthcare in south-eastern Nigeria: Results from a mixed methods study. *Soc Sci Med* 204:84–91
38. Cameron KA, Song J, Manheim LM, Dunlop DD (2010) Gender disparities in health and healthcare use among older adults. *J women's health* 19:1643–1650
39. Bertakis KD, Azari R, Helms LJ, Callahan EJ, Robbins JA (2000) Gender differences in the utilization of health care services. *J Fam Pract* 49
40. Cheng EM, Chen A, Cunningham W (2007) Primary language and receipt of recommended health care among Hispanics in the United States. *J Gen Intern Med* 22:283–288
41. Timmins CL (2002) The impact of language barriers on the health care of Latinos in the United States: a review of the literature and guidelines for practice, vol 47. *Journal of midwifery & women's health*, pp 80–96
42. Diamond LC, Tuot DS, Karliner LS (2012) The use of Spanish language skills by physicians and nurses: policy implications for teaching and testing. *J Gen Intern Med* 27:117–123
43. DuBard CA, Gizlice Z Language spoken and differences in health status, access to care, and receipt of preventive services among US Hispanics. *Am J public health* 2008, 98, 2021–2028
44. Al-Amer R, Ramjan L, Glew P, Darwish M, Salamonson Y (2015) Translation of interviews from a source language to a target language: Examining issues in cross-cultural health care research. *J Clin*

45. Gil-Salmerón A, Katsas K, Riza E, Karnaki P, Linos A (2021) Access to healthcare for migrant patients in Europe: healthcare discrimination and translation services. *Int J Environ Res Public Health* 18:7901
46. Matías-Carrelo LE, Chávez LM, Negrón G, Canino G, Aguilar-Gaxiola S, Hoppe S (2003) The Spanish translation and cultural adaptation of five mental health outcome measures. *Cult Med Psychiatry* 27:291–313
47. Beal A, Hernandez S (2010) Patient reports of the quality of care in community health centers: the importance of having a regular provider. *J Health Care Poor Underserved* 21:591–605
48. Lambrew JM, Defrieze GH, Carey TS, Ricketts TC, Biddle AK (1996) The effects of having a regular doctor on access to primary care. *Med Care* 34:138–151
49. Ettner SL (1999) The relationship between continuity of care and the health behaviors of patients: does having a usual physician make a difference? *Med Care* 37:547–555
50. Mahmood A, Mosalpuria K, Wyant DK, Bhuyan SS (2019) Association between having a regular health provider and access to services linked to electronic health records. *Hosp Top* 97:1–10

Figures




Figure 1

Proposed model relating chronic illness burden, regular provider status, and telehealth use. Chr Ill: number of chronic illnesses; Reg Prov: regular healthcare provider; Educ: educational attainment; Language: Spanish; Gender: male.