


Title:

Cytokines and vascular mediators fingerprints based on interaction networks in kidney transplanted and non-transplanted moderately ill COVID-19 patients.

Authors:

Milena Karina Coló Brunialti, Giuseppe Gianini Figueirêdo Leite, Orlei Ribeiro de Araújo², Gabriela Strafolino Eburneo, Paula M Peçanha-Pietrobom, Paulo Roberto Abrão Ferreira, Nancy Junqueira Bellei, Jaqueline Sonoe Ota Arakaki, Lúcio R. Requião-Moura, Reinaldo Salomao

Supplementary material 1: Comparison of the means of the different mediators in the groups of patients with COVID-19 and healthy controls (HC) on the day of ward admission.



Two-tailed unpaired Student's t-test was applied; * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, and **** $P < 0.0001$ are represented in the graph.

Abbreviations: G-CSF, granulocyte colony-stimulating factor; GDF-15, growth/differentiation factor; IL, interleukin; MPO, myeloperoxidase; NGAL, neutrophil gelatinase-associated lipocalin; P-Sel, P-selectin; sICAM-1, soluble intercellular adhesion molecule-1; sVCAM-1, soluble circulating vascular cell adhesion molecule-1.