

Table 1

Demographics and Clinicopathologic Characteristics of SEER Patients With cNET

Category	Training cohort (n=628)		Total cohort (n=894)	P
	No. of patients	(%)		
Age(year)				0.498
<44	268 (42.7%)		372 (41.6%)	
44-67	263 (41.9%)		377 (42.2%)	
>67	97 (15.4%)		145 (16.2%)	
Year at diagnosis				0.188
2004-2009	294 (46.8%)		432 (48.3%)	
2010-2015	334 (53.2%)		462 (51.7%)	
Marital status				0.901
Married	281 (44.7%)		401 (44.9%)	
Single	247 (39.4%)		348 (38.9%)	
Other	100 (15.9%)		145 (16.2%)	
Race				0.275
White	456 (72.6%)		636 (71.1%)	
Black	101 (16.1%)		148 (16.6%)	
Other [†]	71 (11.3%)		110 (12.3%)	
Grade[‡]				0.208
Low	44 (7.0%)		58 (6.5%)	
High	584 (93.0%)		836 (93.5%)	
Stage				0.501
I	298 (47.5%)		411 (46.0%)	
II	143 (22.8%)		212 (23.7%)	
III	136 (21.6%)		200 (22.4%)	
IV	51 (8.1%)		71 (7.9%)	
Tumor size				0.726
<2 cm	142 (22.6%)		196 (21.9%)	
2-4 cm	121 (19.3%)		172 (19.2%)	
>4 cm	365 (58.1%)		526 (58.8%)	
Metastasis				0.625
Absent	455 (72.5%)		643 (71.9%)	
Present	173 (27.5%)		251 (28.1%)	
Surgery				0.714
Performed	293 (46.7%)		413 (46.2%)	
None	335 (53.3%)		481 (53.8%)	
Radiotherapy				0.653
Yes	247 (39.3%)		347 (38.8%)	
No	381 (60.7%)		547 (61.2%)	
Chemotherapy				0.917
Yes	406 (64.6%)		577 (64.5%)	
No	222 (35.4%)		317 (35.5%)	

[†]American Indian/Alaskan Native, Asian/Pacific Islander[‡]Low: Grade I (well differentiated) and Grade II (moderately differentiated); High: Grade III (poorly differentiated) and Grade IV (undifferentiated anaplastic).