

Lampela M, Minkinen K, Straková P, Bhuiyan R, He W, Mäkiranta P, Ojanen P, Penttilä T, Laiho R:
Effects of persistent water-table level drawdown on fine-root biomass and
production in different site types of a boreal forested peatland

Supplementary figures

Figure S1. Study site locations in the Lakkasuo area. The site types with their drained counterparts: 1-2) DP, dwarf-shrub pine bog, 3-4) TP, tall-sedge pine fen, 5-6) VS, *Vaccinium myrtillus* spruce swamp, 7-8) HS, herb-rich hardwood-spruce swamp. The three plots in each site are marked with red dots. The green ruling borders a nature conservation area. Drainage ditches are shown with straight blue lines. For general information on the area, see Laine et al. (2004).




Figure S2. Study sites: a) DP, dwarf-shrub pine bog, b) DPdr, drained dwarf-shrub pine bog, c) TP, tall-sedge pine fen, d) TPdr, e) VS, *Vaccinium myrtillus* spruce swamp, f) VSdr, g) HS, herb-rich hardwood-spruce swamp and h) HSdr. Photos: Kari Minkkinen





e) VS




f) VSdr





Figure S3. Cumulative soil temperature sums (d.d. >5°C) in 5 cm depth at the a) DP, DPdr, TP and TPdr sites and b) VS, VSdr, HS and HSdr sites, for the two growing seasons, 2016 and 2017, of the ingrowth core incubation period.

a)



b)




Figure S4. P, K, Ca and Mg concentration profiles measured from the sites and from the peat of some incubated ingrowth cores. Site data are from three pooled peat cores per site, and ingrowth core data are from 4-6 cores per peat type, incubated at different sites.

