

Global, Regional, and National Burden of Multiple Myeloma from 1990 to 2019: Estimates from the 2019 Global Burden of Disease study

Linghui Zhou^{1,2,3,4,6}, Qin Yu^{5,6}, Guoqing Wei^{1,2,3,4}, Linqin Wang^{1,2,3,4}, Yue Huang^{1,2,3,4}, Kejia Hu^{1,2,3,4}, Yongxian Hu^{1,2,3,4}, He Huang^{1,2,3,4}

1. Bone Marrow Transplantation Center, the First Affiliated Hospital, Zhejiang University School of Medicine;
2. Institute of Hematology, Zhejiang University;
3. Zhejiang Province Engineering Laboratory for Stem Cell and Immunity Therapy;
4. Zhejiang Laboratory for Systems & Precision Medicine, Zhejiang University Medical Center;
5. College of Life Science, Zhejiang Chinese Medical University, Hangzhou, China;
6. These authors contribute equally to this work.

Corresponding author:

Yongxian Hu, MD/PhD

Bone Marrow Transplantation Center, The First Affiliated Hospital, School of Medicine, Zhejiang University. No.79 Qingchun Road, Hangzhou, China.

Telephone and Fax: 86-0571-87236706 E-mail: 1313016@zju.edu.cn

He Huang, Professor/MD/PhD

Bone Marrow Transplantation Center, The First Affiliated Hospital, School of Medicine, Zhejiang University. No.79 Qingchun Road, Hangzhou, China.

Telephone and Fax: 86-0571-87236706 E-mail: huanghe@zju.edu.cn

Supp Figures

Supp Figure 1 The change trends of age-standardized incidence, death and DALYs rate among different SDI quintiles.

Supp Figure 2 The correlation between SDI and age-standardized rates and percentage change. A: ASIR; B: percentage change in incidence; C: ASDR; D: percentage change in death rate. The circles represent countries that were available on SDI data. The size of circle represents the number of multiple myeloma patients. The ρ indices Person correlation coefficient and p values were derived from Pearson correlation analysis. ASIR: age-standardized incidence rate; ASDR: age-standardized death; SDI, socio-demographic index.

Supp Figure 3 The percentage change of multiple myeloma in age-standardized rates worldwide in 2019. A: incidence; B: death rate

Supp Figure 4 The ratio of male to female ASIR among different age groups in 1990 and 2019. A: global; B: high SDI; C: high-middle SDI; D: middle SDI; E: middle-low SDI; F: low SDI. SDI, socio-demographic index. ASIR, age-standardized incidence rate


Supp Figure 5 The ASDR of multiple myeloma among regions based on SDI in 2019. ASDR: age standardized death rate (per 100,000 population).

Supp Figure 6 T The ratio of male to female ASDR among different age groups in 1990 and 2019. A: global; B: high SDI; C: high-middle SDI; D: middle SDI; E: middle-low SDI; F: low SDI. SDI, socio-demographic index. ASDR, age-standardized death rate


Supp Figure 7 The age-standardized DALYs rate of multiple myeloma among regions based on SDI in 2019.

Supp Figure 8 The percentage of age-standardized deaths and DALYs attributable to high body-mass index from 1990 to 2019.


Supp Figure 1 The change trends of age-standardized incidence, death and DALYs rate among different SDI quintiles.




Supp Figure 2 The correlation between SDI and age-standardized rates and percentage change. A: ASIR; B: percentage change in incidence; C: ASDR; D: percentage change in death rate. The circles represent countries that were available on SDI data. The size of circle represents the number of multiple myeloma patients. The ρ indices Person correlation coefficient and p values were derived from Pearson correlation analysis. ASIR: age-standardized incidence rate; ASDR: age-standardized death; SDI, socio-demographic index.




Supp Figure 3 The percentage change of multiple myeloma in age-standardized rates worldwide in 2019. A: incidence; B: death rate




Supp Figure 4 The ratio of male to female ASIR among different age groups in 1990 and 2019. A: global; B: high SDI; C: high-middle SDI; D: middle SDI; E: middle-low SDI; F: low SDI. SDI, socio-demographic index. ASIR, age-standardized incidence rate




Supp Figure 5 The ASDR of multiple myeloma among regions based on SDI in 2019. ASDR: age standardized death rate (per 100,000 population).



Supp Figure 6 T The ratio of male to female ASDR among different age groups in 1990 and 2019. A: global; B: high SDI; C: high-middle SDI; D: middle SDI; E: middle-low SDI; F: low SDI. SDI, socio-demographic index. ASDR, age-standardized death rate



Supp Figure 7 The age-standardized DALYs rate of multiple myeloma among regions based on SDI in 2019.



Supp Figure 8 The percentage of age-standardized deaths and DALYs attributable to high body-mass index from 1990 to 2019.

