

Characteristics of study participants

<i>Socio-demographics</i>	<i>Freq.</i>	<i>Percent (%)</i>	<i>Mean</i>	<i>SD</i>
<i>Age</i>				
20-24	17	45.95		
25-29	4	10.81		
30-34	9	24.32		
35-39	4	10.81		
40-44	2	5.41		
44+	1	2.70		
<i>Gender</i>				
<i>Female</i>	21	56.76		
<i>Male</i>	16	43.24		
<i>Religion</i>				
<i>Roman catholic</i>	19	51.35		
<i>Anglican</i>	10	27.03		
<i>Muslim</i>	1	2.70		
<i>Orthodox</i>	0	0		
<i>Pentecostal</i>	4	10.81		
<i>Traditional</i>	0	0		
<i>None</i>	3	8.11		
<i>Program</i>				
<i>BNC</i>	19	51.35		
<i>BNS</i>	18	48.65		
<i>Year of study</i>			2.73	± 0.99
1	3	8.11		
2	15	40.54		
3	8	21.62		
4	11	29.73		
<i>Tuition funding</i>				

<i>Government-funded</i>	3	8.11
<i>NGO-funded</i>	4	10.81
<i>Private funded</i>	14	37.84
<i>Self-funded</i>	16	43.24
<i>Marital status</i>		
<i>Single</i>	24	64.86
<i>Married</i>	13	35.14
<i>Semesters rotated on surgical ward</i>		
1	14	37.84
2	11	29.73
3	12	32.43
<i>Previous training on fluids and electrolytes follow-up for burns patients</i>		
<i>Yes</i>	20	45.95
<i>No</i>	17	54.05