

Table 1. Clinical scores of ferrets infected with SARS-CoV-2

Group		0dpi	2dpi	4dpi	6dpi	8dpi	10dpi
G1 (Juvenile)	Cough	0.00	0.00	0.00	0.00	0.00	0.00
	Runny nose	0.00	0.56±0.50	0.33±0.47	0.00	0.00	0.00
	Movement, activity	0.00	0.44±0.50	0.17±0.37	0.00	0.00	0.00
	Total	0.00	1.00	0.50	0.00	0.00	0.00
G2 (Young adult)	Cough	0.00	0.33±0.47	1.00	0.83±0.37	0.00	0.00
	Runny nose	0.00	0.22±0.42	1.33±0.47	1.17±0.37	0.00	0.00
	Movement, activity	0.00	0.67±0.67	1.83±0.37	1.83±0.37	1.00	0.00
	Total	0.00	1.22	4.17	3.83	1.00	0.00
G3 (Aged adult)	Cough	0.00	0.44±0.50	1.00	0.83±0.37	0.00	0.00
	Runny nose	0.00	0.44±0.68	1.50±0.50	1.50±0.50	0.67±0.47	0.33±0.47
	Movement, activity	0.00	0.78±0.92	2.00	1.83±0.37	1.00	0.67±0.47
	Total	0.00	1.67	4.50	4.00	1.75	1.00

Observational clinical symptoms: Cough, rhinorrhea, movement, and activity.

Score: 0; normal, 1: occasional, mild reduced activity, 2: frequent, reduced activity.

* Scores were measured by observation of clinical symptoms for at least 20 minutes in each group of ferrets based on the following criteria: Cough: 0; no evidence of cough, 1; occasional cough, 2; frequent cough (score 2).

Rhinorrhea: 0; no nasal rattling or sneezing, 1; moderate nasal discharge on external nares, 2; severe nasal discharge on external nares.

Movement, activity: 0; normal movement and activity, 1; mild reduced movement and activity, 2; evidence of reduced movement and activity.