


Pentosan polysulfate induces latent prion infection in Fukuoka-1 strain infected cell culture

Hanae Takatsuki, Morikazu Imamura, Tsuyoshi Mori, Ryuichiro Atarashi


Supplementary Information

Supplementary Figure 1




(A) The full length blot of the cropped image shown in Figure 4B. Two gels (placed at the top and bottom) were transferred to a single polyvinylidene difluoride membrane for Western blotting. **(B)** Merged images of marker imaging and luminescence images. The numbers represent the molecular weight of the marker.

Supplementary Figure 2




Inhibitory effect of PPS on PMCA seeded with FK1. (A) The full length blot of the cropped image shown in Figure 5A. Three gels (placed at the top, middle, and bottom) were transferred to a single polyvinylidene difluoride membrane for Western blotting. **(B)** Blot image of Experiment 2. **(C)** Blot image of Experiment 3.

Supplementary Figure 3




Inhibitory effect of PPS on PMCA seeded with 22L. (A) The full length blot of the cropped image shown in Figure 5A. Three gels (placed at the top, middle, and bottom) were transferred to a single polyvinylidene difluoride membrane for Western blotting. **(B)** Blot image of Experiment 2. **(C)** Blot image of Experiment 3.

Supplementary Figure 4




Inhibitory effect of PPS on PMCA seeded with Chandler. (A) The full length blot of the cropped image shown in Figure 5A. Three gels (placed at the top, middle, and bottom) were transferred to a single polyvinylidene difluoride membrane for Western blotting. **(B)** Blot image of Experiment 2. **(C)** Blot image of Experiment 3.

Supplementary Figure 5



Inhibitory effect of PPS on PMCA seeded with ME7. (A) The full length blot of the cropped image shown in Figure 5A. Three gels (placed at the top, middle, and bottom) gels were transferred to a single polyvinylidene difluoride membrane for Western blotting. **(B)** Blot image of Experiment 2. **(C)** Blot image of Experiment 3.

Supplementary Figure 6



Inhibitory effect of PPS on PMCA seeded with mBSE. (A) The full length blot of the cropped image shown in Figure 5A. Two gels (placed at the top and bottom) were transferred to a single polyvinylidene difluoride membrane for Western blotting. **(B)** Blot image of Experiment 2. **(C)** Blot image of Experiment 3.