

Table 2. Five- and 10-year RFS for ER+HER2- breast cancer patients primarily treated with surgery using the 8th AJCC TNM system (anatomic and prognostic stage group) and Bioscore staging system.

Five-year RFS								
Anatomic stage	n (%)	RFS (%)	Prognostic stage	n (%)	RFS (%)	Bioscore	n (%)	RFS (%)
I (A and B)	2,041 (48.3)	91.1	I (A and B)	2,821 (66.7)	91.8	0	0 (0)	NA
IIA	1,209 (28.6)	90.6	IIA	815 (19.3)	87.6	1	2,179 (51.5)	92.1
IIB	499 (11.8)	85.2	IIB	254 (6.0)	81.5	2	1,222 (28.9)	89.0
IIIA	323 (7.6)	82.0	IIIA	154 (3.6)	77.3	3	453 (10.7)	82.6
IIIC	157 (3.7)	65.0	IIIB	163 (3.9)	66.3	4	169 (4.0)	85.2
			IIIC	22 (0.5)	40.9	5	155 (3.7)	69.7
						6	51 (1.1)	54.9
						7	0 (0)	NA
Ten-year RFS								
Anatomic stage	n (%)	RFS (%)	Prognostic stage	n (%)	RFS (%)	Bioscore	n (%)	RFS (%)
I (A and B)	2,041 (48.3)	83.8	I (A and B)	2,821 (66.7)	84.6	0	0 (0)	NA
IIA	1,209 (28.6)	84.9	IIA	815 (19.3)	80.2	1	2,179 (51.5)	85.1
IIB	499 (11.8)	75.2	IIB	254 (6.0)	70.9	2	1,222 (28.9)	81.5
IIIA	323 (7.6)	68.4	IIIA	154 (3.6)	62.3	3	453 (10.7)	71.5
IIIC	157 (3.7)	52.9	IIIB	163 (3.9)	52.1	4	169 (4.0)	75.1
			IIIC	22 (0.5)	31.8	5	155 (3.7)	52.3
						6	51 (1.1)	47.1
						7	0 (0)	NA

