

Table 3.

Diagnosis at discharge from the ED over the study period (March 9th-May 9th) comparing 2019 and 2020.

	Study period			
	2019 (n=6568)	2020 (n=1133)	Change (95%CI%)	OR (95%CI)
Respiratory	1409 (21.5)	165 (14.6)	-88.3 (86.6 – 90.0)	0.68 (0.58 – 0.80)*
Gastrointestinal	961 (14.6)	107 (9.4)	-88.9 (86.8 – 90.9)	0.65 (0.53 – 0.79)*
Trauma	878 (13.4)	157 (13.4)	-82.1 (79.5 – 84.7)	1.04 (0.87 – 1.23)
Otorhinolaryngology	688 (10.5)	78 (6.9)	-88.7 (86.2 – 91.1)	0.66 (0.52 – 0.83)*
Fever/Viral infection	466 (7.1)	93 (8.2)	-80.0 (76.3 – 83.8)	1.16 (0.93 – 1.46)
Rash/Dermatology	213 (3.2)	31 (2.7)	-85.4 (80.5 – 90.4)	0.84 (0.58 – 1.23)
Seizures/Neurology	151 (2.3)	42 (3.7)	-72.2 (64.7 – 79.7)	1.61 (1.15 – 2.27)
Ophthalmology	74 (1.1)	7 (0.6)	-90.5 (83.2 – 97.9)	0.55 (0.25 – 1.19)
Nephrology/Urology	72 (1.1)	30 (2.7)	-58.3 (46.3 – 70.4)	3.14 (2.13 – 4.64)*
Surgery	56 (0.9)	38 (3.4)	-32.1 (19.0 – 45.3)	3.93 (2.61 – 5.94)*
Cardiovascular	56 (0.9)	20 (1.8)	-64.3 (50.8 – 77.7)	2.07 (1.24 – 3.35)*
Psychiatric emergency	37 (0.6)	11 (1.0)	-70.3 (54.2 – 86.3)	1.72 (0.88 – 3.38)

*Values are expressed as numbers (%). *p-value<0.05.*

Abbreviation. CI: Confidence interval. OR: Odds ratio.