

Table 2 Pathological results in unmatched and matched entire cohort

	Entire unmatched cohort			Propensity score-matched cohort		
	LN yield $\geq$ 15 (<i>n</i> =174)	LN yield<15 (<i>n</i> =204)	<i>p</i>	LN yield $\geq$ 15 (<i>n</i> =117)	LN yield<15 (<i>n</i> =117)	<i>p</i>
Pathological T-stage, <i>n</i> (%)			0.070			0.340
T0	2 (1.1)	4 (2.0)		2 (1.7)	0 (0.0)	
T2a-c	90 (61.7)	123 (60.3)		62 (53.0)	65 (55.6)	
T3a-b	82 (47.1)	74 (36.3)		53 (45.3)	50 (42.7)	
T4	0 (0.0)	3 (1.5)		0 (0.0)	2 (1.7)	
ISUP grade group*, <i>n</i> (%)			0.770			0.261
NA	2 (1.1)	4 (2.0)		2 (1.7)	0 (0.0)	
1-2	22 (12.7)	34 (16.7)		10 (8.6)	20 (17.1)	
3	58 (33.3)	60 (29.4)		38 (32.5)	39 (33.3)	
4	48 (27.6)	49 (24.0)		34 (29.1)	29 (24.8)	
5	44 (25.3)	57 (27.9)		33 (28.2)	29 (24.8)	
Mean LN yield, <i>n</i> (SD)	23.9 (7.5)	3.4 (4.1)	<0.001	24.4 (8.1)	3.6 (4.4)	<0.001
LN metastases, <i>n</i> (%)	35 (20.1)	4 (2.0)	<0.001	20 (17.1)	3 (2.6)	<0.001
Positive resection margin, <i>n</i> (%)	61 (35.1)	50 (24.5)	0.031	39 (33.3)	34 (29.1)	0.573

LN=lymph node; NA=not available; SD=standard deviation; ISUP=International Society of Urological Pathology