

**Supplementary Table 2.** Total overlapped areas in km<sup>2</sup> between lions' (vertical column) and spotted hyenas' (horizontal column) home ranges and core use areas in the (a) Etosha National Park, Namibia; (b) Chobe National Park and Linyanti Conservancy, Botswana. Utilization distributions were generated with the home range (95%) and core use area (50%) *a*-LoCoH (adaptive Local Convex Hull) isopleths. Males are underlined. An asterisk denotes mortality.

(a)

HOME RANGE (95%)

DRY SEASON

WET SEASON

SPOTTED HYENA

GO-33869

TJ-33870

NE-33871

SA-33872

AU-33873

OM-33874

SU-33951

WO-34310

GO-33869

TJ-33870\*

NE-33871

SA-33872

AU-33873\*

OM-33874

SU-33951\*

WO-34310

LION

OK-33863

RE-33864

NU-33865

MO-33866

OJ-33867

SU-33868

OM-34308\*

LU-34308

OF-34309\*

G2-35678

0

38.8

0.1

62

0

72

0

0

0

0

48.7

0

0

0

27.1

0.9

57

58

0

0

43.8

236

0

0

98

260

0.2

43.8

0.7

-

-

124

-

0

4.4

124

100

25.2

-

-

CORE AREA (50%)

DRY SEASON

WET SEASON

SPOTTED HYENA

GO-33869

TJ-33870

NE-33871

SA-33872

AU-33873

OM-33874

SU-33951

WO-34310

GO-33869

TJ-33870\*

NE-33871

SA-33872

AU-33873\*

OM-33874

SU-33951\*

WO-34310

LION

OK-33863

RE-33864

NU-33865

MO-33866

OJ-33867

SU-33868

OM-34308\*

LU-34308

OF-34309\*

G2-35678

0

0.2

0

0

0

21.5

0

0

0

0

4.2

0

0

0

0

0

0.2

0.1

0

0

0

56

0

0

69

0

0

9.7

0

-

-

26.4

-

0

0

40.7

16.0

0.3

-

-

