

Primary care physicians' perceptions concerning engagement in cancer survivor care

Journal name: Supportive Care in Cancer

Miho Kimachi¹, Kenji Omae^{1,2}, Tsukasa Kamitani¹, Shingo Fukuma⁴

¹Department of Healthcare Epidemiology, School of Public Health in the Graduate School of Medicine, Kyoto University, Kyoto, Kyoto, Japan

²Department of Innovative Research and Education for Clinicians and Trainees (DiRECT), Fukushima Medical University, Fukushima, Fukushima, Japan

³Section of Clinical Epidemiology, Department of Community Medicine, Graduate School of Medicine, Kyoto University, Kyoto, Japan

⁴Human Health Sciences, Kyoto University Graduate School of Medicine, Kyoto, Kyoto, Japan

***Corresponding author:** Miho Kimachi, MD, PhD E-mail: kimachi.miho.5c@kyoto-u.jp

Supplemental Table 3. Available cancer survivor care in facilities to which primary care physicians belong when oncologists request cooperation (n=87)

Cancer follow-up care	Available at any time	Available only after cancer therapy	Not available at any time
Regular cleaning of central venous access port device	63 (72.4)	17 (19.5)	7 (8.1)
Prescription of regular medications other than anticancer agents	83 (95.4)	4 (4.6)	0 (0)
Therapy for infection	63 (72.4)	24 (27.6)	0 (0)
Prescription of opioid analgesic medications	76 (87.4)	8 (9.2)	3 (3.5)
Prescription of analgesic medications other than opioids	80 (93.0)	6 (7.0)	0 (0)
Nutritional education	73 (83.9)	14 (16.1)	0 (0)
Stoma care	62 (71.3)	21 (24.1)	4 (4.6)
Maintenance or recovery support of patient's ADL	74 (85.1)	10 (11.5)	3 (3.5)
Support for patients' social reintegration or return to work	70 (80.5)	13 (14.9)	4 (4.6)
Psychological support	78 (90.7)	5 (5.8)	3 (3.5)
Regular screening for recurrence of treated cancer	32 (36.8)	42 (48.3)	13 (14.9)
Screening for occurrence of cancers other than the treated cancer	59 (68.6)	16 (18.6)	11 (12.8)

ADL, activities of daily living

Summarized as n (%)