

Supplementary information

A critical role of calcineurin in stress responses, hyphal formation, and virulence of the pathogenic fungus *Trichosporon asahii*

Yasuhiko Matsumoto^{1*}, Asami Yoshikawa¹, Tae Nagamachi¹, Yu Sugiyama¹, Tsuyoshi Yamada^{2,3}, and Takashi Sugita¹

¹Department of Microbiology, Meiji Pharmaceutical University, 2-522-1, Noshio, Kiyose, Tokyo 204-8588, Japan.

²Teikyo University Institute of Medical Mycology, 359 Otsuka, Hachioji, Tokyo 192-0395, Japan.

³Asia International Institute of Infectious Disease Control, Teikyo University, 2-11-1, Kaga, Itabashi-ku, Tokyo, 173-8605, Japan

*Address correspondence to: Dr. Yasuhiko Matsumoto, Department of Microbiology, Meiji Pharmaceutical University, 2-522-1, Noshio, Kiyose, Tokyo 204-8588, Japan, Tel: +81-42-495-8745, e-mail: ymatsumoto@my-pharm.ac.jp.




Figure 1D




Figure 1H

Supplementary Figure 1 Full-length blots of Figure 1d and 1h. Marker 1: OneSTEP Marker 1 (λ /HindIII digest) (Nippon Gene Co., Ltd., Tokyo, Japan), Marker 2: Gene Ladder 100 (Nippon Gene Co., Ltd., Tokyo, Japan)