


Additional file 5: Sensitivity analyses applying the original WHO-MCS-A severity classification(1,2).

Figure additional file 5: Sensitivity analysis - proportion of women by severity categories using original WHO-MCS-A classification*



* WHO-MCS-A: World Health Organization Multi-Country Study on Abortion(2)

AMoCo study_Severity-Magnitude

Table additional file 5: Proportion of women with abortion complications by severity category in AMoCo Nigeria and CAR study hospitals compared to African hospitals in WHO-MCS-A* using original WHO-MCS-A* classification

	Original WHO-MCS-A* criteria								
	Nigeria AMoCo (N=520)			CAR AMoCo (N=548)			210 African hospitals (11 countries) WHO-MCS-A* (N=13657)		
	Proportion	95%CI		Proportion	95%CI		Proportion	95%CI	
Mild	25.6%	21.9%	29.6%	32.3%	28.4%	36.4%	32.4%	31.6%	33.2%
Moderate	7.5%	5.4%	10.1%	17.0%	13.9%	20.4%	58.2%	57.4%	59.0%
PLTC	62.3%	58.0%	66.5%	46.9%	42.7%	51.2%	7.0%	6.6%	7.4%
SMO	4.6%	3.0%	6.8%	3.8%	2.4%	5.8%	2.4%	2.1%	2.6%
Severe complications (PLTC + SMO)	66.9%	62.7%	71.0%	50.7%	46.5%	55.0%	9.4%	8.9%	9.9%

* WHO-MCS-A: World Health Organization Multi-Country Study on Abortion using original WHO near-miss criteria(2)

PLTC: Potentially Life-Threatening Complications

SMO: Severe Maternal Outcome (= near-miss cases + deaths)

References:

1. Kim CR, Tunçalp Ö, Ganatra B, Gülmezoglu AM. WHO Multi-Country Survey on Abortion-related Morbidity and Mortality in Health Facilities: study protocol. BMJ Glob Heal [Internet]. 2016;1(3):e000113. Available from: <http://gh.bmj.com/lookup/doi/10.1136/bmjgh-2016-000113>
2. Qureshi Z, Mehrtash H, Kouanda S, Griffin S, Filippi V, Govule P, et al. Understanding abortion-related complications in health facilities: results from WHO multicountry survey on abortion (MCS-A) across 11 sub-Saharan African countries. BMJ Glob Heal [Internet]. 2021 Jan 29 [cited 2021 Feb 1];6(1):e003702. Available from: <https://gh.bmj.com/lookup/doi/10.1136/bmjgh-2020-003702>