

Supplemental table 1.

Supplemental table 1. Subtypes of progressive hemorrhagic injury (PHI) and their outcomes.

Characteristics	Category	PHI N = 80 n (%)	Non-PHI N = 124 n (%)	p-value	pIPCH N = 33 n (%)	Non-pIPCH N = 55 n (%)	p-value	pEDH N = 15 n (%)	Non-pEDH N = 25 n (%)	p-value
Age	Mean ±	48.49 ±	48.52 ±	.988	53.00 ±	52.78 ±	.948	38.67 ±	42.92 ±	.334
	SD	18.06	15.31		16.35	14.26		13.52	13.16	
Motor score	1	3 (3.8)	3 (2.4)		0 (0.0)	1 (1.8)		0 (0.0)	1 (4.0)	
	2	3 (3.8)	5 (4.0)		0 (0.0)	1 (1.8)		0 (0.0)	1 (4.0)	
	3	2 (2.5)	4 (3.2)		3 (9.1)	1 (1.8)		1 (6.7)	1 (4.0)	
	4	9 (11.3)	8 (6.5)		4 (12.1)	3 (5.5)		0 (0.0)	0 (0.0)	
	5	23 (28.8)	24 (19.4)		10 (30.3)	11 (20.0)		7 (46.7)	7 (28.0)	

	6	40 (50.0)	80 (64.5)	.376	16 (48.5)	38 (69.1)	.214	7 (46.7)	15 (60.0)	.632
Pupil	Both	67 (83.8)	110 (88.7)		28 (84.8)	50 (90.9)		13	22 (88.0)	
	reactive							(86.7)		
	one	5 (6.3)	5 (4.0)		1 (3.0)	2 (3.6)		2 (13.3)	2 (8.0)	
	reactive									
	Non-	8 (10.0)	9 (7.3)	.587	4 (12.1)	3 (5.5)	.533	0 (0.0)	1 (4.0)	.648
	reactive									
EDH	Yes	27 (33.8)	36 (29.0)	.476	-	-	-	15	25	-
Lesion	Mean ±	22.91 ±	17.93 ±	.183	14.60 ±	9.81 ± 15.37	.159	26.01 ±	22.73 ±	.760
volume	SD	24.40	26.97		15.30			36.61	30.21	
Midline	≥5 mm	13 (16.3)	14 (11.3)	.307	4 (12.1)	2 (3.6)	.138*	2 (13.3)	4 (16)	.600*
shift										
Cistern	normal	56 (70.0)	95 (76.6)		25 (75.8)	46 (83.6)		12 (80.0)	20 (80.0)	

compressi

on

compresses	16 (20.0)	21 (16.9)		5 (15.2)	8 (14.5)		3 (20.0)	3 (12.0)
------------	-----------	-----------	--	----------	----------	--	----------	----------

d

absent	8 (10.0)	8 (6.5)	.517	3 (9.1)	1 (1.8)	.278	0 (0.0)	2 (8.0)	.449
--------	----------	---------	------	---------	---------	------	---------	---------	------

*Fisher Exact test. pEDH: progressive epidural hematoma, pIPCH: progressive intraparenchymal contusion or hematoma, LOS: length of hospital stay, SD: standard deviation.