

Supplementary Table S2. Comparison between polypoidal choroidal vasculopathy (PCV) and typical neovascular age-related macular degeneration (nAMD)

	Typical type 1 nAMD	PCV	P value
	(n: 50)	(n: 44)	
Sex (M/F)	33/17	32/12	0.481
Age	71.2±6.4	66.2±8.9	0.002
Refractive error (diopter)	+0.36±1.37	+0.24±1.39	0.685
Total anti-VEGF injections	6.8±3.7	7.3±3.8	0.505
BCVA (logMAR)			
baseline	0.47±0.29	0.49±0.25	0.671
after 3 loading doses	0.36±0.26	0.31±0.24	0.333
at 1-year	0.33±0.27	0.31±0.25	0.607
ONL thickness (µm)			
baseline	52.0±20.6	57.0±18.3	0.213
after 3 loading doses	63.6±26.2	66.3±21.7	0.594
at 1-year	61.9±24.0	64.5±24.8	0.618
SRF height (µm)			
Baseline	157.7±88.9	191.1±109.9	0.107
after 3 loading doses	27.2±38.5	28.9±47.4	0.850
at 1-year	46.9±67.0	67.9±89.2	0.197
Choroidal thickness (µm)			
Baseline	199.1±56.9	343.4±83.4	<0.001
after 3 loading doses	179.9±59.1	316.9±82.1	<0.001
at 1-year	178.6±66.7	288.5±73.3	<0.001

ONL, outer nuclear layer; SRF, subretinal fluid; BCVA, best-corrected visual acuity; logMAR, log of minimum angle of resolution