

Pedigree reconstruction and population structure using SNP markers in Gir cattle

Arielly Oliveira Garcia¹, Pamela Itajara Otto², Luiz Afonso Glatzl Junior³, Renata de Fátima Bretanha Rocha¹, Mateus Guimarães dos Santos¹, Daniele Alves de Oliveira², Marcos Vinicius Gualberto Barbosa da Silva⁴, João Cláudio do Carmo Panetto⁴, Marco Antônio Machado⁴, Rui da Silva Verneque⁴, and Simone Eliza Facioni Guimarães^{1*}

¹Department of Animal Science, Universidade Federal de Viçosa, Viçosa, MG, 36570-900, Brazil

²Department of Animal Science, Universidade Federal de Santa Maria, Santa Maria, RS, 97105-900, Brazil

³Department of Computer Science, Universidade Federal de Juiz de Fora, Juiz de Fora, MG, 36036-900, Brazil

⁴EMBRAPA - Dairy Cattle Research Center, Juiz de Fora, MG, 36038-330, Brazil

* Corresponding author: sfacioni@ufv.br

Journal of Applied Genetics

Online Resource 2 Distribution of 1,810 SPNs common in all four different chips (GeneSeek Genomic Profiler Indicus (GGPi) with 35K and 50K, GGPi-LD BeadChip, and Bovine HD) along the 29 autosomal chromosomes

