

Pedigree reconstruction and population structure using SNP markers in Gir cattle

Arielly Oliveira Garcia¹, Pamela Itajara Otto², Luiz Afonso Glatzl Junior³, Renata de Fátima Bretanha Rocha¹, Mateus Guimarães dos Santos¹, Daniele Alves de Oliveira², Marcos Vinicius Gualberto Barbosa da Silva⁴, João Cláudio do Carmo Panetto⁴, Marco Antônio Machado⁴, Rui da Silva Verneque⁴, and Simone Eliza Facioni Guimarães^{1*}

¹Department of Animal Science, Universidade Federal de Viçosa, Viçosa, MG, 36570-900, Brazil

²Department of Animal Science, Universidade Federal de Santa Maria, Santa Maria, RS, 97105-900, Brazil

³Department of Computer Science, Universidade Federal de Juiz de Fora, Juiz de Fora, MG, 36036-900, Brazil

⁴EMBRAPA - Dairy Cattle Research Center, Juiz de Fora, MG, 36038-330, Brazil

* Corresponding author: sfacioni@ufv.br

Journal of Applied Genetics

Online Resource 5 Distribution and mean length of runs of homozygosity (ROH) in Mb detected throughout the autosomal genome in Gir cattle. Bar chart shows ROH counts and the dark gray line shows the average ROH size (Mb) per chromosome

