

Intestinal epithelial barrier integrity investigated by label-free techniques in ulcerative colitis patients

Elsie Quansah^{+1,2}, Elena Gardey^{+3,4}, Anuradha Ramoji^{+,1,2,5*}, Tobias Meyer-Zedler^{1,2}, Bianca Goehrig⁶, Astrid Heutelbeck⁶, Stephanie Hoeppener^{4,7}, Michael Schmitt^{1,2}, Maximilian Waldner⁸, Andreas Stallmach^{3,4} and Jürgen Popp^{1,2}.

¹Friedrich Schiller University Jena, Institute of Physical Chemistry and Abbe Center of Photonics (IPC), Helmholtzweg 4, D-07743 Jena, Germany

²Leibniz Institute of Photonic Technology (IPHT), Member of Leibniz Health Technologies, Albert-Einstein-Straße 9, D-07745 Jena, Germany

³Jena University Hospital, Department of Internal Medicine IV (Gastroenterology, Hepatology, Infectious Diseases and Interdisciplinary Endoscopy), Friedrich Schiller University Jena, Am Klinikum 1, D-07747 Jena, Germany

⁴Friedrich Schiller University Jena, Jena Center for Soft Matter (JCSM), Philosophenweg 7, D-07743 Jena, Germany

⁵Jena University Hospital, Center for Sepsis Control and Care (CSCC), Friedrich Schiller University Jena, Erlanger Allee 101, D-07747 Jena, Germany

⁶Jena University Hospital, Institute for occupational, social, and environmental medicine, Am Klinikum 1, D-07747, Jena, Germany

⁷Friedrich Schiller University Jena, Laboratory of Organic and Macromolecular Chemistry (IOMC), Humboldtstraße 10, D-07743 Jena, Germany

⁸Department of Medicine, University of Erlangen-Nuremberg, Erlangen, Germany.

⁺*Equal contributions*

**Corresponding author* anuradha.ramoji@uni-jena.de, Tel. Nr. 03641 9390926

Supporting Information




Figure S1. Immunofluorescence images of healthy colonic mucosa (A) and inflamed colonic mucosa (B). Intact epithelial barriers were identified in the healthy colonic mucosa (green arrows). The loss of barrier integrity is indicated by pink arrows. The immunofluorescence staining is shown in blue (DAPI): nuclei, red (E-cadherin): epithelium, and yellow (CD11b): immune cells.