

Supplementary Information

Neurotoxicity Research

Caffeine and nicotine in a Parkinson's disease's animal model: from olfactory improvement to neuroprotection.

Lais Soares Rodrigues^{1,2}, Adriano D.S. Targa^{1,2,3,4}, Ana C. D. Nosedá^{1,2}, Jessica L. Ilkiw¹, Juliane Fagotti^{1,5}, Luana C. Kmita^{1,6}, Patricia Dominico dos Santos¹, Flavia W.C. Dorieux¹, Priscilla L. S. Silveira¹, Valentín Cóppola Segovia^{7,8}, Silvio Marques Zanata⁷ and Marcelo M. S.Lima^{1,2}.

¹ Neurophysiology Laboratory, Department of Physiology, Federal University of Paraná, Curitiba, Paraná, Brazil.

² Department of Pharmacology, Federal University of Paraná, Curitiba, Paraná, Brazil.

³ Hospital Universitari Arnau de Vilanova-Santa Maria, IRBLleida, Translational Research in Respiratory Medicine, Lleida, Spain.

⁴ Centro de Investigación Biomédica en Red de Enfermedades Respiratorias (CIBERES), Madrid, Spain.

⁵ Department of Neuroscience, Physiology and Pharmacology, Medical Sciences Building, Bloomsbury Campus, University College London, Kings Cross, London.

⁶ Programa de Tecnologia em Saúde, Pontifícia Universidade Católica do Paraná, Curitiba, Paraná, Brazil.

⁷ Neurobiology Laboratory, Department of Basic Pathology, Federal University of Paraná, Curitiba, Paraná, Brazil

⁸ Laboratory of Molecular and Cellular Neurobiology, Department of Pharmacology and Chemical Biology and Center for Neurodegenerative Disease, Emory University School of Medicine, Atlanta, GA, USA.

CORRESPONDING AUTHOR:


MARCELO M. S. LIMA, PhD
Universidade Federal do Paraná
Setor de Ciências Biológicas,
Departamento de Fisiologia
Av. Francisco H. dos Santos s/n
ZIP: 81.531 – 990

Caixa Postal: 19031

Curitiba – Paraná – Brasil

Phone number: 0055-041-3361 1754

E-mail: mmslima@ufpr.br or marcelomslima.neuro@gmail.com



Online Resource 5 Percentage of TH-ir neurons in the glomerular layer within the OB of male rats (number of TH-ir neurons – mm²). **a.** Representative images of dopaminergic neurons within the OB (magnification 50x). **b.** Corresponds to OB neuronal counting analysis. The bars are represented as mean $\pm$ standard error of the mean, N=5-10/group, *P $\leq$ 0.05, **P $\leq$ 0.01. . Two-way ANOVA followed by Bonferroni post hoc test.