

Selective Synthesis of Functionalised Linear Aliphatic Primary Amines Using Photoredox Catalysis

Robin Cauwenbergh^{a,b}, Prakash Kumar Sahoo^{a,b}, Rakesh Maiti^{a,b}, Andrea Guidetti^c and Shoubhik Das^{a*}

^aOrganic Synthesis Division, Department of Chemistry, University of Antwerp, 2020 Antwerp, Belgium.

^bBIMEF Laboratory, Department of Chemistry, University of Antwerp, 2610 Antwerp, Belgium.

^cR.C., P.K.S., and R.M. contributed equally.

E-mail: shoubhik.das@uantwerpen.be

Table of Contents

General considerations	3
Setup for photocatalytic reactions	3
Standard procedures for photocatalytic hydroaminoalkylation	4
Condition A	4
Condition B	4
Condition C	5
Reaction Optimization	5
Optimization using boc-glycine	5
Optimization of free amines	7
Optimization of long chain amines	10
Stern-Volmer experiments.....	12
Synthetic procedures literature protocols	14
Procedure D	14
Procedure E	14
Synthesis of hydroaminoalkylated products	15
Copies of the NMR spectra	30
References	95

1. General considerations

Chemicals and solvents were obtained from commercial suppliers and used without further purification. Non-commercial starting materials were prepared as described below according to literature procedures. All reactions were carried out under Nitrogen atmosphere using Schlenk techniques.

Thin Layer Chromatography (TLC) was performed using TLC plates from Merck (SiO₂, Kieselgel 60 F254 neutral, on aluminum with fluorescence indicator) and compounds were visualized by UV detection (254 nm). Flash column chromatographic purification of products was accomplished using an automated chromatography system with on-line UV and ELSD detection using Grace® Silica flash Cartridges.

Gas chromatography-mass spectrometry (GC-MS) samples were prepared by dissolving 0.1-5 mg of the compound in acetone (GC-MS quality) and further diluted to a concentration of 10⁻⁵ - 10⁻⁶ M. The samples were subsequently filtered using a CHROMAFIL® PET-20/25 syringe filter and 3 µL were injected. The apparatus was an Agilent Technologies 7890 A GC System coupled to an Agilent Technologies 5975 C inert MSD with triple-axis detector. As column an Optima 725820.30 30 m × 250 µm × 0.25 µm was selected. Carrier gas was helium. Inlet temperature heater: 225 °C. Oven program: 70 °C for 3 min, then heating 5 °C min⁻¹ to 160 °C and heating at 160 °C for 2 min. A one-minute post-run at 280 °C ends the oven program.

Nuclear Magnetic Resonance (NMR) spectra were recorded on a Bruker Avance III 400 (101 MHz for ¹³C) Fourier Transform NMR spectrometer at 300 K, using the non or partly deuterated solvent as internal standard (¹H: δ = 7.26 ppm, ¹³C: δ = 77.16 ppm for CDCl₃; ¹H: δ = 2.50 ppm, ¹³C: δ = 39.52 ppm for DMSO-d₆). Chemical shifts (δ) are given in ppm and coupling constants (*J*) are reported in Hertz (Hz). Multiplicities are described as s (singlet), d (doublet), t (triplet), q (quartet), p (pentet), br s (broad singlet) and m (multiplet) or combinations thereof. ¹³C NMR spectra were recorded with complete proton decoupling. ¹H NMR yields were determined by addition of a known amount of an internal standard and dissolving everything in a suitable deuterated solvent, followed by ¹H NMR analysis.

High resolution mass spectrometry (HRMS) samples were prepared by dissolving 0.1-5 mg of the compound in DMSO or CH₃CN/H₂O and further diluted to a concentration of 10⁻⁵ - 10⁻⁶ M. Formic acid (0.1%) was added prior to injection. 10 µL of each sample was injected using the CapLC system (Waters, Manchester, UK) and electrosprayed using a standard electrospray source. Samples were injected with an interval of 3 minutes. Positive ion mode accurate mass spectra were acquired using a Q-TOF II instrument (Waters, Manchester, UK). The MS was calibrated prior to use with a 0.1% H₃PO₄ solution. The spectra were lock mass corrected using the known mass of the nearest H₃PO₄ cluster or a known background ion. Analytes were detected as protonated or as a sodium adduct. All measured masses are within a difference of 5 ppm compared to the calculated mass unless specified otherwise.

2. Setup for photocatalytic reactions

The reaction setup is depicted in Figure S1. The reaction setup consists of a 456 nm Kessil lamp, cooling of the setup was performed by two commercially available 120 mm computer fans to keep the temperature around 30 °C. The light intensity was measured by Ophir StarLite power meter with 3A probe head. The light intensity of Kessil lamp (λ = 456 nm) was 0.34 W/cm² and the distance between tube and light was maintained between 3 to 4 cm.




Figure S1: Photocatalytic set up for the hydroaminoalkylation reactions.

3. Standard procedures for photocatalytic hydroaminoalkylation

3.1. Condition A

An oven dried 20 mL Schlenk Tube was charged with olefin (if solid, 0.2 mmol), 4CzIPN (0.02 equiv.), NaOH (1.0 equiv.), protected amino acid (1.0 equiv.) and a magnetic stirring bar. The reaction mixture was degassed by three freeze/pump/thaw cycles (3 min) using Schlenck techniques and refilled with nitrogen. After, olefin (if liquid, 0.2 mmol) and dry DMSO (1 ml) were added. After, the reaction was allowed to stir under 456 nm blue KESSIL light at room temperature. The mixture was monitored via TLC, quenched with water and extracted with EtOAc (3 times). Lastly, the solvent was removed under reduced pressure. NMR yield was determined by adding 1,3,5-trimethoxybenzene as the internal standard. The pure compound was obtained using Flash column chromatography using heptane:EtOAc (0-50 %) as the eluent.

3.2. Condition B

An oven-dried 20 mL Schlenk Tube was charged with olefin (if solid, 0.2 mmol), 4CzIPN (0.02 equiv.), NaOH (1equiv.), boc-anhydride (1.5 equiv.), unprotected amino acid (1.5 equiv.) and a magnetic stirring bar. The reaction mixture was degassed by three freeze/pump/thaw cycles (3 min) and refilled with nitrogen. After, olefin (if liquid, 0.2 mmol) and dry DMSO (2 ml) were added. After, the reaction was allowed to stir in the dark for 4 hours and subsequently under 456 nm blue KESSIL light at room temperature. The mixture was monitored via TLC, quenched with water and extracted with EtOAc (3 times). Lastly, the solvent was removed under reduced pressure. NMR yield was determined by adding 1,3,5-trimethoxybenzene as the internal standard. The pure compound was obtained using Flash column chromatography using heptane:EtOAc (0-50 %) as the eluent.


3.3. Condition C

An oven dried 20 mL Schlenk Tube was charged with 4CzIPN (0.02 equiv.), NaOH (1.0 equiv.), protected amino acid (1.0 equiv.) and a magnetic stirring bar. The reaction mixture was degassed by three freeze/pump/thaw cycles (3 min) using Schlenck techniques and refilled with nitrogen. After, olefin (0.2 mmol) and dry DMSO (2 ml) were added. Next, the reaction was allowed to stir under 456 nm blue KESSIL light at 70 °C. The mixture was monitored via TLC, quenched with water and extracted with EtOAc (3 times). Lastly, the solvent was removed under reduced pressure. NMR yield was determined by adding 1,3,5-trimethoxybenzene as the internal standard. The pure compound was obtained using Flash column chromatography using heptane:EtOAc (0-50 %) as the eluent.

4. Reaction Optimization

4.1. Optimization using boc-glycine

Table S1: Screening of different catalysts^a




Entry	Photocatalyst (PC)	Yield (%) ^b
1	Ir[dF(CF ₃)ppy] ₂ (dtbpy))PF ₆	93
2	DPZ	12
3	Mes-Acr	0
4	Fluorenone	0
5	4CzIPN	98
6	4CzIPN (4 mol%)	91
7	4CzIPN (3 mol%)	95
8	4CzIPN (2 mol%)	94
9	4CzIPN (1 mol%)	65
10	No PC	0
11	No light	0

^aReaction conditions: **1** (0.2 mmol), **2** (0.2 mmol), Cs₂CO₃ (0.2 mmol) and PC (5 mol%, unless stated otherwise) in DMSO (2mL) under N₂ atmosphere and light irradiation (456 nm 40 W KESSIL) at room temperature for 24h. ^bYield determined by ¹H NMR using 1,3,5-trimethoxybenzene as internal standard.

Table S2: Screening of different solvents^a

Entry	Solvent	Yield (%) ^b
1	DMSO	94
2	DMA	95
3	DMF	95
4	MeCN	61
5	Toluene	0
6	CHCl ₃	0
7	Dioxane	83
8	MeOH	0
9	DCM	0
10	THF	86
11	2-Me THF	35
12	DMSO (1 mL)	94

^aReaction conditions: **1** (0.2 mmol), **2** (0.2 mmol), Cs₂CO₃ (0.2 mmol) and 4CzIPN (2 mol%) in solvent (2 mL, unless stated otherwise) under N₂ atmosphere and light irradiation (456 nm 40 W KESSIL) at room temperature for 24h. ^bYield determined by ¹H NMR using 1,3,5-trimethoxybenzene as internal standard.


Table S3: Screening of different bases^a


Entry	Base	Yield (%) ^b
1	CS ₂ CO ₃	94
2	Li ₂ CO ₃	93
3	Na ₂ CO ₃	80
4	K ₂ CO ₃	66
5	KOAc	56
6	NaOH	95
7	K ₃ PO ₄	92
8	DIPEA	0
9	No base	0

^aReaction conditions: **1** (0.2 mmol), **2** (0.2 mmol), Base (0.2 mmol) and 4CzIPN (2 mol%) in DMSO (1 mL) under N₂ atmosphere and light irradiation (456 nm 40 W KESSIL) at room temperature for 24h.


^bYield determined by ¹H NMR using 1,3,5-trimethoxybenzene as internal standard.

4.2. Optimization of free amines

Table S4: Screening of base amount^a


Entry	Amount of NaOH (mol%)	Yield (%) ^b
1	0	81
2	5	83
3	10	82
4	20	95
5	50	88
6	100	90


^aReaction conditions: **1** (0.2 mmol), **2** (0.3 mmol), Boc-anhydride (0.3 mmol), NaOH and 4CzIPN (5 mol%) in DMSO (2 mL) under N₂ atmosphere in the dark for 4h and subsequently under light irradiation (456 nm 40 W KESSIL) at room temperature for 24h. ^bYield determined by ¹H NMR using 1,3,5-trimethoxybenzene as internal standard.

Table S5: Screening of catalyst loading^a

Entry	Amount of 4CzIPN	Yield (%) ^b
1	1 mol%	72
2	2 mol%	88
3	3 mol%	89
4	4 mol%	93
5	5 mol%	93

^aReaction conditions: **1** (0.2 mmol), **2** (0.3 mmol), Boc-anhydride (0.3 mmol), NaOH (20 mol%) and 4CzIPN in DMSO (2 mL) under N₂ atmosphere in the dark for 4h and subsequently under light irradiation (456 nm 40 W KESSIL) at room temperature for 24h. ^bYield determined by ¹H NMR using 1,3,5-trimethoxybenzene as internal standard.


Table S6: Screening of Solvent



Entry	Solvent	Yield (%) ^b
1	DMSO	96
2	DMA	95
3	DMF	95
4	MeCN	43
5	Dioxane	73
6	THF	82
7	DMSO (1 mL)	86

^aReaction conditions: **1** (0.2 mmol), **2** (0.3 mmol), Boc-anhydride (0.3 mmol), NaOH (20 mol%) and 4CzIPN (4 mol%) in solvent (2 mL, unless stated otherwise) under N₂ atmosphere in the dark for 4h and subsequently under light irradiation (456 nm 40 W KESSIL) at room temperature for 24h. ^bYield determined by ¹H NMR using 1,3,5-trimethoxybenzene as internal standard.

Table S7: Screening of glycine and boc-anhydride ratio




Entry	Glycine:boc-anhydride (equivalents)	Yield (%) ^b
1	1.0:1.0	70
2	1.1:1.1	72
3	1.2:1.2	72
4	1.3:1.3	75
5	1.4:1.4	86
6	1.5:1.5	93
7	1.0:1.1	60
8	1.0:1.2	63
9	1.0:1.3	54
10	1.0:1.4	50
11	1.0:1.5	56

^aReaction conditions: **1** (0.2 mmol), **2**, Boc-anhydride, NaOH (20 mol%) and 4CzIPN (4 mol%) in DMSO (2 mL) under N₂ atmosphere in the dark for 4h and subsequently under light irradiation (456 nm 40 W KESSIL) at room temperature for 24h. ^bYield determined by ¹H NMR using 1,3,5-trimethoxybenzene as internal standard.

4.3. Optimization of long chain amino acids


Table S8: Catalyst loading screening



Entry	Loading	Yield (%) ^b
1	0 mol%	0
2	1 mol%	0
3	2 mol%	Trace
4	3 mol%	5
5	4 mol%	15
6	5 mol%	20

^aReaction conditions: **1** (0.2 mmol), **2** (0.2 mmol), NaOH (0.2 mmol) and 4CzIPN in 2 mL DMSO under N₂ atmosphere and light irradiation (456 nm 40 W KESSIL) at room temperature for 72h. ^bYield determined by ¹H NMR using 1,3,5-trimethoxybenzene as internal standard.


Table S9: Base screening



Entry	Base	Yield (%) ^b
1	CS ₂ CO ₃	43
2	Li ₂ CO ₃	16
3	Na ₂ CO ₃	10
4	NaOH	20
5	K ₃ PO ₄	17
6	DIPEA	0
7	No base	0

^aReaction conditions: **1** (0.2 mmol), **2** (0.2 mmol), Base (0.2 mmol) and 4CzIPN (5 mol%) in 2 mL DMSO under N₂ atmosphere and light irradiation (456 nm 40 W KESSIL) at room temperature for 72h. ^bYield determined by ¹H NMR using 1,3,5-trimethoxybenzene as internal standard.

Table S10: Solvent screening



Entry	Solvent	Yield (%) ^b
1	DMSO	43
2	DMF	40
3	DMA	38
4	MeCN	10
5	THF	16
6	DMSO (1 mL)	19

^aReaction conditions: **1** (0.2 mmol), **2** (0.2 mmol), Cs₂CO₃ (0.2 mmol) and 4CzIPN (5 mol%) in 2 mL solvent under N₂ atmosphere and light irradiation (456 nm 40 W KESSIL) at room temperature for 72h.

^bYield determined by ¹H NMR using 1,3,5-trimethoxybenzene as internal standard.

Table S11: Temperature screening



Entry	Temperature (°C)	Yield (%) ^b
1	r.t.	43
2	50	48
3	70	62

^aReaction conditions: **1** (0.2 mmol), **2** (0.2 mmol), Cs₂CO₃ (0.2 mmol) and 4CzIPN (5 mol%) in DMSO (2 mL) under N₂ atmosphere and light irradiation (456 nm 40 W KESSIL) at a given temperature for 72h.

^bYield determined by ¹H NMR using 1,3,5-trimethoxybenzene as internal standard.

5. Stern-Volmer experiments




Figure S2. Stern-Volmer experiment between 4CzIPN and DPE.




Figure S3. Stern-Volmer experiment between 4CzIPN and NaOH.




Figure S4. Stern-Volmer experiment between 4CzIPN and deprotonated boc-glycine.

6. Synthetic procedures literature protocols


6.1. Procedure D

Procedure A (modification from Ahmad *et al.*¹): Unprotected amino acid (0.03 mol) was dissolved in 25 mL of 10% sodium hydroxide solution (2.5g in 25 mL) and benzoyl chloride (4,14 mL, 0,038 mol) was added in five equal portions and stirred vigorously after each addition until all the benzoyl chloride has reacted. The reaction mixture was neutralized carefully with concentrated hydrochloric acid and the precipitate was washed with methylene chloride to remove any little benzoic acid formed and was then recrystallised in boiling water. The pure solid was allowed to dry for 24 hours in vacuo at 25 °C.

6.2. Procedure E

Procedure B: A slurry of unprotected amino acid (25 mmol) in 70 mL methylene chloride was treated with Chlorotrimethylsilane (2 equiv.) and was allowed to reflux for 140 minutes. The reaction mixture was cooled to 0 °C and was then treated with Triethylamine (1.5 equiv.). After this mixture stirred for about 20 min, a solution of benzoyl chloride (1 equiv.) in 10 mL of methylene chloride was added dropwise to the reaction mixture over a period of 15 min. The reaction mixture was allowed to stir for 30 min. at 0 °C and then for 18 hours at 25 °C. Methylene chloride was removed in vacuo and 100 mls of NaOH solution (2N) was added to the residue. This mixture was allowed to stir for 1 hour before the mixture was acidified to pH = 1 with hydrochloric acid solution (2M). The acidified mixture was then extracted with ethyl acetate (2 × 100 mls), dried over sodium sulfate, and concentrated in vacuo. The resulting white solid was recrystallized from a 50% ethanol/water mixture yielding a white solid, which was allowed to dry for 24 hours in vacuo at 25 °C.

2,4,5,6-tetrakis(carbazol-9-yl)-4,6-dicyanobenzene (4CzIPN)




4CzIPN was synthesized according to previous literature.²

NaH (60% in oil, 1.2 g, 30 mmol) was added slowly to a stirred solution of carbazole (3.34 g, 20.0 mmol) in dry THF (80 mL) under a nitrogen atmosphere at r.t. After 30 min, tetrafluoroisophthalonitrile (0,800 g, 4 mmol) was added. After, the mixture was stirred at r.t. for 12 h and 4 mL water was added to the reaction mixture to quench the excess NaH. The resulting mixture was then concentrated under reduced pressure and purified by column chromatography (gradient towards hexane/DCM 1:1). NMR spectrum matched with previous

report.³


Benzoylglycine



Benzoylglycine was prepared using Procedure D.

¹H NMR (400 MHz, (CD₃)₂SO) δ 8.82 (t, *J* = 5.7 Hz, 1H), δ 7.89-7.87 (m, 2H), δ 7.57-7.47 (m, 3H), δ 3.95 (d, *J* = 5.9 Hz, 2H), δ 3.39 (br, 1H). ¹³C NMR (101 MHz, (CD₃)₂SO) δ 171.8, 167.0, 134.3, 131.9, 128.8, 127.7, 41.7.


4-(Benzoylamino)butanoic acid



4-(Benzoylamino)butanoic acid was prepared using procedure D. ¹H NMR (400 MHz, DMSO) δ 8.46 (t, *J* = 5.2 Hz, 1H), 7.90 – 7.77 (m, 2H), 7.58 – 7.36 (m, 3H), 3.29 (dd, *J* = 12.7, 6.8 Hz, 2H), 2.29 (t, *J* = 7.4 Hz,

2H), 1.86 – 1.69 (m, 2H). ¹³C NMR (101 MHz, CDCl₃) 174.7, 166.7, 135.1, 131.5, 128.7, 127.6, 39.1, 31.7, 25.0.

12-(Benzoylamino)dodecanoic acid




12-(Benzoylamino)dodecanoic acid was prepared using procedure E.

¹H NMR (400 MHz, (CD₃)₂SO) δ 8.40 (br, 1H), δ 7.84-7.82 (m, 2H), δ 7.53-7.43 (m, 3H), δ 3.25 (dd, *J* = 13.0, 6.8 Hz, 2H), δ 2.18 (t, *J* = 7.4 Hz, 2H), δ (dt, *J* = 14.0, 6.8 Hz, 4H), δ 1.29-1.25 (m, 14H). ¹³C NMR (101 MHz, (CD₃)₂SO) δ 174.9, 166.5, 135.2, 131.4, 128.7, 127.6, 39.7, 34.1, 29.6, 29.4, 29.4, 29.4, 29.2, 29.2, 29.0, 27.0, 25.0.

7. Synthesis of hydroaminoalkylated products


Tert-butyl (3,3-diphenylpropyl)carbamate (from 3)



The compound was synthesized according to Condition A, using 1,1-diphenylethylene and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 7.29-7.15 (m, 10H), δ 4.47 (br, 1H), δ 3.95 (t, *J* = 7.9 Hz, 1H), δ 3.07 (dd, *J* = 12.9, 6.2 Hz, 2H), δ 2.25 (dd, *J* = 14.6, 7.6 Hz, 2H), δ 1.42 (s, 9H). ¹³C NMR (101 MHz, CDCl₃) δ 155.9, 144.3, 128.6, 127.8, 126.3, 79.1, 48.9, 39.4, 35.8, 28.4. HRMS (ESI) [M+Na]⁺ calcd. 334.1778, found 334.1763.


Tert-butyl (3,3-bis(4-fluorophenyl)propyl)carbamate (from 4)



The compound was synthesized according to Condition A, using 4,4'-(ethene-1,1-diyl)bis(fluorobenzene) and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 7.15 (dd, *J* = 8.5, 5.4 Hz, 4H), 6.96 (t, *J* = 8.7 Hz, 4H), 4.51 (s, 1H), 3.93 (t, *J* = 7.8 Hz, 1H), 3.05 (dd, *J* = 13.0, 6.3 Hz, 2H), 2.19 (dd, *J* = 14.9, 7.8 Hz, 2H), 1.43 (s, 9H). ¹³C NMR (101 MHz, CDCl₃) δ 162.8, 160.4, 156.0, 140.0, 139.9, 129.3, 129.2, 115.7, 115.5, 79.4, 47.4, 39.4, 36.2, 28.5. HRMS (ESI) [M+Na]⁺ calcd. 370.1589, found 370.1594.


Tert-butyl (3-phenylpropyl)carbamate (from 5)



The compound was synthesized according to Condition A, using styrene and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 7.32 – 7.21 (m, 2H), 7.18 (t, *J* = 6.8 Hz, 3H), 4.54 (s, 1H), 3.14 (q, *J* = 6.2 Hz, 2H), 2.62 (d, *J* = 7.9 Hz, 2H), 1.86 – 1.73 (m, 2H), 1.44 (s, 9H). ¹³C NMR (101 MHz, CDCl₃) δ 156.1, 144.5, 142.4, 141.7, 128.7, 128.5, 128.5, 128.4, 127.8, 126.5, 126.0, 125.8, 79.2, 40.4, 33.8, 33.3, 31.9, 28.6. HRMS (ESI) [M+Na]⁺ calcd. 258.1464, found 258.1457.


Tert-butyl (3-(4-bromophenyl)propyl)carbamate (from 6)



The compound was synthesized according to Condition A, using *p*-bromostyrene and boc-glycine.

$^1\text{H NMR}$ (400 MHz, CDCl_3) $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.39 (d, J = 8.2 Hz, 2H), 7.04 (d, J = 8.2 Hz, 2H), 4.55 (s, 1H), 3.13 (q, J = 6.2 Hz, 2H), 2.64 – 2.52 (m, 2H), 1.84 – 1.72 (m, 2H), 1.44 (s, 9H). $^{13}\text{C NMR}$ (101 MHz, CDCl_3) δ 156.1, 140.7, 131.6, 130.2, 119.8, 79.3, 40.2, 32.6, 31.7, 28.5. **HRMS (ESI)** $[\text{M}+\text{Na}]^+$ calcd. 336.0570, found 336.0568.


Tert-butyl (3-(4-(trifluoromethyl)phenyl)propyl)carbamate (from 7)



The compound was synthesized according to Condition A, using *p*-trifluoromethylstyrene and boc-glycine.

$^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.53 (d, J = 8.0 Hz, 1H), 7.29 (d, J = 8.0 Hz, 1H), 4.54 (s, 1H), 3.16 (q, J = 6.3 Hz, 1H), 2.75 – 2.64 (m, 1H), 1.83 (dt, J = 14.7, 7.4 Hz, 1H), 1.44 (s, 5H). $^{13}\text{C NMR}$ (101 MHz, CDCl_3) δ 156.1, 145.9, 128.8, 128.7, 128.4, 125.8, 125.6, 125.5, 125.5, 125.5, 123.1, 79.4, 40.3, 33.1, 31.7, 28.6. **HRMS (ESI)** $[\text{M}+\text{Na}]^+$ calcd. 326.1338, found 326.1347.


Tert-butyl (3-(4-cyanophenyl)propyl)carbamate (from 8)



The compound was synthesized according to Condition A, using *p*-cyanostyrene and boc-glycine.

$^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.57 (d, J = 8.1 Hz, 2H), δ 7.28 (d, J = 8.1 Hz, 2H), δ 4.56 (s, 1H), δ 3.15 (m, 2H), δ 2.7 (t, J = 7.7 Hz, 2H), δ 1.82 (quint J = 7.6 Hz), δ 1.44 (s, 9H). $^{13}\text{C NMR}$ (101 MHz, CDCl_3) δ 156.0, 147.3, 132.3, 129.2, 119.0, 110.0, 79.4, 41.0, 33.3, 31.4, 28.4. **HRMS (ESI)** $[\text{M}+\text{Na}]^+$ calcd. 283.1417, found 283.148.


Tert-butyl (3-(4-(tert-butyl)phenyl)propyl)carbamate (from 9)



The compound was synthesized according to Condition A, using *p*-*t*-butylstyrene and boc-glycine.

$^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.30 (d, J = 8.2 Hz, 2H), 7.11 (d, J = 8.2 Hz, 2H), 4.50 (s, 1H), 3.15 (q, J = 6.1 Hz, 2H), 2.68 – 2.51 (m, 2H), 1.89 – 1.71 (m, 2H), 1.44 (s, 9H), 1.30 (s, 9H). $^{13}\text{C NMR}$ (101 MHz, CDCl_3) δ 156.1, 148.9, 138.6, 128.2, 125.4, 79.3, 40.5, 34.5, 32.7, 31.8, 31.5, 28.6. **HRMS (ESI)** $[\text{M}+\text{H}]^+$ calcd. 292.2271, found 292.2279.


Tert-butyl (3-(4-(trimethylsilyl)phenyl)propyl)carbamate (from 10)



The compound was synthesized according to Condition A, using *p*-trimethylsilylstyrene and boc-glycine.


$^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.44 (d, J = 7.7 Hz, 2H), 7.17 (d, J = 7.7 Hz, 2H), 4.50 (s, 1H), 3.15 (q, J = 6.2 Hz, 2H), 2.67 – 2.57 (m, 2H), 1.90 – 1.73 (m, 2H), 1.44 (s, 9H), 0.25 (s, 9H). $^{13}\text{C NMR}$ (101 MHz, CDCl_3) δ 156.1, 142.3, 137.8, 133.7, 128.0, 79.3, 40.5, 33.3, 31.8, 28.6, 0.1. **HRMS (ESI)** $[\text{M}+\text{Na}]^+$ calcd. 330.2945, found 330.1979.

Tert-butyl (3-([1,1'-biphenyl]-4-yl)propyl)carbamate (from 11)




The compound was synthesized according to Condition A, using p-phenylstyrene and boc-glycine. **¹H NMR** (400 MHz, CDCl₃) δ 7.59 – 7.54 (m, 2H), 7.51 (d, *J* = 8.1 Hz, 2H), 7.42 (t, *J* = 7.6 Hz, 1H), 7.31 (t, *J* = 7.3 Hz, 2H), 7.25 – 7.21 (m, 2H), 4.55 (s, 1H), 3.18 (q, *J* = 6.2 Hz, 2H), 2.68 (t, 1H), 1.89 – 1.79 (m, 2H), 1.45 (s, 9H). **¹³C NMR** (101 MHz, CDCl₃) δ 156.0, 141.1, 140.7, 139.0, 128.8, 128.7, 127.2, 127.1, 127.0, 79.2, 77.4, 77.0, 76.7, 40.3, 32.8, 31.7, 28.4. **HRMS (ESI)** [M+Na]⁺ calcd. 334.1777, found 334.1763.

Tert-butyl (3-(2-chlorophenyl)propyl)carbamate (from 12)




The compound was synthesized according to Condition A, using o-chlorostyrene and boc-glycine. **¹H NMR** (400 MHz, CDCl₃) δ 7.40 – 7.27 (m, 1H), 7.24 – 7.07 (m, 3H), 4.57 (br, 1H), 3.17 (q, *J* = 6.4 Hz, 2H), 2.80 – 2.71 (m, 2H), 1.81 (dt, *J* = 14.7, 7.3 Hz, 2H), 1.45 (s, 9H). **¹³C NMR** (101 MHz, CDCl₃) δ 156.0, 139.2, 133.9, 130.4, 129.5, 127.5, 126.8, 79.2, 40.2, 30.8, 30.1, 28.4. **HRMS (ESI)** [M+H]⁺ calcd. 292.1075, found 292.1088

Tert-butyl (3-mesitylpropyl)carbamate (from 13)




The compound was synthesized according to Condition A, using 2,4,6-trimethylstyrene and boc-glycine. **¹H NMR** (400 MHz, CDCl₃) δ 6.82 (s, 2H), 4.55 (s, 1H), 3.20 (q, *J* = 6.1 Hz, 2H), 2.64 – 2.53 (m, 2H), 2.27 (s, 6H), 2.23 (s, 3H), 1.68 – 1.58 (m, 2H), 1.45 (s, 9H). **¹³C NMR** (101 MHz, CDCl₃) δ 156.2, 136.0, 135.6, 135.2, 129.1, 79.3, 41.1, 29.8, 28.6, 26.7, 20.9, 19.8. **HRMS (ESI)** [M+Na]⁺ calcd. 300.1934, found 300.1945

Tert-butyl (3-(perfluorophenyl)propyl)carbamate (from 14)




The compound was synthesized according to Condition A, using pentafluorostyrene and boc-glycine. **¹H NMR** (400 MHz, CDCl₃) δ 4.58 (s, 1H), 3.16 (q, *J* = 6.3 Hz, 2H), 2.73 (t, *J* = 7.6 Hz, 2H), 1.83 – 1.74 (m, 2H), 1.44 (s, 9H). **¹³C NMR** (101 MHz, CDCl₃) δ 156.0, 144.1, 140.3, 138.6, 114.6, 79.6, 40.0, 29.7, 28.5, 19.8. **HRMS (ESI)** [M+H]⁺ calcd. 326.1174, found 326.1159

Tert-butyl (3-phenylbutyl)carbamate (from 15)



The compound was synthesized according to Condition A, using α-methylstyrene and boc-glycine. **¹H NMR** (400 MHz, CDCl₃) δ 7.28 (dt, *J* = 8.1, 3.3 Hz, 2H), 7.18 (dd, *J* = 7.3, 5.6 Hz, 3H), 4.43 (s, 1H), 3.00 (dt, *J* = 13.2, 6.8 Hz, 2H), 2.84 – 2.62 (m, 1H), 1.77 (ddd, *J* = 14.7, 9.5, 5.4 Hz, 2H), 1.42 (s, 9H), 1.26 (d, *J* = 7.0 Hz, 3H). **¹³C NMR** (101 MHz, CDCl₃) δ 156.0, 146.8, 128.6, 127.0, 126.3, 79.2, 39.3, 38.5, 37.8, 28.5, 22.4. **HRMS (ESI)** [M+H]⁺ calcd. 250.1802, found 250.1813.


Tert-butyl (3-(4-chlorophenyl)butyl)carbamate (from 16)



The compound was synthesized according to Condition A, using 1-chloro-4-(prop-1-en-2-yl)benzene and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 7.26-7.24 (m, 2H), δ 7.12-7.10 (m, 2H), δ 4.41 (br, 1H), δ 3.08-2.95 (m, 2H), δ 2.73 (sext, *J* = 7.2 Hz, 1H), δ (1.81-1.67, 2H), δ 1.42 (s, 9H), 1.24, *J* = 7.0 Hz, 3H). **¹³C NMR** (101 MHz, CDCl₃) δ 155.9, 145.1, 131.8, 128.6, 128.3, 79.2, 39.1, 38.3, 37.2, 28.4, 22.2. **HRMS (ESI)** [M+H]⁺ calcd. 306.1231, found 306.1240.


Tert-butyl (3-(9H-fluoren-2-yl)butyl)carbamate (17)



The compound was synthesized according to Condition A, using 2-(prop-1-en-2-yl)-9H-fluorene and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 7.71 (dd, *J* = 17.2, 7.6 Hz, 2H), 7.51 (d, *J* = 7.4 Hz, 1H), 7.37 – 7.32 (m, 2H), 7.29 – 7.23 (m, 1H), 7.18 (d, *J* = 7.9 Hz, 1H), 4.43 (s, 1H), 3.86 (s, 2H), 3.03 (dt, *J* = 12.7, 6.4 Hz, 2H), 2.82 (dd, *J* = 14.3, 7.1 Hz, 1H), 1.81 (dt, *J* = 14.4, 7.0 Hz, 2H), 1.42 (s, 9H), 1.31 (d, *J* = 6.9 Hz, 3H). **¹³C NMR** (101 MHz, CDCl₃) δ 156.1, 145.6, 143.8, 143.3, 141.8, 140.1, 126.8, 126.5, 125.7, 125.1, 123.6, 120.0, 119.8, 79.2, 39.4, 38.7, 38.0, 37.0, 28.6, 22.6. **HRMS (ESI)** [M+Na]⁺ calcd. 360.1933, found 360.1939.


Tert-butyl (3-(naphthalen-2-yl)butyl)carbamate (from 18)



The compound was synthesized according to Condition A, using 2-(prop-1-en-2-yl)naphthalene and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 7.79 (dt, *J* = 7.5, 3.8 Hz, 1H), 7.61 (s, 1H), 7.49 – 7.38 (m, 1H), 7.34 (dd, *J* = 8.5, 1.5 Hz, 1H), 4.42 (s, 1H), 3.05 (dd, *J* = 13.3, 6.7 Hz, 1H), 2.93 (dd, *J* = 14.2, 7.1 Hz, 1H), 1.87 (q, *J* = 7.4 Hz, 1H), 1.42 (s, 3H), 1.36 (d, *J* = 6.9 Hz, 1H). **¹³C NMR** (101 MHz, CDCl₃) δ 156.1, 144.2, 133.8, 132.5, 128.4, 127.7, 127.7, 126.1, 125.6, 125.4, 125.3, 79.4, 40.5, 38.4, 38.0, 28.6, 22.4. **HRMS (ESI)** [M+H]⁺ calcd. 322.1777, found 322.1805.


Tert-butyl (2,3,3-triphenylpropyl)carbamate (from 19)



The compound was synthesized according to Condition A, using triphenylethylene and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 7.42 (d, *J* = 7.4 Hz, 2H), 7.31 (t, *J* = 7.6 Hz, 2H), 7.21 – 7.00 (m, 10H), 6.95 (d, *J* = 7.2 Hz, 1H), 4.12 (d, *J* = 11.4 Hz, 2H), 3.71 (s, 1H), 3.55 (s, 1H), 3.10 – 2.97 (m, 1H), 1.34 (s, 9H). **¹³C NMR** (101 MHz, CDCl₃) δ 155.7, 143.2, 141.2, 129.0, 128.7, 128.5, 128.3, 128.2, 128.2, 126.8, 126.7, 126.0, 79.1, 56.6, 49.7, 45.3, 35.6, 28.5. **HRMS (ESI)** [M+Na]⁺ calcd. 410.2090, found 410.2091.


Tert-butyl (2-cyano-3,3-diphenylpropyl)carbamate (from 20)



The compound was synthesized according to Condition A, using 3,3-diphenylacrylonitrile and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 7.37 – 7.23 (m, 10H), 4.93 (s, 1H), 4.12 (d, *J* = 8.9 Hz, 1H), 3.76 (d, *J* = 4.0 Hz, 1H), 3.52 – 3.44 (m, 1H), 3.01-3.08 (m, 1H), 1.44 (s, 9H). **¹³C NMR** (101 MHz, CDCl₃) δ 155.7, 140.20, 140.0, 129.2, 129.1, 128.3, 127.8, 127.7, 127.6, 120.4, 80.3, 51.3, 42.0, 38.1, 35.6, 28.4, 26.6, 26.5, 23.0. **HRMS (ESI)** [M+Na]⁺ calcd. 359.1730, found 359.1743.


Tert-butyl (2-cyano-3-phenylpropyl)carbamate (from 21)



The compound was synthesized according to Condition A, using cinnamionitrile and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 7.35-7.24 (m, 5H), δ 4.97 (br, 1H), δ 3.48-3.45 (m, 1H), δ 3.26-3.19 (m, 1H), 3.16-3.10 (m, 1H), δ 2.94-2.83 (m, 2H), δ 1.45 (s, 9H). **¹³C NMR** (101 MHz, CDCl₃) δ 155.7, 136.2, 129.0, 128.9, 127.4, 120.5, 80.3, 42.1, 35.8, 35.0, 28.3. **HRMS (ESI)** [M+Na]⁺ calcd. 283.1417, found 283.1427.


Tert-butyl (2-methyl-3-phenylpropyl)carbamate (from 22)



The compound was synthesized according to Condition A, using β-methylstyrene and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 7.27 (dd, *J* = 10.5, 4.1 Hz, 2H), 7.17 (dd, *J* = 20.0, 7.2 Hz, 2H), 4.54 (s, 1H), 3.06 (ddd, *J* = 40.2, 12.9, 6.3 Hz, 1H), 2.69 (dd, *J* = 13.4, 5.7 Hz, 1H), 2.37 (dd, *J* = 13.5, 8.4 Hz, 1H), 1.92 (dq, *J* = 13.4, 6.6 Hz, 1H), 1.44 (s, 9H), 0.87 (d, *J* = 6.7 Hz, 3H). **¹³C NMR** (101 MHz, CDCl₃) δ 156.2, 140.6, 129.2, 128.4, 126.1, 79.2, 46.6, 41.1, 36.0, 28.6, 17.5. **HRMS (ESI)** [M+Na]⁺ calcd. 272.1621, found 272.1610.


Methyl 3-((tert-butoxycarbonyl)amino)-2-(4-chlorobenzyl)propanoate (from 23)



The compound was synthesized according to Condition A, using methyl (*E*)-3-(4-chlorophenyl)acrylate and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 7.24 (d, *J* = 8.4 Hz, 2H), 7.09 (d, *J* = 8.4 Hz, 2H), 4.85 (s, 1H), 3.64 (s, 3H), 3.42 – 3.21 (m, 2H), 2.98 – 2.86 (m, 2H), 2.77 (d, *J* = 6.8 Hz, 1H), 1.43 (s, 9H). **¹³C NMR** (101 MHz, CDCl₃) δ 174.5, 155.9, 137.0, 132.6, 130.3, 128.8, 79.7, 77.5, 77.2, 76.8, 52.0, 47.5, 41.7, 35.3, 28.5. **HRMS (ESI)** [M+Na]⁺ calcd. 350.1130, found 350.1139.


Methyl 3-((tert-butoxycarbonyl)amino)-2-(4-methylbenzyl)propanoate (from 24)



The compound was synthesized according to Condition A, using methyl (*E*)-3-(4-methylphenyl)acrylate and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 7.05 (dd, *J* = 18.3, 8.0 Hz, 4H), 4.84 (s, 1H), 3.64 (s, 3H), 3.23-3.27 (m, 2H), 2.95 – 2.86 (m, 2H), 2.83 – 2.67 (m, 1H), 2.30 (s, 3H), 1.42 (s, 9H). **¹³C NMR** (101 MHz, CDCl₃) δ 174.8, 155.9, 136.2, 135.3, 129.3, 128.8, 79.5, 51.9, 47.6, 41.6, 35.6, 28.5, 21.1. **HRMS (ESI)** [M+H]⁺ calcd. 308.1856, found 308.1869.


Tert-butyl (2,3-diphenylpropyl)carbamate (from 25)



The compound was synthesized according to Condition A, using *trans*-stilbene and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 7.27 (dd, *J* = 12.9, 5.3 Hz, 2H), 7.19 (dd, *J* = 12.9, 7.1 Hz, 3H), 7.12 (d, *J* = 7.0 Hz, 3H), 7.02 (d, *J* = 7.1 Hz, 2H), 4.33 (s, 1H), 3.55 (d, *J* = 5.4 Hz, 1H), 3.24 (d, *J* = 8.8 Hz, 1H), 3.06 (s, 1H), 2.91 (qd, *J* = 13.6, 7.4 Hz, 2H), 1.38 (s, 9H). **¹³C NMR** (101 MHz, CDCl₃) δ 156.0, 142.2, 139.7, 129.18, 128.7, 128.4, 128.1, 126.9, 126.2, 79.3, 48.0, 45.5, 40.8, 28.5. **HRMS (ESI)** [M+Na]⁺ calcd. 334.1777, found 334.1775.


Tert-butyl (2-phenyl-2-(phenylamino)ethyl)carbamate (from 26)



The compound was synthesized according to Condition A, using (*E*)-*N*,1-diphenylmethanimine and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 7.34 (dt, *J* = 15.0, 7.5 Hz, 4H), 7.25 (t, *J* = 6.9 Hz, 1H), 7.07 (t, *J* = 7.8 Hz, 2H), 6.63 (t, *J* = 7.2 Hz, 1H), 6.49 (d, *J* = 7.9 Hz, 2H), 4.91 (s, 1H), 4.78 (s, 1H), 4.47 – 4.36 (m, 1H), 3.54 – 3.36 (m, 2H), 1.46 (s, 9H). **¹³C NMR** (101 MHz, CDCl₃) δ 157.3, 147.6, 141.3, 129.2, 129.0, 127.75, 126.7, 117.4, 113.4, 80.1, 60.2, 47.5, 28.5. **HRMS (ESI)** [M+H]⁺ calcd. 312.1838, found 312.1830.


Tert-butyl (3-(pyridin-2-yl)propyl)carbamate (from 27)



The compound was synthesized according to Condition A, using 2-vinylpyridine and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 8.51 (d, *J* = 4.4 Hz, 1H), δ 7.59 (td, *J* = 7.7, 1.7 Hz, 1H), δ 7.15 (d, *J* = 7.8 Hz, 1H), δ 7.10 (dd, *J* = 7.1, 5.3 Hz, 1H), δ 4.77 (br, 1H), δ 3.17 (dd, *J* = 12.6, 6.2 Hz, 2H), δ 2.83 (t, 7.6 Hz, 2H), δ 1.93 (p, *J* = 7.6 Hz, 2H), δ 1.44 (s, 9H). **¹³C NMR** (101 MHz, CDCl₃) δ 161.4, 156.0, 149.3, 136.5, 122.9, 121.2, 79.0, 40.1, 35.5, 29.9, 28.5. **HRMS (ESI)** [M+H]⁺ calcd. 237.1598, found 237.1586.


Tert-butyl (3-phenyl-3-(pyridin-2-yl)propyl)carbamate (from 28)



The compound was synthesized according to Condition A, using 2-(1-phenylvinyl)pyridine and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 8.56 (d, *J* = 4.0 Hz, 1H), 7.55 (td, *J* = 7.7, 1.8 Hz, 1H), 7.34 – 7.27 (m, 4H), 7.19 (t, *J* = 7.0 Hz, 1H), 7.15 – 7.06 (m, 2H), 4.61 (s, 1H), 4.12 (t, *J* = 7.7 Hz, 1H), 3.20 – 2.97 (m, 2H), 2.47 (td, *J* = 14.2, 7.2 Hz, 1H), 2.26 (td, *J* = 14.3, 7.3 Hz, 1H), 1.41 (s, 9H). **¹³C NMR** (101 MHz, CDCl₃) δ 163.29, 156.04, 149.36, 143.35, 136.68, 128.77, 128.11, 126.76, 123.10, 121.59, 79.17, 51.33, 39.40, 35.21, 28.57. **HRMS (ESI)** [M+H]⁺ calcd. 313.1911, found 313.1919.

Tert-butyl (2-(benzo[*b*]thiophen-2-yl)ethyl)carbamate (from 29)




The compound was synthesized according to Condition A, using 2-vinylbenzo[*b*]thiophene and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 7.74 (d, *J* = 7.9 Hz, 1H), δ 7.65 (d, *J* = 7.6 Hz, 1H), δ 7.31-7.25 (m, 2H), δ 7.01 (s, 1H), δ 4.57 (br, 1H), 3.21 (dd, *J* = 12.9, 7.4 Hz, 2H), δ 2.93 (t, *J* = 7.5 Hz, 2H), δ 1.93 (p, *J* = 7.6 Hz,

2H), δ 1.44 (s, 9H). **¹³C NMR** (101 MHz, CDCl₃) δ 156.0, 145.3, 140.2, 139.3, 124.1, 123.6, 122.8, 122.1, 120.9, 79.3, 40.0, 31.4, 28.4, 28.1. **HRMS (ESI)** [M+Na]⁺ calcd. 314.1185, found 314.1197.


Tert-butyl (3-(4-methylthiazol-5-yl)propyl)carbamate (from 30)



The compound was synthesized according to Condition A, using 4-methyl-5-vinylthiazole and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 8.55 (s, 1H), δ 4.57 (br, 1H), δ (dd, J = 12.5, 6.1 Hz, 2H), δ 2.79 (t, J = 7.7 Hz, 2H), δ 2.38 (s, 3H), δ 1.81 (p, J = 7.2 Hz, 2H), δ 1.45 (s, 9H). **¹³C NMR** (101 MHz, CDCl₃) δ 155.9; 149.0, 148.7, 130.9, 79.4, 40.0, 32.0, 28.4, 23.6, 14.8. **HRMS (ESI)** [M+Na]⁺ calcd. 279.1138, found 279.1149.


Ethyl 4-((tert-butoxycarbonyl)amino)butanoate (from 31)



The compound was synthesized according to Condition A, using ethyl acrylate and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 4.63 (s, 1H), 4.13 (q, J = 7.1 Hz, 2H), 3.16 (q, J = 6.4 Hz, 2H), 2.34 (t, J = 7.3 Hz, 2H), 1.81 (p, J = 7.1 Hz, 2H), 1.44 (s, 9H), 1.26 (t, J = 7.1 Hz, 3H). **¹³C NMR** (101 MHz, CDCl₃) δ 173.4, 156.1, 79.3, 60.6, 40.1, 31.8, 28.5, 25.5, 14.3. **HRMS (ESI)** [M+Na]⁺ calcd. 254.1363, found 254.1375.


Adamantan-1-yl 4-((tert-butoxycarbonyl)amino)butanoate (from 32)



The compound was synthesized according to Condition A, using adamantyl acrylate and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 4.62 (br, 1H), δ 3.15 (dd, J = 21.2, 5.9 Hz, 1H), δ 2.25 (t, J = 7.3 Hz, 1H), δ 2.16-2.10 (m, 9H), δ 1.76 (p, J = 6.8 Hz, 2H), δ 1.69-1.63 (m, 6H), δ 1.44 (s, 9H). **¹³C NMR** (101 MHz, CDCl₃) δ 172.4, 155.9, 80.6, 79.1, 41.4, 40.1, 36.2, 33.1, 30.9, 28.4, 25.4. **HRMS (ESI)** [M+H]⁺ calcd. 338.2326, found 338.2327.


Tert-butyl (4-(dimethylamino)-4-oxobutyl)carbamate (from 33)



The compound was synthesized according to Condition A, using *N,N*-dimethylacrylamide and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 4.84 (br, 1H), δ 3.17 (q, J = 6.4 Hz, 2H), δ 3.00 (s, 3H), δ 2.94 (s, 3H), δ 2.36 (t, J = 7.1 Hz, 2H), δ 1.83 (p, J = 6.9 Hz, 2H), δ 1.44 (s, 9H). **¹³C NMR** (101 MHz, CDCl₃) δ 172.5, 156.1, 79.0, 40.3, 37.2, 35.5, 30.6, 28.4, 25.2. **HRMS (ESI)** [M+H]⁺ calcd. 231.1703, found 231.1709.


Tert-butyl (2-methyl-4-oxopentyl)carbamate (from 34)



The compound was synthesized according to Condition A, using (*E*)-pent-3-en-2-one and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 4.57 (br, 1H), δ 3.12 (dd, J = 12.9, 6.5 Hz, 2H), δ 2.47-2.40 (m, 4H), δ 1.76 (p, J = 7.0 Hz, 2H), δ 1.44 (s, 9H), δ 1.06 (t, J = 7.3 Hz, 3H). **¹³C NMR** (101 MHz, CDCl₃) δ 211.1, 156.0, 79.2, 40.1, 39.4, 36.0, 28.4, 24.2, 7.8. **HRMS (ESI)** [M+Na]⁺ calcd. 238.1414, found 238.1412.


Methyl 4-((tert-butoxycarbonyl)amino)-2-methylbutanoate (from 35)



The compound was synthesized according to Condition A, using methyl methacrylate and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 4.57 (s, 1H), 3.68 (s, 3H), 3.15 (s, 2H), 2.51 (dd, *J* = 13.9, 6.9 Hz, 1H), 1.83 (dt, *J* = 14.6, 7.1 Hz, 1H), 1.44 (s, 9H), 1.18 (d, *J* = 7.0 Hz, 3H). **¹³C NMR** (101 MHz, CDCl₃) δ 176.9, 156.01, 79.4, 77.5, 77.2, 76.8, 51.8, 38.7, 37.2, 34.0, 28.6, 17.2. **HRMS (ESI)** [M+Na]⁺ calcd. 254.1363, found 254.1375.


Dimethyl 2-(2-((tert-butoxycarbonyl)amino)ethyl)succinate (from 36)



The compound was synthesized according to Condition A, using dimethyl 2-methylenesuccinate and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 4.74 (s, 1H), 3.71 (s, 3H), 3.67 (s, 3H), 3.16 (ddd, *J* = 26.8, 13.2, 6.7 Hz, 2H), 2.90 (td, *J* = 8.2, 4.2 Hz, 1H), 2.75 (dd, *J* = 16.7, 8.8 Hz, 1H), 2.49 (dd, *J* = 16.6, 5.3 Hz, 1H), 1.88 – 1.69 (m, 2H), 1.44 (s, 9H). **¹³C NMR** (101 MHz, CDCl₃) δ 175.0, 172.2, 155.9, 79.4, 52.1, 51.9, 38.8, 38.3, 35.8, 32.0, 28.5. **HRMS (ESI)** [M+Na]⁺ calcd. 312.1418, found 312.1416.


Dimethyl 2-(((tert-butoxycarbonyl)amino)methyl)succinate (from 37)



The compound was synthesized according to Condition A, using dimethyl maleate and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 4.91 (br, 1H), δ 3.72 (s, 3H), δ 3.69 (s, 3H), δ 3.46-3.36 (m, 2H), δ 3.03 (p, *J* = 6.8 Hz, 1H), δ 2.64 (ddd, *J* = 22.7, 16.8, 6.8 Hz, 2H), δ 1.43 (s, 9H). **¹³C NMR** (101 MHz, CDCl₃) δ 173.7, 172.0, 155.9, 79.6, 52.2, 41.9, 41.8, 41.4, 33.5, 28.4. **HRMS (ESI)** [M+H]⁺ calcd. 298.1261, found 298.1255.


Methyl 3-(((tert-butoxycarbonyl)amino)methyl)-5-methylhexanoate (from 38)



The compound was synthesized according to Condition A, using methyl (E)-5-methylhex-2-enoate and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 4.67 (s, 1H), 3.67 (s, 3H), 3.27 – 2.91 (m, 2H), 2.28 (d, *J* = 6.6 Hz, 2H), 2.16 – 2.01 (m, 1H), 1.72 – 1.57 (m, 2H), 1.44 (s, 9H), 1.15 (qd, *J* = 13.8, 6.9 Hz, 2H), 0.89 (dd, *J* = 7.7, 6.9 Hz, 6H). **¹³C NMR** (101 MHz, CDCl₃) δ 173.8, 156.2, 79.2, 51.7, 44.4, 41.7, 37.5, 33.9, 28.5, 25.4, 22.8. **HRMS (ESI)** [M+Na]⁺ calcd. 296.1832, found 296.1840.


Tert-butyl (3-(phenylsulfonyl)propyl)carbamate (from 39)



The compound was synthesized according to Condition A, using (vinylsulfonyl)benzene and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 7.91 (d, *J* = 7.4 Hz, 2H), δ 7.62 (dt, *J* = 15.2, 7.4 Hz, 3H), δ 4.68 (br, 1H), δ 3.23 (dd, *J* = 12.7, 6.3 Hz, 2H), δ 3.16-3.12 (m, 2H), δ 1.96-1.89 (m, 2H), δ 1.41 (s, 9H). **¹³C NMR** (101 MHz, CDCl₃) δ 155.9, 139.1, 133.8, 129.4, 128.0, 79.6, 53.7, 38.9, 28.3, 23.6. **HRMS (ESI)** [M+Na]⁺ calcd. 322.1083, found 322.1095.


Tert-butyl ((2-methyl-3-oxo-5-(prop-1-en-2-yl)cyclohexyl)methyl)carbamate (from 40)



The compound was synthesized according to Condition A, using carvone and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 4.79 (s, 1H), 4.75 (s, 1H), 4.48 (br, 1H), 3.22 – 3.14 (m, 1H), 2.90 (s, 1H), 2.71 – 2.58 (m, 2H), 2.53 – 2.41 (m, 1H), 2.31 (dd, *J* = 16.6, 8.4 Hz, 2H), 2.01 (d, *J* = 13.0 Hz, 1H), 1.74 (s, 4H), 1.43 (s, 9H), 1.08 (d, *J* = 6.9 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 212.2, 156.0, 147.4, 110.3, 110.1, 79.6, 47.1, 46.6, 46.1, 44.8, 41.5, 41.0, 39.1, 35.0, 32.0, 28.5, 20.8, 20.5, 11.8, 11.6. HRMS (ESI) [M+Na]⁺ calcd. 304.1883, found 304.1877.


Tert-butyl (3-cyanopropyl)carbamate (from 41)



The compound was synthesized according to Condition A, using acrylonitrile and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 4.69 (br, 1H), δ 3.25 (q, *J* = 6.5 Hz, 2H), δ 2.4 (t, *J* = 7.2 Hz, 2H), δ 1.87 (p, *J* = 6.8 Hz, 2H), δ 1.45 (s, 9H). ¹³C NMR (101 MHz, CDCl₃) δ 156.0, 119.2, 79.8, 39.4, 28.4, 26.1, 14.6. HRMS (ESI) [M+Na]⁺ calcd. 207.1104, found 207.1094.


Tert-butyl (4,4-diphenylbutyl)carbamate (from 42)



The compound was synthesized according to Condition A, using 1,1-diphenylethylene and 3-((Tert-butoxycarbonyl)(methyl)amino)-propanoic acid.

¹H NMR (400 MHz, CDCl₃) δ 7.28-7.21 (m, 8H), δ 7.18-7.14 (m, 2H), δ 4.43 (br, 1H), δ 3.89 (t, *J* = 7.8 Hz, 1H), δ 3.12 (dd, *J* = 13, 7.2, 2H), δ 2.08-2.03 (m, 2H), δ 1.47-1.40 (m, 11H). ¹³C NMR (101 MHz, CDCl₃) δ 155.9, 144.8, 128.5, 127.8, 126.2, 79.1, 51.0, 40.5, 32.8, 28.6, 28.4. HRMS (ESI) [M+Na]⁺ calcd. 348.1934, found 348.1918.


Tert-butyl (5,5-diphenylpentyl)carbamate (from 43)



The compound was synthesized according to Condition A, using 1,1-diphenylethylene and 4-((tert-butoxycarbonyl)amino)-butanoic acid.

¹H NMR (400 MHz, CDCl₃) δ 7.28 – 7.12 (m, 10H), 4.42 (s, 1H), 3.87 (t, *J* = 7.8 Hz, 1H), 3.06 (q, *J* = 6.4 Hz, 2H), 2.05 (dd, *J* = 15.6, 7.8 Hz, 2H), 1.49 (dd, *J* = 14.8, 7.4 Hz, 2H), 1.42 (s, 9H), 1.28 (ddd, *J* = 8.0, 6.7, 4.0 Hz, 2H). ¹³C NMR (101 MHz, CDCl₃) δ 156.1, 145.2, 128.6, 128.0, 126.3, 79.2, 51.5, 40.6, 35.5, 30.2, 28.6, 25.4. HRMS (ESI) [M+Na]⁺ calcd. 362.2090, found 362.2096.

Tert-butyl (6,6-diphenylhexyl)carbamate (from 44)




The compound was synthesized according to Condition A, using 1,1-diphenylethylene and 5-((tert-butoxycarbonyl)amino)-pentanoic acid.

¹H NMR (400 MHz, CDCl₃) δ 7.33 – 7.06 (m, 10H), 4.44 (s, 1H), 3.87 (t, *J* = 7.8 Hz, 1H), 3.05 (q, *J* = 6.1 Hz, 2H), 2.03 (dd, *J* = 14.9, 7.7 Hz, 2H), 1.42 (d, *J* = 9.2 Hz, 11H), 1.36 – 1.16 (m,

4H). **¹³C NMR** (101 MHz, CDCl₃) δ 156.1, 145.3, 128.5, 128.0, 126.2, 79.2, 51.4, 40.7, 35.8, 30.1, 28.6, 27.9, 26.9. **HRMS (ESI)** [M+Na]⁺ calcd. 376.2247, found 376.2240.


Tert-butyl (7,7-diphenylheptyl)carbamate (from 45)



The compound was synthesized according to Condition A, using 1,1-diphenylethylene and 6-((tert-butoxycarbonyl)-amino)hexanoic acid.

¹H NMR (400 MHz, CDCl₃) δ 7.30 – 7.20 (m, 8H), 7.19 – 7.11 (m, 2H), 4.45 (s, 1H), 3.87 (t, *J* = 7.8 Hz, 1H), 3.06 (q, *J* = 6.2 Hz, 2H), 2.02 (dd, *J* = 15.0, 7.7 Hz, 2H), 1.43 (s, 10H), 1.36 – 1.12 (m, 5H). **¹³C NMR** (101 MHz, CDCl₃) δ 156.1, 145.4, 128.5, 128.0, 126.2, 79.1, 77.5, 77.2, 76.8, 51.5, 40.7, 35.8, 30.2, 29.4, 28.6, 28.1, 26.8. **HRMS (ESI)** [M+Na]⁺ calcd. 390.2403, found 390.2389.


Tert-butyl (2-((2-((3,3-diphenylpropyl)amino)-2-oxoethyl)amino)-2-oxoethyl)carbamate (from 46)



The compound was synthesized according to Condition B, using 1,1-diphenylethylene and glycylglycylglycine.

¹H NMR (400 MHz, CDCl₃) δ 7.32 – 7.26 (m, 1H), 7.26 – 7.21 (m, 1H), 7.21 – 7.15 (m, 1H), 5.45 (s, 1H), 4.28 (s, 1H), 4.02 (s, 1H), 3.94 (t, *J* = 7.9 Hz, 1H), 3.28 (dd, *J* = 13.4, 6.5 Hz, 1H), 2.30 (dd, *J* = 14.3, 7.6 Hz, 1H), 1.55 (s, 1H). **¹³C NMR** (101 MHz, CDCl₃) δ 174.7, 163.7, 143.1, 127.7, 126.7, 125.5, 83.7, 48.3, 47.2, 40.3, 38.1, 34.0, 27.0. **HRMS (ESI)** [M+H]⁺ calcd. 426.2387, found 426.2399.


Tert-butyl (2-hydroxy-5,5-diphenylpentyl)carbamate (from 47)



The compound was synthesized according to Condition B, using 1,1-diphenylethylene and 4-amino-3-hydroxybutanoic acid.

¹H NMR (400 MHz, CDCl₃) δ 7.24 – 7.05 (m, 10H), 4.76 (s, 1H), 3.80 (t, *J* = 7.8 Hz, 1H), 3.69 (s, 1H), 3.61 (d, *J* = 4.6 Hz, 1H), 3.20 – 2.73 (m, 2H), 2.12 (ddd, *J* = 21.3, 19.9, 7.7 Hz, 1H), 2.06 – 1.92 (m, 1H), 1.32 (s, 9H). **¹³C NMR** (101 MHz, CDCl₃) δ 161.7, 145.0, 144.9, 130.7, 129.4, 129.2, 128.6, 128.5, 128.0, 128.0, 126.3, 93.1, 79.8, 71.8, 55.5, 51.5, 46.9, 35.7, 35.2, 33.5, 31.7, 28.5, 21.2. **HRMS (ESI)** [M+Na]⁺ calcd. 378.2040, found 378.2027.


N-(5,5-diphenylpentyl)benzamide (from 48)



The compound was synthesized according to Condition C, using 1,1-diphenylethylene and 4-benzamidobutanoic acid.

¹H NMR (400 MHz, CDCl₃) δ 7.71 – 7.62 (m, 2H), 7.46 (dd, *J* = 8.5, 6.2 Hz, 1H), 7.38 (t, *J* = 7.4 Hz, 2H), 7.30 – 7.19 (m, 8H), 7.15 (dd, *J* = 9.1, 4.4 Hz, 2H), 6.09 (s, 1H), 3.89 (t, *J* = 7.8 Hz, 1H), 3.39 (dd, *J* = 13.2, 6.9 Hz, 2H), 2.08 (dd, *J* = 15.6, 7.8 Hz, 2H), 1.63 (dt, *J* = 14.8, 7.4 Hz, 2H), 1.38 – 1.28 (m, 2H). **¹³C NMR** (101 MHz, CDCl₃) δ 167.6, 145.1, 135.0, 131.4, 128.6, 128.6, 127.9, 127.0, 126.3, 51.3, 39.9, 35.3, 29.6, 25.4. **HRMS (ESI)** [M+H]⁺ calcd. 344.2009, found 344.1991.


N-(12,12-diphenyldodecyl)benzamide (from 49)



The compound was synthesized according to Condition C, using 1,1-diphenylethylene and 12-benzamidododecanoic acid.

¹H NMR (400 MHz, CDCl₃) δ 7.767.74 (m, 2H), δ 7.50-7.40 (m, 3H), 7.28-7.13 (m, 10H), δ 6.06 (br, 1H), δ 3.87 (t, *J* = 7.8 Hz, 1H), δ 3.45 (dd, *J* = 13.1, 7.1 Hz, 2H), δ 2.02 (dd, *J* = 14.8, 7.8 Hz, 2H), δ 1.61 (p, *J* = 7.6 Hz, 2H), δ 1.37-1.22 (m, 18H). **¹³C NMR** (101 MHz, CDCl₃) δ 167.5, 145.4, 135.0, 131.3, 128.6, 128.3, 127.9, 126.8, 126.0, 51.4, 40.1, 35.7, 29.7, 29.6, 29.6, 29.5, 29.5, 29.5, 29.3, 28.0, 27.0. **HRMS (ESI)** [M+Na]⁺ calcd. 456.3261, found 456.3282.


N-(5-phenylpentyl)benzamide (50)



The compound was synthesized according to Condition C, using 4-benzamidobutanoic acid and styrene.

¹H NMR (400 MHz, CDCl₃) δ 7.77 – 7.70 (m, 1H), 7.50 – 7.35 (m, 2H), 7.31 – 7.12 (m, 3H), 6.22 (s, 1H), 3.47 – 3.39 (m, 1H), 2.66 – 2.57 (m, 1H), 1.72 – 1.58 (m, 2H), 1.46 – 1.36 (m, 1H). **¹³C NMR** (101 MHz, CDCl₃) δ 167.5, 142.4, 134.9, 131.3, 128.5, 128.5, 128.4, 128.4, 128.4, 128.3, 128.3, 126.9, 125.7, 40.0, 35.8, 31.0, 29.5, 26.5. **HRMS (ESI)** [M+Na]⁺ calcd. 290.1515, found 290.1520.


N-(13-phenyltridecyl)benzamide (from 51)



The compound was synthesized according to Condition C, using styrene and 12-benzamidododecanoic acid.

¹H NMR (400 MHz, CDCl₃) δ 7.76 – 7.74 (m, 1H), 7.45 (ddd, *J* = 26.1, 11.0, 5.9 Hz, 2H), 7.32 – 7.09 (m, 3H), 6.12 (br, 1H), 3.44 (dd, *J* = 13.1, 7.1 Hz, 1H), 2.63 – 2.56 (m, 1H), 1.61 (dt, *J* = 14.7, 7.2 Hz, 2H), 1.35 – 1.19 (m, 9H). **¹³C NMR** (101 MHz, CDCl₃) δ 167.5, 143.0, 135.0, 131.3, 128.5, 128.4, 128.4, 128.3, 128.2, 127.8, 126.8, 125.6, 125.5, 45.6, 40.1, 36.0, 31.5, 29.7, 29.6, 29.6, 29.6, 29.6, 29.5, 29.3, 27.5, 27.0. **HRMS (ESI)** [M+H]⁺ calcd. 380.2948, found 380.2942.


N-(13-(pyridin-2-yl)tridecyl)benzamide (from 52)



The compound was synthesized according to Condition A, using 2-vinylpyridine and 12-benzamidododecanoic acid.

¹H NMR (400 MHz, CDCl₃) δ 8.52-8.50 (m, 1H), δ 7.76-7.74 (m, 2H), δ 7.57 (td, *J* = 7.7, 1.8 Hz, 1H) δ 7.50-7.46 (m, 1H), δ 7.43-7.40 (m, 1H), δ 7.14-7.12 (m, 1H), δ 7.10-7.06 (m, 1H), δ 6.16 (br, 1H), δ 3.45 (dd, *J* = 13.1, 7.1 Hz, 2H), δ 2.78 (t, *J* = 7.7 Hz, 2H), δ 1.72 (dt, *J* = 15.4, 7.6 Hz, 2H), δ 1.61 (dt, *J* = 14.8, 7.3 Hz, 2H), δ 1.37-1.23 (m, 18H). **¹³C NMR** (101 MHz, CDCl₃) δ 167.5, 162.6, 149.2, 136.2, 135.0, 131.3, 128.5, 126.8, 122.7, 120.8, 40.1, 38.5, 29.9, 29.7, 29.6, 29.6, 29.5, 29.5, 29.4, 29.3, 27.0. **HRMS (ESI)** [M+H]⁺ calcd. 381.2900, found 381.2891.


Methyl 14-benzamido-2-methyltetradecanoate (**53**)



The compound was synthesized according to Condition A, using methyl methacrylate and 12-benzamidododecanoic acid.

¹H NMR (400 MHz, CDCl₃) δ 7.78 – 7.75 (m, 2H), 7.48 – 7.39 (m, 3H), 6.28 (br, 1H), 3.66 (s, 3H), 3.46 – 3.41 (m, 2H), 2.47 – 2.38 (m, 1H), 1.67 – 1.57 (m, 3H), 1.42 – 1.25 (m, 19H), 1.14 (d, *J* = 7.0 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 177.4, 167.5, 135.0, 131.3, 128.5, 126.8, 51.4, 40.1, 39.5, 33.8, 29.7, 29.6, 29.5, 29.5, 29.5, 29.5, 29.3, 27.2, 27.0, 17.1. HRMS (ESI) [M+H]⁺ calcd. 376.2846, found 376.2835.


3,3-diphenyl-N-(1-phenylpropan-2-yl)propan-1-amine (Prenylamine, **58**)



The compound **54** was synthesized according to Condition A, using 1,1-diphenylethelene and boc-glycin. The pure compound **54** (0.2 mmol) was first dissolved in DCM (2 mL). Then trifluoroacetic acid (20% V/V) was added dropwise to the solution and continued stirring for 5 h. After completion of the reaction (monitored with TLC), the solvent was evaporated using vacuum. After that, the crude mixture (product **56**) was then used in the next-step without purification. To the solution of **56** in DCM, AcOH (1 equiv.) followed by NaBH(OAc)₃ (1.4 equiv.) were added under N₂ atmosphere. The reaction mixture was left to stir for 36h and then the reaction mixture was quenched with H₂O and NaHCO₃. The aqueous solution was extracted with 5.0 mL DCM for three times. The organic layer was then dried over Na₂SO₄ and concentrated in vacuo. The reaction mixture was directly loaded onto a short silica gel column, followed by gradient elution with EtOAc/Heptane (20% - 50%). Removing the solvent in vacuo, afforded product prenylamine (**58**).

¹H NMR (400 MHz, CDCl₃) δ 8.02 (s, 1H), 7.26 (ddd, *J* = 9.8, 6.7, 3.4 Hz, 3H), 7.22 – 7.07 (m, 10H), 7.04 (d, *J* = 6.4 Hz, 2H), 3.92 (t, *J* = 8.0 Hz, 1H), 3.22 – 3.13 (m, 1H), 3.10 (dd, *J* = 13.1, 4.1 Hz, 1H), 2.94 – 2.71 (m, 2H), 2.55 (dd, *J* = 12.9, 10.0 Hz, 1H), 2.44 (q, *J* = 7.9 Hz, 2H), 1.06 (d, *J* = 6.4 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 162.5, 162.2, 143.2, 143.2, 136.6, 129.4, 128.9, 127.6, 127.6, 127.1, 126.8, 126.8, 55.7, 48.8, 44.2, 40.1, 32.1, 16.1. HRMS (ESI) [M+Na]⁺ calcd. 352.2036, found 352.2025.


N,N-dimethyl-3-phenyl-3-(pyridin-2-yl)propan-1-amine (Pheniramine, **59**)



The compound **55** was synthesized according to Condition A, using 2-vinylpyridine and boc-glycin. The pure compound **55** (0.2 mmol) was first dissolved in DCM (2 mL). Then trifluoroacetic acid (20% V/V) was added dropwise to the solution and continued stirring for 5h. After completion of the reaction

(monitored with TLC), the solvent was evaporated using vacuum. The crude product **57** was then without further purification was directly used to synthesize pheniramine following previous reported method.⁴ The analytical data of the compound **59** matched with the reported literature data.⁴

(8R,9S,13S,14S)-13-methyl-17-oxo-7,8,9,11,12,13,14,15,16,17-decahydro-6H-cyclopenta[a]phenanthren-3-yl 4-(3-((tert-butoxycarbonyl)amino)propyl)benzoate (60)




The compound was synthesized according to Condition A, using estrone derivative and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 8.10 (d, *J* = 8.2 Hz, 2H), 7.32 (t, *J* = 7.9 Hz, 3H), 6.97 (dd, *J* = 8.4, 2.3 Hz, 1H), 6.94 (s, 1H), 4.54 (s, 1H), 3.17 (d, *J* = 6.4 Hz, 2H), 2.98 – 2.89 (m, 2H), 2.79 – 2.67 (m, 2H), 2.58 – 2.47 (m, 1H),

2.47 – 2.38 (m, 1H), 2.37 – 2.26 (m, 1H), 2.21 – 1.93 (m, 4H), 1.91 – 1.79 (m, 2H), 1.71 – 1.47 (m, 6H), 1.45 (s, 9H), 0.92 (s, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 202.9, 165.5, 156.1, 149.1, 148.0, 138.2, 137.5, 130.5, 128.7, 127.6, 126.6, 121.9, 119.1, 50.6, 48.1, 44.4, 38.2, 36.0, 33.4, 32.5, 31.7, 31.6, 29.7, 29.6, 28.6, 26.5, 25.9, 21.7, 14.0. HRMS (ESI) [M+Na]⁺ calcd. 554.2877, found 554.2853.

Methyl 4-(3-((tert-butoxycarbonyl)amino)-1-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydronaphthalen-2-yl)propyl)benzoate (61)




The compound was synthesized according to Condition A, using bexarotene methyl ester and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 7.99 (d, *J* = 8.3 Hz, 1H), 7.29 (d, *J* = 8.3 Hz, 1H), 7.22 (s, 1H), 7.00 (s, 1H), 4.50 (s, 1H), 4.15 (s, 1H), 3.11 (d, *J* = 6.1 Hz, 1H), 2.23 (d, *J* = 6.8 Hz, 1H), 2.16 (s, 1H), 1.66 (s, 3H), 1.42 (s, 5H), 1.29 (s, 1H), 1.25 (s, 3H), 1.23 (s, 1H). ¹³C NMR (101 MHz, CDCl₃) δ 171.5, 156.0, 150.6, 143.0, 142.7, 138.01, 133.2, 130.5, 128.7, 128.5, 127.6, 124.6, 79.4, 36.3, 35.6, 35.3, 34.2, 33.9, 32.9, 32.2, 32.1, 32.0, 31.8, 28.5, 26.6, 26.5, 23.0, 19.6. HRMS (ESI) [M+H]⁺

calcd. 502.2928, found 502.2914. HRMS (ESI) [M+Na]⁺ calcd. 502.2928, found 502.2923.

Isopropyl 2-(4-(3-((tert-butoxycarbonyl)amino)-1-(4-chlorophenyl)propyl)phenoxy)-2-methylpropanoate (62)




The compound was synthesized according to Condition A, using protected fenofibrate and boc-glycine.

¹H NMR (400 MHz, CDCl₃) δ 7.23 (d, *J* = 8.4 Hz, 2H), 7.12 (d, *J* = 8.5 Hz, 2H), 7.04 (d, *J* = 8.6 Hz, 2H), 6.76 (d, *J* = 8.6 Hz, 2H), 5.05 (hept, *J* = 6.3 Hz, 1H), 4.47 (s, 1H), 3.86 (t, *J* = 7.8 Hz, 1H), 3.03 (d, *J* = 6.5 Hz, 2H), 2.16 (dd, *J* = 14.5, 7.5 Hz, 2H), 1.42 (s, 9H), 1.19 (d, *J* = 6.3 Hz, 6H). ¹³C NMR (101 MHz, CDCl₃) δ 173.8, 156.0,

154.3, 143.2, 137.3, 132.1, 129.2, 128.8, 128.4, 119.3, 79.3, 69.0, 47.5, 39.4, 36.0, 28.5, 25.5, 25.5, 21.7. HRMS (ESI) [M+Na]⁺ calcd. 350.1129, found 350.1122.

(8R,9S,13S,14S)-13-methyl-17-oxo-7,8,9,11,12,13,14,15,16,17-decahydro-6H-cyclopenta[a]phenanthren-3-yl 4-(5-((tert-butoxycarbonyl)amino)pentyl)benzoate (63)




The compound was synthesized according to Condition A, using estrone derivative and 4-benzamidobutanoic acid.

¹H NMR (400 MHz, CDCl₃) δ 8.09 (d, *J* = 8.2 Hz, 2H), 7.31 (dd, *J* = 14.1, 8.3 Hz, 3H), 7.02 – 6.87 (m, 2H), 4.49 (s, 1H), 3.18 – 3.02 (m, 2H), 3.00 – 2.87

(m, 2H), 2.70 (t, *J* = 7.6 Hz, 2H), 2.51 (dd, *J* = 18.8, 8.6 Hz, 1H), 2.47 – 2.38 (m, 1H), 2.37 – 2.25 (m, 1H), 2.21 – 1.92 (m, 4H), 1.76 – 1.47 (m, 12H), 1.44 (s, 9H), 0.91 (d, *J* = 8.5 Hz, 3H). **¹³C NMR** (101 MHz, CDCl₃) δ 202.9, 165.6, 156.1, 149.1, 148.9, 138.2, 137.5, 130.4, 128.7, 127.4, 126.6, 121.9, 119.1, 50.6, 48.1, 44.4, 40.6, 38.2, 36.1, 36.0, 35.6, 31.7, 30.9, 30.1, 29.6, 28.6, 26.5, 26.5, 25.9, 21.7, 14.0. **HRMS (ESI)** [M+Na]⁺ calcd. 582.3190, found 582.3190.

4-(13-benzamido-1-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydronaphthalen-2-yl)tridecyl)benzoic acid (64)




The compound was synthesized according to Condition A, using bexarotene methyl ester and 12-benzamidododecanoic acid acid.

¹H NMR (400 MHz, CDCl₃) δ 7.98 (d, *J* = 8.3 Hz, 2H), 7.81 – 7.69 (m, 2H), 7.52 – 7.44 (m, 1H), 7.44 – 7.37 (m, 2H), 7.31 – 7.21 (m, 3H), 6.99 (s, 1H), 6.14 (s, 1H), 4.04 (t, *J* = 7.5 Hz, 1H), 3.50 – 3.39 (m, 2H), 2.15 (s, 3H), 1.98 (dt, *J* = 22.0, 13.0 Hz, 2H), 1.69 – 1.55 (m, 6H), 1.41 – 1.16 (m, 30H). **¹³C NMR** (101 MHz, CDCl₃) δ 171.0, 167.8, 151.8, 142.7, 142.4, 140.0, 138.9, 135.0, 133.3, 131.5, 130.3, 128.7, 128.6,

128.5, 127.1, 127.0, 125.2, 124.6, 119.9, 47.4, 40.3, 36.3, 35.4, 34.2, 33.9, 32.2, 32.1, 32.0, 31.9, 31.0, 29.8, 29.7, 29.7, 29.6, 29.5, 29.5, 28.0, 27.1, 19.6. **HRMS (ESI)** [M+Na]⁺ calcd. 646.4231, found 646.4231.

N-(5,5-diphenylpentyl)nicotinamide (65)




The compound was synthesized according to Condition A, using 1,1-diphenylethylene and picamilon.

¹H NMR (400 MHz, CDCl₃) δ 8.88 (s, 1H), 8.66 (d, *J* = 3.4 Hz, 1H), 8.01 (d, *J* = 7.9 Hz, 1H), 7.31 (dd, *J* = 7.7, 4.9 Hz, 1H), 7.28 – 7.19 (m, 8H), 7.18 – 7.12 (m, 2H), 6.37 (s, 1H), 3.88 (t, *J* = 7.8 Hz, 1H), 3.39 (dd, *J* = 13.1, 6.9 Hz, 2H), 2.08 (dd, *J* = 15.5, 7.8 Hz, 2H), 1.71

– 1.56 (m, 2H), 1.42 – 1.25 (m, 2H). **¹³C NMR** (101 MHz, CDCl₃) δ 165.7, 152.1, 147.9, 145.0, 135.1, 130.6, 128.6, 127.9, 126.3, 123.5, 51.3, 40.0, 35.2, 29.5, 25.4. **HRMS (ESI)** [M+H]⁺ calcd. 345.1961, found 345.1949.


Tert-butyl (2-(4-chlorophenyl)-5,5-diphenylpentyl)carbamate (Baclofen, 66)



The compound was synthesized according to Condition A, using 1,1-diphenylethylene and 4-((tert-butoxycarbonyl)amino)-3-(4-chlorophenyl)butanoic acid.

¹H NMR (400 MHz, CDCl₃) δ 7.27-7.20 (m, 6H), δ 7.16-7.10 (m, 6H), 7.02 (d, *J* = 8.3 Hz, 2H), δ 4.29 (br, 1H), δ 3.80 (t, *J* = 7.8 Hz, 1H), δ 3.40-3.32 (m, 1H), δ (ddd, *J* = 13.7, 8.9, 5.1 Hz, 1H), δ 2.79-2.70 (m, 1H), δ 2.03-1.74 (m, 2H), δ 1.66-1.58 (m, 2H), δ 1.37 (s, 9H). **¹³C NMR** (101 MHz, CDCl₃) δ 155.8, 144.9, 144.4, 132.4, 129.2, 128.8, 128.4, 127.8, 127.7, 126.2, 126.2, 79.3, 51.5, 46.2, 45.8, 33.3, 31.8, 28.3. **HRMS (ESI)** [M+H]⁺ calcd. 472.2014, found 472.1998.


Tert-butyl ((1-(3,3-diphenylpropyl)cyclohexyl)methyl)carbamate (from 67)



The compound was synthesized according to Condition A, using 1,1-diphenylethylene and boc-protected Gabapentin.

¹H NMR (400 MHz, CDCl₃) δ 7.30 – 7.20 (m, 8H), 7.18 – 7.12 (m, 2H), 4.25 (s, 1H), 3.78 (t, *J* = 7.7 Hz, 1H), 3.02 (d, *J* = 6.3 Hz, 2H), 2.02 – 1.90 (m, 2H), 1.43 (s, 11H), 1.31 (d, *J* = 6.8 Hz, 4H), 1.25 – 1.12 (m, 6H). **¹³C NMR** (101 MHz, CDCl₃) δ 156.3, 145.2, 128.6, 127.9, 126.3, 79.0, 52.3, 46.8, 36.3, 35.4, 33.6, 29.1, 28.6, 26.5, 26.3, 21.9, 21.5. **HRMS (ESI)** [M+H]⁺ calcd. 408.2897, found 408.2889.


Tert-butyl (E)-(6,6-diphenylhex-1-en-1-yl)carbamate (68a)



The compound was synthesized according to Condition A, using 1,1-diphenylethylene and 2-((tert-butoxycarbonyl)amino)-2-cyclopropylacetic acid.

¹H NMR (400 MHz, CDCl₃) δ 7.31 – 7.20 (m, 8H), 7.19 – 7.12 (m, 2H), 6.38 (t, *J* = 9.6 Hz, 1H), 6.03 (d, *J* = 9.3 Hz, 1H), 4.51 (d, *J* = 7.6 Hz, 1H), 3.88 (t, *J* = 7.8 Hz, 1H), 2.06 (dd, *J* = 15.6, 7.8 Hz, 2H), 1.93 (qd, *J* = 7.3, 1.4 Hz, 2H), 1.47 (s, 9H), 1.41 (t, *J* = 5.0 Hz, 1H), 1.39 – 1.30 (m, 2H). **¹³C NMR** (101 MHz, CDCl₃) δ 152.9, 145.2, 128.6, 128.0, 126.3, 122.4, 107.9, 51.4, 35.3, 28.6, 28.4, 28.0, 25.5. **HRMS (ESI)** [M+Na]⁺ calcd. 374.2090, found 374.2082.

tert-butyl (Z)-(6,6-diphenylhex-1-en-1-yl)carbamate (68b)





The compound was synthesized according to Condition A, using 1,1-diphenylethylene and 2-((tert-butoxycarbonyl)amino)-2-cyclopropylacetic acid.



¹H NMR (400 MHz, CDCl₃) δ 7.30 – 7.18 (m, 8H), 7.18 – 7.11 (m, 2H), 6.40 (t, *J* = 11.8 Hz, 1H), 6.03 (d, *J* = 8.8 Hz, 1H), 4.94 – 4.68 (m, 1H), 3.86 (t, *J* = 7.8 Hz, 1H), 2.02 (tt, *J* = 10.0, 7.5 Hz, 4H), 1.45 (s, 9H), 1.37 – 1.24 (m, 2H). **¹³C NMR** (101 MHz, CDCl₃) δ 145.3, 128.5, 128.0, 126.2, 123.9, 109.6, 51.4, 35.3, 29.8, 28.7, 28.4. **HRMS (ESI)** [M+Na]⁺ calcd. 374.2090, found 374.2082.

8. Copies of the NMR spectra



Benzoyl glycine





12-(Benzoylamino)dodecanoic acid





4-(Benzoylamino)butanoic acid




Tert-butyl (3,3-diphenylpropyl)carbamate (from 3)





Tert-butyl (3,3-bis(4-fluorophenyl)propyl)carbamate (from 4)





Tert-butyl (3-phenylpropyl)carbamate (from 5)





Tert-butyl (3-(4-bromophenyl)propyl)carbamate (from 6)





Tert-butyl (3-(4-(trifluoromethyl)phenyl)propyl)carbamate (from 7)





Tert-butyl (3-(4-cyanophenyl)propyl)carbamate (from 8)





Tert-butyl (3-(4-(tert-butyl)phenyl)propyl)carbamate (from 9)





Tert-butyl (3-(4-(trimethylsilyl)phenyl)propyl)carbamate (from 10)





Tert-butyl (3-([1,1'-biphenyl]-4-yl)propyl)carbamate (from 11)





Tert-butyl (3-(2-chlorophenyl)propyl)carbamate (from 12)





Tert-butyl (3-mesitylpropyl)carbamate (from 13)





Tert-butyl (3-(perfluorophenyl)propyl)carbamate (from 14)





Tert-butyl (3-phenylbutyl)carbamate (from 15)





Tert-butyl (3-(4-chlorophenyl)butyl)carbamate (from 16)





Tert-butyl (3-(9H-fluoren-2-yl)butyl)carbamate (from 17)





Tert-butyl (3-(naphthalen-2-yl)butyl)carbamate (from 18)





Tert-butyl (2,3,3-triphenylpropyl)carbamate (from 19)





Tert-butyl (2-cyano-3,3-diphenylpropyl)carbamate (from 20)





Tert-butyl (2-cyano-3-phenylpropyl)carbamate (from 21)





Tert-butyl (2-methyl-3-phenylpropyl)carbamate (from 22)





Methyl 3-((tert-butoxycarbonyl)amino)-2-(4-chlorobenzyl)propanoate (from 23)





Methyl 3-((tert-butoxycarbonyl)amino)-2-(4-methylbenzyl)propanoate (from 24)





Tert-butyl (2,3-diphenylpropyl)carbamate (from 25)





Tert-butyl (2-phenyl-2-(phenylamino)ethyl)carbamate (from 26)





Tert-butyl (3-(pyridin-2-yl)propyl)carbamate (from 27)





Tert-butyl (3-phenyl-3-(pyridin-2-yl)propyl)carbamate (from 28)





Tert-butyl (2-(benzo[b]thiophen-2-yl)ethyl)carbamate (from 29)





Tert-butyl (3-(4-methylthiazol-5-yl)propyl)carbamate (from 30)





Ethyl 4-((tert-butoxycarbonyl)amino)butanoate (from 31)





Adamantan-1-yl 4-((tert-butoxycarbonyl)amino)butanoate (from 32)





Tert-butyl (4-(dimethylamino)-4-oxobutyl)carbamate (from 33)





Tert-butyl (2-methyl-4-oxopentyl)carbamate (from 34)





Methyl 4-((tert-butoxycarbonyl)amino)-2-methylbutanoate (from 35)





Dimethyl 2-((tert-butoxycarbonyl)amino)ethylsuccinate (from 36)





Dimethyl 2-(((tert-butoxycarbonyl)amino)methyl)succinate (from 37)





Methyl 3-(((tert-butoxycarbonyl)amino)methyl)-5-methylhexanoate (from 38)





Tert-butyl (3-(phenylsulfonyl)propyl)carbamate (from 39)





Tert-butyl ((2-methyl-3-oxo-5-(prop-1-en-2-yl)cyclohexyl)methyl)carbamate (from 40)





Tert-butyl (3-cyanopropyl)carbamate (from 41)





Tert-butyl (4,4-diphenylbutyl)carbamate (from 42)





Tert-butyl (5,5-diphenylpentyl)carbamate (from 43)





Tert-butyl (6,6-diphenylhexyl)carbamate (from 44)





Tert-butyl (7,7-diphenylheptyl)carbamate (from 45)





Tert-butyl (2-((2-((3,3-diphenylpropyl)amino)-2-oxoethyl)amino)-2-oxoethyl)carbamate (from 46)





Tert-butyl (2-hydroxy-5,5-diphenylpentyl)carbamate (from 47)





N-(5,5-diphenylpentyl)benzamide (from 48)





N-(12,12-diphenyldodecyl)benzamide (from 49)





N-(5-phenylpentyl)benzamide (50)




N-(13-phenyltridecyl)benzamide (from 51)




N-(13-(pyridin-2-yl)tridecyl)benzamide (from 52)




Methyl 14-benzamido-2-methyltetradecanoate (53)





3,3-diphenyl-N-(1-phenylpropan-2-yl)propan-1-amine (Prenylamine, 58)







(8R,9S,13S,14S)-13-methyl-17-oxo-7,8,9,11,12,13,14,15,16,17-decahydro-6H-cyclopenta[a]phenanthren-3-yl 4-(3-((tert-butoxycarbonyl)amino)propyl)benzoate (60)







methyl 4-(3-((tert-butoxycarbonyl)amino)-1-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydronaphthalen-2-yl)propyl)benzoate (61)







isopropyl 2-(4-(3-((tert-butoxycarbonyl)amino)-1-(4-chlorophenyl)propyl)phenoxy)-2-methylpropanoate (62)







(8R,9S,13S,14S)-13-methyl-17-oxo-7,8,9,11,12,13,14,15,16,17-decahydro-6H-cyclopenta[a]phenanthren-3-yl 4-(5-((tert-butoxycarbonyl)amino)pentyl)benzoate (63)







4-(13-benzamido-1-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydronaphthalen-2-yl)tridecyl)benzoic acid (64)







N-(5,5-diphenylpentyl)nicotinamide (65)







Tert-butyl (2-(4-chlorophenyl)-5,5-diphenylpentyl)carbamate (66)







Tert-butyl ((1-(3,3-diphenylpropyl)cyclohexyl)methyl)carbamate (67)







Tert-butyl (E)-(6,6-diphenylhex-1-en-1-yl)carbamate (68a)





Tert-butyl (Z)-(6,6-diphenylhex-1-en-1-yl)carbamate (68b)





9. References

- 1 MESAİK, M. A. *et al.* Synthesis and immunomodulatory properties of selected oxazolone derivatives. *Bioorganic & Medicinal Chemistry* **12**, 2049-2057, doi:<https://doi.org/10.1016/j.bmc.2004.02.034> (2004).
- 2 Yang, H. *et al.* Organic photoredox catalytic decarboxylative cross-coupling of gem-difluoroalkenes with unactivated carboxylic acids. *Organic Chemistry Frontiers* **6**, 2365-2370, doi:10.1039/C9QO00495E (2019).
- 3 Luo, J. & Zhang, J. Donor–Acceptor Fluorophores for Visible-Light-Promoted Organic Synthesis: Photoredox/Ni Dual Catalytic C(sp³)–C(sp²) Cross-Coupling. *ACS Catalysis* **6**, 873-877, doi:10.1021/acscatal.5b02204 (2016).
- 4 Yin, Y. *et al.* Conjugate Addition–Enantioselective Protonation of N-Aryl Glycines to α -Branched 2-Vinylazaarenes via Cooperative Photoredox and Asymmetric Catalysis. *Journal of the American Chemical Society* **140**, 6083-6087, doi:10.1021/jacs.8b01575 (2018).