

















a




b




c




d




e




f




a




b








c











Supplementary Figure 5











Supplementary Figure 6



Supplementary Figure 7





Supplementary Figure 9

Supplementary Legends

Supplementary figure 1. mRNA expression of *MNX1* determined by quantitative real time qPCR, normalized to *Hprt* and represented as relative expression from both *invitro* FL cells (a) and adult bone marrow cells (b). Western blot analysis of HA in the transfected FL cells. Actin was used as loading control (c). Expected size for MNX1 (41-48 kDa) and MNX1::ETV6 fusion (72 kDa).

Supplementary figure 2. Kaplan-Meier survival curves of (C57BL/6) mice transplanted with either FL (a) cells or ABM (c) cells after lethal radiation (8.5 Gy) with rescue bone marrow. Cells were retrovirally transfected with either ectopic expression of *MNX1*, *ETV6*, *MNX1::ETV6* fusion or empty vector (Ctrl). Mice were euthanized at the end of the experiment and were analyzed for spleen weight, white blood cell number and hemoglobin concentration (HGB) (b,d). Kaplan-Meier survival curves of (C57BL/6) mice transplanted with FL cells after sub-lethal radiation (5.5 Gy) with no rescue bone marrow (e). Mice were euthanized at the end of the experiment and were analyzed for spleen weight, white blood cell number and hemoglobin concentration (f). Results of control ($n \geq 6$) and transfected mice ($n \geq 6$) were analyzed using the log-rank test (ns: non-significant).

Supplementary figure 3. Percentage of GFP/YFP chimerism in peripheral blood of (C57BL/6) mice transplanted with FL cells after lethal radiation (8.5 Gy) with rescue bone marrow, as determined by flowcytometry every two weeks (a). Percentage of MNX1 engraftment in bone marrow of (C57BL/6) in mice transplanted after lethal radiation (8.5 Gy) with rescue bone marrow versus mice transplanted after sublethal radiation (5.5 Gy) with no rescue BM, as determined by GFP/YFP in bone marrow of the euthanized mice with flowcytometry (b). Quantification of flow cytometry analysis of BM from (C57BL/6) mice

transplanted with FL cells after sub-lethal radiation (5.5 Gy) with no rescue bone marrow with the indicated antibodies (c). Data represent mean $\pm$ SD of at least three experiments. Kaplan-Meier survival curves of NOD.Cg-Kit^{W-41J} Tyr⁺ Prkdc^{scid} Il2rg^{tm1Wjl}/ThomJ (NBSGW) mice transplanted with *MNX1*, *MNX1::ETV6* fusion or empty vector after lethal radiation (1.6 Gy) with rescue BM (d, left panel), or after sublethal radiation (0.9 Gy) with no rescue BM (d, middle, panel). Mice from sublethal radiation experiment were euthanized at the end and were analyzed for spleen weight, white blood cell number and hemoglobin concentration (d, right panel). Results of control ($n \geq 6$) and transfected mice ($n \geq 3$) were analyzed using the log-rank test. Kaplan-Meier survival curves of (NBSGW) mice transplanted with either primary FL cells retrovirally transfected with *MNX1* (red) or secondary transplanted with BM from leukemia mice (purple) and no radiation (e). Kaplan-Meier survival curves of (NBSGW) mice secondary transplanted with BM after sublethal radiation (0.9 Gy) (f, left panel). Mice were euthanized at the end of the experiment and were analyzed for spleen weight, white blood cell number and hemoglobin concentration (f, right panel).

Supplementary figure 4. Kaplan-Meier survival curves of (NBSGW) mice transplanted with either FL cells with either ectopic expression of *MNX1*, *ETV6*, *MNX1::ETV6* fusion or empty vector after Sub-lethal radiation (0.9 Gy) with no rescue bone marrow (a). Results of control ($n \geq 5$) and transfected mice ($n \geq 3$) were analyzed using the log-rank test (ns: non-significant). Mice were euthanized at the end of the experiment and were analyzed for spleen weight, white blood cell number and hemoglobin concentration (b). Quantification of flow cytometry analysis of the euthanized mice at the end of the experiment with the indicated antibodies (c). Data represent mean $\pm$ SD of at least three experiments and is considered significant using two sided student t-test, non-significant (ns) at $p > 0.05$.

Supplementary figure 5. Quantification of flow cytometry analysis of FL cells (a) or ABM cells (b-d) transfected with ectopic expression of *MNX1*, *ETV6*, *MNX1::ETV6* fusion or empty

vector with the indicated antibodies. Data represent mean $\pm$ SD of at least three experiments. CFU (Colony forming unit assay) of transfected FL cells showing different types of colonies after 2 weeks (e). Bright-field microscopy image of FL cells transfected with *MNX1* or empty vector (Ctrl), showing number of cells after 5 days (f).

Supplementary figure 6. mRNA expression of *MNX1* determined by quantitative real time qPCR, normalized to *Hprt* and represented as relative expression from FL cells (a). Most enriched pathways for differentially expressed genes of leukemia BM cells with *MNX1* ectopic expression in comparison with empty vector using gene-ontology biological process pathways (b). LogFC heat map of downregulated (blue) and upregulated (red) differentially expressed genes of leukemia BM cells with *MNX1* ectopic expression in comparison with BM from mice transplanted with empty vector (Ctrl). Differentially expressed genes were selected based on a LogFC1 and FDR $\leq$ 0.05 (c). PCA scatter plot of normalized gene expression using variance Stabilizing Transformation (d). Gene set enrichment analysis (GSEA) using the Gene ontology biological pathways (GO) gene set showing normalized enrichment score for pathways with NOM p-value $\leq$ 0.05 (e)

Supplementary figure 7. Immunofluorescence of ABM cells stained with γ H2AX –FITC antibody (green) and counterstained with DAPI (blue) (left panel, a). Quantification of the number of γ H2AX foci/cell. At least 50 cells were counted (right panel, a). Flowcytometry analysis of FL and ABM cells after staining with DAPI for cell cycle distribution (b). Median DNA values at G1 phase and G2 phase of cell cycle for flowcytometry analysis of BM from leukemia NBSGW mice transfected with *MNX1*-FL cells in comparison with NBSGW mice transfected with empty vector Ctrl (c). Quantification of the binding of H3K4me3 and H3K27me3 to the promoter regions of *Cep164* in FL cells with either *MNX1* or empty vector

determined by ChIP-qPCR (d). Western blot analysis of H3K4me1, H3K4me2, H3K4me3 and H3K27me3 in the BM of leukemia mice transfected with *MNX1* ectopic expression in comparison with FL cells transfected with empty vector. Actin was used as loading control (e). The binding of Histone H3 was detected by immunoprecipitation of MNX1 using HA-antibody followed by western blot using histone H3 antibody in FL cells (upper panel ,f) or BM of leukemia mice transfected with *MNX1* ectopic expression (lower panel, f) in comparison to FL cells transfected with empty vector (ctrl). ETV6 with HA tag was used as a negative control. Total protein input was used as loading control (upper panel, f). Most enriched pathways for differentially accessible regions of leukemia BM cells with *MNX1* ectopic expression in comparison with FL cells with empty vector using gene-ontology biological process pathways (g). M: MNX1, C: Control.

Supplementary figure 8. In-silico predicted transcription factor binding sites by DIFFTF package from ATAC-Seq showing top differentially activated (FDR < 0.05) transcription factors (TFs) binding between MNX1 BM from leukemia mice and FL Ctrl cells (a). Correlation map between ATAC-Seq and RNA-Seq expression analysis (b). The binding of Mat2a was detected by immunoprecipitation of MNX1 using HA-antibody followed by western blot using Mat2a antibody in BM of leukemia mice transfected with *MNX1* ectopic expression in comparison to FL cells transfected with empty vector. Total protein input was used as loading control (c). Western blot analysis of Mat2a, Ahcy and Histone H3 in the FL cells transfected with *MNX1*, *ETV6* or *MNX1::ETV6* fusion in comparison with FL cells transfected with empty vector (ctrl). Actin was used as loading control (d). Various amino acid concentration was determined by LC-ESI-MSMS from leukemia BM cells transfected with MNX1 ectopic expression normalized against total protein concentration and represented as fold difference relative to control (e). SAH concentration was determined by ELISA from FL cells expressing MNX1 (Invitro-FL) and leukemia BM cells (leukemia mice BM), normalized

against total protein concentration and represented as fold difference relative to control (f). Recombinant Histone H3.1& H3.3 were subjected to an invitro methyltransferase reaction using MNX1 complex pulled down with HA antibody from FL cells transfected with MNX1 (M), in comparison with FL cells with empty vector (C) in the presence of SAM and DTT. The reactions were terminated by boiling in SDS sample buffer. Separation of samples in 12% SDS-PAGE was followed by immunoblotting (IB) with H3K4me1 antibody. Reblotting was made for detection of HA and total Histone H3. As indicated, negative controls were obtained by omitting Histone H3, or pulled down immune complex (upper panel, g). Quantification of the invitro methyltransferase reaction (lower panel, g). Data represent mean $\pm$ SD of at least three experiments. M: MNX1, C: Control.

Supplement figure 9. ImageJ quantification of the western blot analysis of H3K4me1, H3K4me2, H3K4me3 and H3K27me3 after treatment as indicated (a). Immunofluorescence of FL cells stained with γ H2AX –FITC antibody (green) and counterstained with DAPI (blue) after treatment with vehicle or 5 μ M Sinefungin (MNX1+S) (b). Quantification of the flowcytometry analysis of FL cells with 5 μ M Sinefungin (Ctrl+S, MNX+S) or without treatment (Ctrl, MNX1) after staining with DAPI for cell cycle distribution (c). Percentage of MNX1 engraftment in bone marrow of (NBSGW) in mice transplanted with FL cells, with 5 μ M Sinefungin (Ctrl+S, MNX+S) or without treatment (Ctrl, MNX1), as determined by GFP/YFP in bone marrow of the euthanized mice with flowcytometry (d). Mice were euthanized at the end and were analyzed for spleen weight, white blood cell number and hemoglobin concentration (e). mRNA expression of *MNX1* determined by quantitative real time qPCR, normalized to *Hprt* and represented as relative expression from FL cells with 5 μ M Sinefungin (Ctrl+S, MNX+S) or without treatment (Ctrl, MNX1) (f). Percentage of GFP/YFP chimerism in peripheral blood of (NBSGW) mice transplanted with FL with 5 μ M Sinefungin (Ctrl+S, MNX+S) or without treatment (Ctrl, MNX1), as determined by flowcytometry every

two weeks (g). LogFC heat map of downregulated (blue) and upregulated (red) differentially expressed genes of leukemia BM cells with *MNX1* ectopic expression in comparison with FL cells with empty vector (Ctrl), with 5 μ M Sinefungin (Ctrl+S, MNX+S) or without treatment (Ctrl, MNX1). Differentially expressed genes were selected based on a LogFC1.5 and FDR $\leq$ 0.05 (h). Data represent mean $\pm$ SD of at least three experiments

Supplementary Table .2

Plasmid sequences

HA-MNX1-ETV6 fusion

(MNX1 exon 1 is mouse codon optimized)

```
ATGGAGGCCTATCCTTACGATGTGCCTGATTATGCACTCTGGTAGTACTGCGCAGTCGA  
CGGAAAAGAGCAAGAACCTTTAGAATAGACGCCCTTTTGGCTGTCGACCCACCTC  
GCGCAGCTAGTGACAAAGCGCTCCTCTGGCCTTGGTTACATCCCTGGCTGCTGC  
CGCATCTGGAACCGGTGGCGGAGGCGGGGGCGGCGGAGCCTCTGGGGGGACTTC  
CGGTAGCTGTAGTCCTGCAAGTTCCGAACCACAGCCGCCCCAGCCGACCGCCTG  
CGGGCCGAGAGCCCATCTCCACCAAGGCTGTTGGCTGCTCACTGTGCCTTGCTCC  
CAAAACCCGGATTCTCTGGCGCAGGCGGAGGAGGGGGCGGGACTGGAGGAGGA  
CACGGAGGCCCTCATCACCATGCTCATCCTGGCGCAGCAGCTGCAGCAGCTGCTG  
CCGCAGCTGCCGCCGCAGCAGGGGGCCTGGCTCTGGGCCTGCACCCCGGTGGCG  
CTCAGGGCGGAGCTGGCCTGCCGGCTCAGGCCGCACTTTACGGCCACCCGGTCTA  
TGGATACAGCGCTGCCGCCGCCGCCGCCGCTTGGCTGGACAACACCCAGCACTT  
TCTTACAGTTATCCACAGGTGCAGGGAGCCCACCCGGCCCCATCCCGCAGACCCTA  
TTAAACTGGGGGCTGGGACGTTCCAGCTGGACCAGTGGCTCAGAGCATCCACTG  
CTGGGATGATCCTGCCCAAGATGCCTGACTTCAACCTGCTTGCAGCCAATTTACTG  
GAGCAGGGATGACGTAGCCCAGTGGCTCAAGTGGGCTGAAAATGAGTTTTCTTT  
AAGGCCAATTGACAGCAACACGTTTGAAATGAATGGCAAAGCTCTCCTGCTGCT  
GACCAAAGAGGACTTTTCGCTATCGATCTCCTCATTGAGGTGATGTGCTCTATGAA  
CTCCTTCAGCATATTCTGAAGCAGAGGAAACCTCGGATTCTTTTTTCACCATTCTT  
CCACCCTGGAACTCTATACACACACAGCCGGAGGTCATACTGCATCAGAACCA  
TGAAGAAGATAACTGTGTCCAGAGGACCCCCAGGCCATCCGTGGATAATGTGCA  
CCATAACCCTCCCACCATTGAACTGTTGCACCGCTCCAGGTCACCTATCACGACA  
AATCACCGGCCTTCTCCTGACCCCGAGCAGCGGCCCTCCGGTCCCCCTGGACA  
ACATGATCCGCCGCTCTCCCCGGCTGAGAGAGCTCAGGGACCCAGGCCGCACC  
AGGAGAACAACCACAGGAGTCTTACCCTCTGTCACTGTCTCCCATGGAGAATA  
ATCACTGCCCAGCGTCTTCCGAGTCCCACCCGAAGCCATCCAGCCCCCGGCAGG  
AGAGCACACGCGTGATCCAGCTGATGCCAGCCCCATCATGCACCCTCTGATCCT  
GAACCCCCGGCACTCCGTGGATTTCAAACAGTCCAGGCTCTCCGAGGACGGGCT  
GCATAGGGAAGGGAAGCCCATCAACCTCTCTCATCGGGAAGACCTGGCTTACAT  
GAACCACATCATGGTCTCTGTCTCCCCGCTGAAGAGCACGCCATGCCATTGGG  
AGAATAGCAGACTGTAGACTGCTTTGGGATTACGTCTATCAGTTGCTTTCTGACA  
GCCGGTACGAAAACCTTCATCCGATGGGAGGACAAAGAATCCAAAATATTCCGGA  
TAGTGGATCCCAACGGACTGGCTCGACTGTGGGGAAACCATAAGAACAGAACAA  
ACATGACCTATGAGAAAATGTCCAGAGCCCTGCGCCACTACTACAACTAAACA  
TTATCAGGAAGGAGCCAGGACAAAGGCTTTTGTTTCAGGTTTATGAAAACCCAG  
ATGAAATCATGAGTGGCCGAACAGACCGTCTGGAGCACCTAGAGTCCCAGGAGC  
TGGATGAACAAATATACCAAGAAGATGAATGCTGA
```

HA-ETV6

ATGGAGGCCTATCCTTACGATGTGCCTGATTATGCACTCTGGTAGTACTGCGCAGTCGACGCTCTGAGACTCCTGCTCAGTGTAGCATTAAGCAGGAACGAATTTTCATATACACCTCCAGAGAGCCCAGTGCCGAGTTACGCTTCCTCGACGCCACTTCATGTTCCAGTGCTCGAGCGCTCAGGATGGAGGAAGACTCGATCCGCCTGCCTGCGCACCTGCGCTTGCAGCCAATTTACTGGAGCAGGGATGACGTAGCCCAGTGGCTCAAGTGGGCTGAAAATGAGTTTTCTTTAAGGCCAATTGACAGCAACACGTTTGAAATGAATGGCAAAGCTCTCCTGCTGCTGACCAAAGAGGACTTTTCGCTATCGATCTCCTCATTTCAGGTGATGTGCTCTATGAACTCCTTCAGCATATTCTGAAGCAGAGGAAACCTCGGATTCTTTTTTTCACCATTCTTCCACCCTGGAACTCTATACACACACAGCCGGAGGTCACTATGCATCAGAACCATGAAGAAGATAACTGTGTCCAGAGGACCCCCAGGCCATCCGTGGATAATGTGCACCATAACCCTCCCACCATTGAACTGTTGCACCGCTCCAGGTCACCTATCACGACAAATCACCGGCCTTCTCCTGACCCCGAGCAGCGGCCCTCCGGTCCCCCCTGGACAACATGATCCGCCGCTCTCCCCGGCTGAGAGAGCTCAGGACCCAGGCCGACCCAGGAGACAACACCAGGAGTCCTACCCTCTGTCAGTGTCTCCCATGGAGAATAATCACTGCCCAGCGTCCTCCGAGTCCCACCCGAAGCCATCAGCCCCCGGCAGGAGAGCACACGCGTGATCCAGCTGATGCCCAGCCCCATCATGCACCCTCTGATCCTGAACCCCCGGCACTCCGTGGATTTCAAACAGTCCAGGCTCTCCGAGGACGGGCTGCATAGGGAAGGGAAGCCCATCAACCTCTCTCATCGGGGAACCTGGCTTACATGAACCACATCATGGTCTCTGTCTCCCCGCCTGAAGAGCACGCCATGCCCATTTGGGAGAATAGCAGACTGTAGACTGCTTTGGGATTACGTCTATCAGTTGCTTTCTGACAGCCGGTACGAAAACCTTCATCCGATGGGAGGACAAAGAATC CAAAATATTCCGGATAGTGGATCCCAACGGACTGGCTCGACTGTGGGGAAACCA TAAGAACAGAACAAACATGACCTATGAGAAAATGTCCAGAGCCCTGCGCCACTA CTACAAACTAAACATTATCAGGAAGGAGCCAGGACAAAGGCTTTTGTTCAGGTT TATGAAAACCCAGATGAAATCATGAGTGGCCGAACAGACCGTCTGGAGCACCT AGAGTCCCAGGAGCTGGATGAACAAATATACCAAGAAGATGAATGCTGA

HA-MNX1

ATGGAGGCCTATCCTTACGATGTGCCTGATTATGCACTCTGGTAGTACTGCGCAGTCGACGGAAGAGCAAGAACTTTAGAAATAGACGCCCTTTTGGCTGTGCGACCCACCTGCGCAGCTAGTGACAAAGCGCTCCTCTGGCCTTGGTTACATCCCTGGCTGCTGCGCATCTGGAACCGGTGGCGGAGGCGGGGGCGGCGGAGCCTCTGGGGGGGACTTC CGGTAGCTGTAGTCCTGCAAGTTCCGAACCACCAGCCGCCCCAGCCGACCGCCTG CGGGCCGAGAGCCCATCTCCACCAAGGCTGTTGGCTGCTCACTGTGCCTTGCTCC CAAAACCCGGATTCTCTGGCGCAGGCGGAGGAGGGGGCGGGACTGGAGGAGGGA CACGGAGGCCCTCATCACCATGCTCATCCTGGCGCAGCAGCTGCAGCAGCTGCTG CCGCAGCTGCCGCCGAGCAGGGGGCCTGGCTCTGGGCCTGCACCCCGGTGGCG CTCAGGGCGGAGCTGGCCTGCCGGCTCAGGCCGCACTTTACGGCCACCCGGTCTA TGGATACAGCGCTGCCGCCGCGCCGCCGCTTGGCTGGACAACACCAGCACTT TCTTACAGTTATCCACAGGTGCAGGGAGCCCACCCGGCCCATCCCGCAGACCCTA TTAAACTGGGGGCTGGGACGTTCCAGCTGGACCAGTGGCTCAGAGCATCCACTG

CTGGGATGATCCTGCCCCAAGATGCCTGACTTCAACAGTCAGGCTCAGTCAAACCT
TCTGGGTAAGTGCCGACGACCACGGACAGCTTTCACATCACAGCAACTGTTGGA
ACTGGAGCACCAATTTAAGCTGAATAAGTACCTCTCCAGACCCAAGAGGTTCTGA
AGTTGCCACCAGCCTGATGCTTACTGAAACCCAAGTCAAGATATGGTTTCAGAAC
AGACGCATGAAGTGGAAGCGAAGTAAGAAGGCTAAGGAACAGGCCGCACAGGA
GGCCGAAAAGCAGAAGGGGGGAGGCGGCGGTGCCGGTAAAGGCGGTGCCGAGG
AGCCAGGAGCAGAGGAGCTGCTGGGCCCACCAGCTCCTGGCGACAAAGGGTCAG
GGAGACGGCTCCGGGATCTCCGAGACAGCGATCCAGAGGAGGACGAGGACGAG
GATGATGAAGATCACTTTCCTACTCCAACGGGGCTAGTGTTACACGCAGCCAGCT
CAGACTGTAGTAGCGAAGACGATTCCCCCCCACCTAGACCTTCTACCAACCTGC
CCCCCAATGA

HA-tag

Linker region

| Border between MNX1 exon 1 and ETV6 exon 3

Primer sequences

Primers for SYBR assays (for codon optimized transcripts)

Primer name	Sequence 5'-3'
hMNX1ex2-3 fwd	GCT TAC TGA AAC CCA AGT CAA GA
hMNX1ex2-3 rev	TTC TGC TTT TCG GCC TCC T
hETV6ex2-3 fwd	CTC GAC GCC ACT TCA TGT TC
hETV6ex2-3 rev	ATT TTC AGC CCA CTT GAG CC
hM-Efu fwd	CCT GAC TTC AAC TGC TTG CA
hM-Efu rev	TGC TGT CAA TTG GCC TTA AAG A

TaqMan Gene Expression Assays from Applied Biosystems

Gene	Assay
mHprt (ref gene)	Mm01545399_m1
mMnx1	Mm01222622_m1
mEtv6	Mm00468390_m1
mCep164	Mm00553106_m1

Primers used for ChIP-qPCR

Gene	Assay
Cep164-fwd	CCAGCAACCACTTCCGGAAA
Cep164-rev	ACCAATCCAATGCCAACGCT

Supplementary materials and methods

Western blotting

Cells were lysed in modified RIPA buffer (50 mm Tris, pH 7.4, 150 mm NaCl, 1% Nonidet P-40, 1 mm EDTA, 0.25% sodium deoxycholate) containing protease (Roche Applied Science, Mannheim, Germany) and phosphatase (Sigma) inhibitors and processed as described previously (47). Cell lysate subjected to sonication using bioruptor for 5 minutes (30 second ON/ 30 seconds OFF). Primary antibodies used in this study included the following: rabbit-anti HA # 11846, rabbit-anti actin # 4970, rabbit-anti H3K4me1 #5326, rabbit-anti H3K4me2 #9725 from Cell Signaling Technology, rabbit-anti Mat2A#NB110-94158, rabbit-anti Ahcy #NBP2-48817 from Novous biological, rabbit-anti H3K4me3 (catalog no. C15410003-50), rabbit anti-H3K27me3 (catalog no. C15410069), from Diagenode and rabbit anti-Histone 3 (catalog no. Ab18521) from Abcam.

Immunoprecipitation and mass spectrometry

Anti HA-magnetic beads from Thermo Fisher Scientific (catalog no. 88836) were used for immunoprecipitation. After blocking with 0.1% BSA in PBS, 0.7-1 mg of cell lysates were incubated with HA antibody for 1h at RT. For IgG antibody, the antibody was cross-linked to magnetic protein G Dynabeads using Dynabeads® antibody coupling kit 14311D (Invitrogen). The immune complexes were washed three times with lysis buffer and eluted by boiling in SDS sample buffer (Invitrogen). Precipitates were separated with SDS-PAGE and co-precipitated protein bands were visualized using Coomassie Blue. Prominent gel bands were excised for LC-MS using a hybrid LTQ-Orbitrap Velos mass spectrometer (Thermo Fisher Scientific) as described in ¹.

Invitro histone methyltransferase

Invitro histone methyltransferase was done as described in ². Briefly, 0.25-0.5 mg of cell lysates were incubated with HA antibody for 1h at RT. The immune complexes were washed three times with lysis buffer and once with BC100 buffer (20 mM Tris-HCl pH 8.0, 100 mM KCl, 0.2 mM EDTA, 20% glycerol, 1 mM DTT). The immune complexes was incubated with 4 µg of human recombinant histone H3.1 and H3.3 (New England biolabs # M2507S, # M2503S) at 30 °C for 60 min in 25 µl of methylation buffer buffer (50 mM Tris-HCl pH 8.5, 5 mM MgCl₂) in the presence of 10 µM SAM (New England biolabs # B9003S) and 4mM DTT. Omission of the recombinant histone H3, or the immune complexes served as negative controls. The reactions were terminated by boiling in SDS sample buffer (Invitrogen), followed by separations in 12% SDS-PAGE and immunoblottings using H3k4m31 (Cell signaling), Histone H3 (Abcam) and anti-methylated lysines (Abcam #ab23366). Quantification was done using ImageJ ³.

Histological preparation

Moribund mice or mice at the end of the experiment were euthanized. Liver, spleen, and sternum were harvested by fixing in 4% formaldehyde at 4°C, dehydrated in 70–100% ethanol, cleared in xylene and embedded in paraffin. Sections (4–5 µm). Paraffin-embedded tissue sections were prepared at Histocentre (Gothenburg, Sweden) and stained with HandE. Blood samples was collected directly from the heart after euthanizing the mice , prepared as thin blood films, Giemsa-stained and examined under the microscope (Nikon Inc., Japan).

Colony forming unit assay

Methocult M3434 (STEMCELL Technologies) was used for the CFU assays where 5000 cells were seeded/petri dish and incubated for 7-14 days according to the manufacturer's protocol.

Proliferation assay

Cell proliferation was determined using MTT (3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide) assay from abcam # ab211091. Briefly, FL cells transduced with our constructs were plated, in triplicate, in a 96-well culture plate at 37° C in CO2 incubator for 7 days. Each day an equal volume of MTT was added to the cell culture medium. The plate was incubated at 37° C for 4 h followed by addition of MTT solvent. The plate was then swirled gently until all crystals are dissolved. Absorbance values were blanked against cell culture medium without cells. The absorbance was read at 590 nm using a microplate reader (*Infinite F50*® Tecan Group Ltd., Swiss). The absorbance was plotted against the number of days

Sinefungin treatment

Sinefungin was purchased from Calbiochem #567051 as Insolution. Final doses of 5µM was used to treat the cells. Daily change of medium with fresh Sinefungin was added after viral transfection for one week.

Gene expression analysis

Total RNA was isolated using the miRNeasy Plus micro kit (Qiagen) and cDNA synthesized with SuperScript III First-Strand Synthesis SuperMix for qRT-PCR (Invitrogen/Thermo Fisher Scientific). TaqMan Gene Expression Assays and TaqMan Universal MasterMix II (Applied Biosystems/ Thermo Fisher Scientific) were used for probe based qPCR assays, listed in

supplement table 2. A SYBR Green assay was used for the MNX1, ETV6 and MNX1(ex1)::ETV6(ex3) fusion detection, with primers synthesized at IDT (Integrated DNA Technologies) and Power SYBR Green PCR Master Mix (Applied Biosystems/Thermo Fisher Scientific), listed in supplement table 2. The reactions were performed on the ABI 7900HT Sequence Detection System (Applied Biosystems/ Thermo Fisher Scientific) with *hprt* as the reference gene. Gene expression was calculated using the deltaCt method and are presented as $2^{-\text{deltaCt}}$.

Next-generation sequencing

Total RNA was extracted from transduced FL cells after one week of transfection using RepliePrep RNA miniprep systems (Promega). 5-20µg of total RNA were utilized for RNA-Seq using DNBseq technology at Beijing Genomics Institute (BGI, Hong Kong), with RIN value of ≥ 7.5 and 100bp paired-end reads. An average of 40 million reads were generated for each sample. The gene expression level was quantified by BGI Europe. Only samples with $28S/18S \geq 1$, $Q20_rate \geq 97$ and $GC\% \geq 51\%$ were used. For RNA-seq analysis, alignment was carried out to reference mouse genome GRCm38 using STAR (version 2.9.0a) with the following parameters: outSAMtype BAM SortedByCoordinate. Quantification was carried out using HTSeq-count (Python package HTSeq, python v 3.8.1) software using the following parameters -f bam -r pos -s no -m union with the reference annotation file. The mouse reference genome and annotation files were obtained from the Ensembl database. Differential gene expression was performed using the R package “Deseq2”⁴. Genes with $\log_2$ (foldchange) ≥ 1 and false discovery rate (FDR) adjusted p-value < 0.05 were considered differentially expressed genes. Principal component (PCA) analysis, heatmap analysis (pheatmap) and enhanced volcano plot (enhancedVolcano) were performed in order to examine the level of differences in the gene expression profiles between control and MNX1 samples. Gene

Ontology Go-biological process database was used for pathway analysis using the default parameters. A pathway was considered to be significantly enriched for a FDR < 0.05, p-value < 0.05 of the respective pathway. Gene Set Enrichment Analysis (GSEA) was used with the Gene ontology GO-biological process gene set. A pathway in GSEA analysis was regarded significant at nominal p-value < 0.05.

Assay for transposase-accessible chromatin using sequencing (ATACseq)

ATACseq libraries were prepared according to a modified Omni-ATAC protocol ⁵ as follows. Cells stored at -80 °C in culture medium containing 10% DMSO were thawed on ice, counted with a Countess counter (Invitrogen), pelleted at 1000 × g for 10 min at 4 °C, and a suitable aliquot for both ATACseq and antibody-guided chromatin tagmentation (ACTseq) was resuspended in cold phosphate-buffered saline (PBS) at a titer of 10⁶ viable cells/ml. An aliquot of 50000 viable cells was pelleted at 600 × g for 5 min at 4 °C, and cells were resuspended in 50 µl cold RSB containing each 0.1% of NP40 and Tween-20 and 0.01% digitonin. Proper lysis was monitored microscopically using Trypan blue, then 200 µl RSB containing 0.1% Tween-20 were mixed in. Lysed cells were spun with 1000 x g for 10 min at 4 °C and resuspended in 48 µl transposition buffer (10 mM Tris-acetate, pH 7.6, 5 mM MgCl₂, 10% dimethylformamide, 30% PBS, 0.01% digitonin and 0.1% Tween-20). The suspension was completed by addition of 2 µl Tn5 transposome generated as described by Picelli et al. ⁶. (The Tn5 transposase was kindly provided by the EMBL-Heidelberg protein expression and purification core facility). Tagmentation was performed at 37 °C for 30 minutes in a thermomixer with 1000 rpm. DNA was purified after addition of 20 µl 5 M guanidinium-thiocyanate with 140 µl HighPrep beads (Biozym) and eluted with 20 µl H₂O. Sequencing libraries were generated by addition of 25 µl NEBNext High Fidelity 2× Master Mix (NEB), each 2.5 µl of 10 µM primer Tn5mCP1n and Tn5mCBar ⁷ and 0.5 µl of 100x SYBRGreen I

using a LightCycler 480 real-time machine with 72 °C for 5 min, 98 °C for 30 sec and cycling at 98 °C, 10 sec; 63 °C, 30 sec; 72 °C, 30 sec. Amplification was stopped when an increase of about 10 fluorescence units was observed, usually between cycle 12 and 15. . Libraries were bead-purified with 70 µl HighPrep beads and eluted with 12 µl EB (Qiagen). Concentrations and fragment lengths were determined with the Qubit dsDNA High sensitivity kit (Invitrogen) and the TapeStation D1000 High sensitivity assay (Agilent), respectively. Up to ten libraries with compatible barcodes were pooled equimolarly, and pools of 5-10 nM were sequenced in the DKFZ Genomics and Proteomics Core Facility (GPCF) using an Illumina NextSeq 550 system with 75 bp paired-end, mid output mode. Sequencing reads were bioinformatically analyzed as detailed by Hey et al.⁸. In short, peak calling of processed reads was done using MACS2 callpeak v. 2.1.0.20140616 with a q-value cutoff of 0.05 and default parameters. The analysis is a publicly accessible fully containerized workflow using Common Workflow Language v. 1.0

(<https://doi.org/10.6084/m9.figshare.3115156.v2>); Highly Portable Processing of Data Generated by ChIP-seq, ChIPmentation, Cut&Run, and ACT-seq. Version 1.1.2, Zenodo: Temple, FL, USA). Differentially accessible regions (DARs) were identified with the DiffBind R package (R package version 2.12.0)⁹. Common peaks were considered those found in at least two samples. Differential analysis was performed using edgeR¹⁰. Regions with an FDR < 0.05 were considered differentially accessible. Further annotation of DARs was performed with R package ChIPseeker (R package version 1.20.0)¹¹. Differential transcription factor activity was predicted using diffTF¹² and in silico predicted transcription factor binding sites based on the HOCOMOCO 10 database were used as a reference¹³.

Antibody-guided chromatin tagmentation followed by sequencing (ACTseq)

ACTseq¹⁴ was essentially performed as detailed by Liu et al.¹⁵. As described in the ATACseq methods chapter, frozen cells were thawed, washed with PBS and, per antibody, aliquots of 50000 viable cells were lysed. Antibodies used for the pA-Tn5 transposome-antibody complexes were directed against H3K27me3 (Cell signaling, 9733), IgG (Millipore PP64B) and yeast histone H2B (M3930, Boster Bio. Tech., Pleasanton, CA, USA), the latter complexes being used for spike-in to enable sequence read normalization between biological replicates¹⁵. Sequencing libraries were generated, sequenced and bioinformatically analyzed as described in the ATACseq section.

For data processing, Trim Galore v. 0.4.4¹⁶ (https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/; accessed on 13 December 2020) was applied together with Cutadapt v. 1.14¹⁷ using “-paired”, “-nextera”, “-length_1 35”, and “-length_2 35” parameters. Trimmed reads were mapped against the Genome Reference Consortium Human Build version 37 by means of Bowtie2 v. 2.2.6¹⁸ using the “-very-sensitive” flag and a maximum insertion length of 2500 bp. Mappings belonging to the same lane-multiplexed library were combined using SAMtools merge v. 1.5¹⁹. Discordant alignments and mappings with a Phred score below 20 were removed using SAMtools²⁰, as fragments obtained from tagmentation cannot be smaller than 38 bp. Thus, all alignments corresponding to fragment sizes below that threshold were removed. Read ends were shifted to represent the center of the transposition event as previously described by²¹. Additionally, trimmed reads were aligned against the *Saccharomyces cerevisiae* R64 reference genome followed by post-alignment filtering as described above. To calculate a library-specific scaling factor, we derived the multiplicative inverse of the number of filtered reads mapped to the yeast

Cancers 2021, 13, 2455 9 of 26 genome. Coverage tracks were generated by means of the bamCoverage functionality of Deeptools v. 3.1.1²² using the non-default parameters “--ignoreForNormalization chrM chrY chrX” and “--effectiveGenomeSize 2652783500” as well as the “--scaleRatio” option to specify the spike-in-based scaling factor. The analysis procedure was implemented as fully containerized workflow using the Common Workflow Language v. 1.0. Differentially bound regions were identified as described for the ATACseq data.

Chromatin immunoprecipitation (ChIP):

ChIP assays were performed using iDeal ChIP-qPCR kit (C01010180, Diagenode). Cells were subject to fixation and crosslinking by formaldehyde (net concentration 1%) for 10-15 minutes at RT. Formaldehyde fixation is quenched by adding Glycine 125uM and incubating the fixed cellular suspension at RT for 5-10 minutes. The pellet is then washed twice by cold PBS and subjected then to lysis according to the specifications of the manufacturer. Next, cell lysates is subjected to sonication using bioruptor for 20 minutes (30 second ON/ 30 seconds OFF), so that the sheared smear is supposed to range from 200-600 bp long. The sheared chromatin is then subject for immunoprecipitation using high quality ChIP-grade antibodies against: HA (cell signaling), H3K27me3 (Diagenode), H3K4me3 (Diagenode), and polyclonal rabbit anti-IgG (c15410206, Diagenode) as negative control. Immunoprecipitation relies on that each antibody is crosslinked to protein A/G-coated magnetic beads, and the reaction takes up to 16 hours at 4°C, and then the beads crosslinked with the antibody-DNA is then subject for washing by buffers containing gradient concentrations of detergent. The washed beads-antibody-DNA is then subjected for de-crosslinking in the presence of proteinase K, so that all DNA-binding proteins, histones and beads are released, and the DNA sequences are then eluted and used for downstream ChIP-real time PCR. The final eluted DNA was analyzed by real-time quantitative PCR with the Power SYBR Green PCR master mix (Applied Biosystems). The PCR conditions

were: 50 °C for 2 min, 95 °C for 10 min, and 40 cycles of 95 °C for 15 s and 60 °C for 1 min. The PCR analysis was performed with the 7900HT fast real-time PCR system instrument and software (Applied Biosystems). All primers used to amplify the genomic target sites were designed by using PRIMER3 PLUS online tools. Primer sequences are listed in Supplement table 2.

The calculation of enrichment was normalized to the respective inputs and done according to the following formulas:

- The % of IP to input formula: $\% \text{ recovery} = 2^{[(\text{Ct input} - \log_2(5\%)) - \text{Ct sample}]} * 100\%$
- The enrichment was calculated as: $\% \text{ recovery IP} / \% \text{ recovery IgG}$

Methionine concentration:

Amino acid quantification was done at the Swedish Metabolomics Centre, Umeå, Sweden by LC-MSMS. This was done as following:

Standards and Calibration Curve

Amino acid standards (alanine, arginine, aspartic acid, cysteine, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tyrosine, valine, glutamine, asparagine, GABA, citrulline, ornithine, taurine, tryptophan, 5-HTP and kynurenine) were purchased from Sigma (St. Louis, MO, USA). Isotopically labeled amino acid standards (alanine ($^{13}\text{C}_3$, ^{15}N), arginine ($^{13}\text{C}_6$, $^{15}\text{N}_4$), aspartic acid ($^{13}\text{C}_4$, ^{15}N), cystine ($^{13}\text{C}_6$, $^{15}\text{N}_2$), glutamic acid ($^{13}\text{C}_5$, ^{15}N), glycine ($^{13}\text{C}_2$, ^{15}N), histidine ($^{13}\text{C}_6$, $^{15}\text{N}_3$), isoleucine ($^{13}\text{C}_6$, ^{15}N), leucine ($^{13}\text{C}_6$, ^{15}N), lysine ($^{13}\text{C}_6$, $^{15}\text{N}_2$), methionine ($^{13}\text{C}_5$, ^{15}N), phenylalanine ($^{13}\text{C}_9$, ^{15}N), proline ($^{13}\text{C}_5$, ^{15}N), serine ($^{13}\text{C}_3$, ^{15}N), threonine ($^{13}\text{C}_4$, ^{15}N), tyrosine ($^{13}\text{C}_9$, ^{15}N), valine ($^{13}\text{C}_5$, ^{15}N), Citrulline (d4), GABA ($^{13}\text{C}_4$), glutamine ($^{13}\text{C}_5$), asparagine ($^{13}\text{C}_4$), ornithine (d6), tryptophan (d8), kynurenine (d4)) were obtained from Cambridge Isotope Laboratories (Andover, MA). Stock solutions of each compound were prepared at a concentration of 500 ng/ μL and stored at -80°C . A 10-point calibration curve (0.01-100 pmol/ μL) was prepared by serial dilutions and spiked with internal standards at a final concentration of 5 pmol/ μL . Mass spectrometry grade formic acid was purchased from Sigma-Aldrich (St Louis, MO, USA) and HPLC grade acetonitrile from Fisher Scientific (Fair Lawn, NJ, USA). T

Amino acid extraction

One million cells were extracted with 400 μL 90:10 (v/v) Methanol:water solution containing internal standards at 2.5 nmol/ μL . One tungsten bead were added and the samples were shaken at 30 Hz for 2 min in a mixer mill. After removing the bead, the samples were centrifuged at 4°C , 14000 RPM, for 10 min. The supernatant collected and stored at -80°C until analysis.

Amino acid derivatization with AccQ-Tag

Extracted samples were derivatized by AccQ-Tag™ (Waters, Milford, MA, USA) according to the manufacturer's instructions. Briefly, 10 µL of extract were added to 70 µL of AccQ•Tag Ultra Borate buffer and finally 20 µL of the freshly prepared AccQ•Tag derivatization solution was added, and the sample was immediately vortex for 30s.

Samples were kept at room temperature for 30 minutes followed by 10 minutes at 55°C. For each batch quality control samples and procedure blanks were included.

Calibration curves were prepared in a similar way by adding 10 µL of each point of the curve to 70 µL of AccQ•Tag Ultra Borate buffer and 20 µL of the freshly prepared AccQ•Tag derivatization solution.

Amino acids Quantification by LC-ESI-MSMS

Derivatized solutions were analyzed using a 1290 Infinitely system from Agilent Technologies (Waldbronn, Germany), consisting G4220A binary pump, G1316C thermostated column compartment and G4226A autosampler with G1330B autosampler thermostat coupled to an Agilent 6460 triple quadrupole mass spectrometer equipped with a jet stream electrospray source operating in positive ion mode.

Separation was achieved injecting 1 µL of each sample onto a BEH C₁₈ 2.1x100 mm, 1.7 µm column (Waters, Milford, MA, USA) held at 50°C in a column oven. The gradient eluents used were H₂O 0.1% formic acid (A) and acetonitrile 0.1% formic acid (B) with a flow rate of 500 µL/min. The initial conditions consisted of 0% B, and the following gradient was used with linear increments: 0.54-3.50 minutes (0.1-9.1% B), 3.50-7.0 (9.1-17.0% B), 7.0-8.0 (17.0-19.70% B), 8.0-8.5 (19.7% B), 8.5-9.0 (19.7-21.2% B), 9.0-10.0 (21.2-59.6% B), 10.0-11.0 (59.6-95.0% B), 11.0-11.5 (95.0% B), 11.5-15.0 (0% B). From 13.0 minutes to 14.8 minutes the flow rate was set at 800 µL/min for a faster equilibration of the column.

The MS parameters were optimized for each compound as described in Supporting Information. MRM transitions for the derivatized amino acids were optimized using MassHunter MS Optimizer software (Agilent Technologies Inc., Santa Clara, CA, USA). The fragmentor voltage was set at 380 V, the cell accelerator voltage at 7 V and the collision energies from 14-45V, nitrogen was used as collision gas. Jet-stream gas temperature was 290°C with a gas flow of 11 L/min, sheath gas temperature 325°C, sheath gas flow of 12 L/min. The nebulizer pressure was set to 20 psi and the capillary voltage was set at 4 kV. The QqQ was run in Dynamic MRM Mode with 2 min retention time windows and 500 msec cycle scans. The data was quantified using MassHunter™ Quantitation software B08.00 (Agilent Technologies Inc., Santa Clara, CA, USA) and the amount of each amino acid was calculated based on the calibration curves.

Retention times (rt), MRM-transition stages monitored (precursor ion and product ions) and collision energies of analyzed compounds.

Compounds	MRM transition		rt (min)	Collision Energy (V)
	Precursor Ion	Product Ion		
alanine	260.1	171	4.1	14
arginine	345.1	171	2.8	35
aspartic acid	304.1	171	3.4	18
cystine	581	171	5.2	25
glutamic acid	318.1	171	3.6	22
glycine	246.1	171	3.1	18
histidine	326.1	171	2.6	26
isoleucine	302.1	171	7.8	18
leucine	302.1	171	7.6	18
lysine	487.2	171	5.1	26
methionine	320.1	171	6.0	22
phenylalanine	336.1	171	8.1	18

proline	286.1	171	4.4	14
serine	276.1	171	3.1	14
threonine	290.1	171	3.7	18
tyrosine	352.1	171	5.7	18
valine	288.1	171	6.1	18
citrulline	346.2	171	3.4	30
GABA	274.1	171	4.1	18
glutamine	317.1	171	3.0	22
asparagine	303.1	171	2.8	18
ornithine	473.2	171	4.7	34
tryptophan	375.2	171	8.4	26
kynurenin	379.2	171	7.6	37

Labelled internal standards

alanine (13C3, 15N)	264.07	171	4.1	14
arginine (13C6, 15N4)	355.1	171	2.8	35
aspartic acid (13C4, 15N)	309.07	171	3.3	18
cystine (13C6, 15N2)	589	171	5.2	25
glutamic acid (13C5, 15N)	324.09	171	3.5	22
glycine (13C2, 15N)	249.05	171	3.1	18
histidine (13C6, 15N3)	335	171	2.6	26
isoleucine (13C6, 15N)	309.12	171	7.9	18
leucine (13C6, 15N)	309.12	171	7.6	18
lysine (13C6, 15N2)	495.1	171	5.2	26
methionine (13C5, 15N)	326.17	171	6.0	22
phenylalanine (13C9, 15N)	346	171	8.2	18
proline (13C5, 15N)	292.09	171	4.4	14
serine (13C3, 15N)	280.6	171	3.0	14
threonine (13C4, 15N)	295.08	171	3.7	18
tyrosine (13C9, 15N)	362.12	171	5.7	18
valine (13C5, 15N)	294.1	171	6.1	18
Citrulline (d4)	350.1	171	3.4	30
GABA (13C4)	278.1	171	4.0	18

glutamine (13C5)	322.1	171	3.1	22
asparagine(13C4)	307.1	171	2.8	18
ornithine (d6)	479	171	4.6	34
tryptophan (d8)	383.2	171	8.3	26
kynurenine (d4)	383	171	7.7	37

SAH concentration

Intracellular metabolites were isolated after lysating the cells in modified ripa buffer. Cell lysate was subjected to sonication using bioruptor for 5 minutes (30 second ON/ 30 seconds OFF). Resultant free cell supernatant was snap frozen and stored at -80 °C. Protein concentrations were determined using BCA (Pierce Biotechnology, Rockford, IL) and equal amounts of protein were used for SAH quantification. Quantification of intracellular SAH was done using S-Adenosylmethionine (SAM) and S-Adenosylhomocysteine (SAH) ELISA Combo Kit # MET-5151-C (Cell biolabs, San Diego, CA, USA) as instructed.

Reference List for supplementary materials and methods

1. Waraky A, Lin Y, Warsito D, Haglund F, Aleem E, Larsson O. Nuclear insulin-like growth factor 1 receptor phosphorylates proliferating cell nuclear antigen and rescues stalled replication forks after DNA damage. *J Biol Chem* 2017 Nov 3; **292**(44): 18227-18239.
2. Jacob Y, Voigt P. In Vitro Assays to Measure Histone Methyltransferase Activity Using Different Chromatin Substrates. *Methods Mol Biol* 2018; **1675**: 345-360.
3. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* 2012 Jul; **9**(7): 671-675.
4. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 2014; **15**(12): 550.
5. Corces MR, Trevino AE, Hamilton EG, Greenside PG, Sinnott-Armstrong NA, Vesuna S, *et al.* An improved ATAC-seq protocol reduces background and enables interrogation of frozen tissues. *Nat Methods* 2017 Oct; **14**(10): 959-962.
6. Picelli S, Bjorklund AK, Reinius B, Sagasser S, Winberg G, Sandberg R. Tn5 transposase and tagmentation procedures for massively scaled sequencing projects. *Genome Res* 2014 Dec; **24**(12): 2033-2040.
7. Weichenhan D, Wang Q, Adey A, Wolf S, Shendure J, Eils R, *et al.* Tagmentation-Based Library Preparation for Low DNA Input Whole Genome Bisulfite Sequencing. *Methods Mol Biol* 2018; **1708**: 105-122.
8. Hey J, Paulsen M, Toth R, Weichenhan D, Butz S, Schatterny J, *et al.* Epigenetic reprogramming of airway macrophages promotes polarization and inflammation in muco-obstructive lung disease. *Nat Commun* 2021 Nov 11; **12**(1): 6520.
9. Ross-Innes CS, Stark R, Teschendorff AE, Holmes KA, Ali HR, Dunning MJ, *et al.* Differential oestrogen receptor binding is associated with clinical outcome in breast cancer. *Nature* 2012 Jan 4; **481**(7381): 389-393.
10. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 2010 Jan 1; **26**(1): 139-140.

11. Yu G, Wang LG, He QY. ChIPseeker: an R/Bioconductor package for ChIP peak annotation, comparison and visualization. *Bioinformatics* 2015 Jul 15; **31**(14): 2382-2383.
12. Berest I, Arnold C, Reyes-Palomares A, Palla G, Rasmussen KD, Giles H, *et al.* Quantification of Differential Transcription Factor Activity and Multiomics-Based Classification into Activators and Repressors: diffTF. *Cell Rep* 2019 Dec 3; **29**(10): 3147-3159 e3112.
13. Kulakovskiy IV, Vorontsov IE, Yevshin IS, Sharipov RN, Fedorova AD, Rumynskiy EI, *et al.* HOCOMOCO: towards a complete collection of transcription factor binding models for human and mouse via large-scale ChIP-Seq analysis. *Nucleic Acids Res* 2018 Jan 4; **46**(D1): D252-D259.
14. Carter B, Ku WL, Kang JY, Hu G, Perrie J, Tang Q, *et al.* Mapping histone modifications in low cell number and single cells using antibody-guided chromatin tagmentation (ACT-seq). *Nat Commun* 2019 Aug 20; **10**(1): 3747.
15. Liu CS, Toth R, Bakr A, Goyal A, Islam MS, Breuer K, *et al.* Epigenetic Modulation of Radiation-Induced Diacylglycerol Kinase Alpha Expression Prevents Pro-Fibrotic Fibroblast Response. *Cancers (Basel)* 2021 May 18; **13**(10).
16. Kechin A, Boyarskikh U, Kel A, Filipenko M. cutPrimers: A New Tool for Accurate Cutting of Primers from Reads of Targeted Next Generation Sequencing. *J Comput Biol* 2017 Nov; **24**(11): 1138-1143.
17. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnetjournal* 2011; **17**(1).
18. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nature Methods* 2012; **9**(4): 357-359.
19. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 2009; **25**(16): 2078-2079.
20. Adey A, Shendure J. Ultra-low-input, tagmentation-based whole-genome bisulfite sequencing. *Genome Res* 2012 Jun; **22**(6): 1139-1143.
21. Buenrostro JD, Giresi PG, Zaba LC, Chang HY, Greenleaf WJ. Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position. *Nature Methods* 2013; **10**(12): 1213-1218.

22. Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, *et al.* deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic Acids Research* 2016; **44**(W1): W160-W165.